



RESEARCH ARTICLE

Changes in Nrf2 Expression in the Spinal Cord in Mouse Model of Experimental Autoimmune Encephalomyelitis

Sung Min Nam^{1,§}, Jun Ho Lee^{2,§}, Byung-Joon Chang¹, Jong-Hwan Lee^{1,*} and Hyun-Jin Tae^{3,*}

¹Department of Anatomy, College of Veterinary Medicine, Konkuk University, Seoul 05030, Korea

²Institute of BioInnovation Research, Kolon Life Science, Inc., Magokdong-ro, Gangseo-gu, Seoul 07793, Korea

³Bio-Safety Research Institute, College of Veterinary Medicine, Chonbuk National University, Iksan 54596, Korea

*Corresponding author: hjtae@jbnu.ac.kr; jhlee21@konkuk.ac.kr

ARTICLE HISTORY (17-203)

Received: June 07, 2017
Revised: July 02, 2018
Accepted: October 13, 2018
Published online: December 27, 2018

Key words:

Experimental autoimmune encephalomyelitis
Heme oxygenase-1
Multiple sclerosis
NAD(P)H quinone oxidoreductase 1
Nuclear factor erythroid-derived 2-related factor 2

ABSTRACT

In this study, we investigated the expression of nuclear factor erythroid-derived 2-related factor 2 (Nrf2) in the spinal cord using an experimental autoimmune encephalomyelitis (EAE) model. MOG₃₅₋₅₅ was injected into female mice to induce an autoimmune response. Clinical scoring and a behavioral rotarod test were conducted to assess the successful modeling of EAE. Animals were sacrificed 1, 2, and 3 weeks after immunization. Spinal cords were prepared for morphometry, immunohistochemistry, and western blotting. Multifocal demyelination was observed in the white matter. Nrf2 immuno-reactivity was detected in astrocytes and neurons in spinal cord. Nrf2 immuno-reactive neurons and astrocytes had increased by 2 weeks after immunization, and their up-regulation was maintained until 3 weeks after immunization. Additionally, time-dependent up-regulation of Nrf2 in the spinal cord was investigated by western blotting. The level of NAD(P)H quinone oxidoreductase-1 protein, an Nrf2 target gene, was not changed during week 1, but showed a significant decrease 2 and 3 weeks after immunization. Parallel to up-regulated Nrf2, heme oxygenase-1, another target gene of Nrf2, was significantly augmented until 3 weeks after immunization. We conclude that Nrf2 expression in neurons is involved in an autoimmune attack in the spinal cord in an EAE model.

©2018 PVJ. All rights reserved

To Cite This Article: Nam SM, Lee JH, Chang BJ, Lee JH and Tae HJ, 2018. Changes in Nrf2 expression in the spinal cord in mouse model of experimental autoimmune encephalomyelitis. Pak Vet J. <http://dx.doi.org/10.29261/pakvetj/2018.119>

INTRODUCTION

Multiple sclerosis (MS) is a demyelinating disease developed by an autoimmune attack on the central nervous system, characterized by destruction of myelin and multi-centric inflammation. It leads to serious motor disturbance and clinical disorders in MS range from relapsing-remitting to chronic-progressive form, associated with permanent neurological problems such as motor paralysis and ataxia (Tsujimura *et al.*, 2008; Dendrou *et al.*, 2015; Ransohoff *et al.*, 2015;).

Nuclear factor erythroid-derived 2-related factor 2 (Nrf2) is encoded by the NFE2L3 gene and is a basic leucine-zipper transcription factor that regulates the expression of cytoprotective proteins including antioxidant enzymes (Kraft *et al.*, 2004). Nrf2 has been reported to mediate neuro-protection in Parkinson's

disease and cerebral ischemia (Shih *et al.*, 2005; Chen *et al.*, 2009) and to direct neuroprotection in the autoimmune neuro-inflammatory response in MS (Kraft *et al.*, 2004; Johnson *et al.*, 2010). Recent findings highlight the importance of Nrf2 in MS pathogenesis; in support of this, Nrf2 enhancement has been reported in the brains of various animal models of MS (Solis *et al.*, 2002; Kraft *et al.*, 2004; Johnson *et al.*, 2010; van Horssen *et al.*, 2010). In addition, Nrf2 was shown to be prominently increased in demyelinated lesions in the spinal cord of human MS patients (Licht-Mayer *et al.*, 2015).

However, previous studies using animal models of MS focused predominantly on Nrf2 expression in the brain. Lately, Nrf2 expression and its association with MS in the spinal cord of MS animal models are beginning to be reported (Linker *et al.*, 2011; Higashi *et al.*, 2017). In the present study, we attempted to investigate the expression of Nrf2 in a cellular subpopulation in the spinal cord using an experimental autoimmune encephalomyelitis (EAE) model,

[§]These authors contributed equally to this work.

which serves as an *in vivo* validation model for MS. Additionally, the EAE model in dogs is also a viable model for canine necrotizing encephalitis (Moon *et al.*, 2015). In the spinal cord, Nrf2 expression was studied using immunohistochemistry and western blotting 1, 2 and 3 weeks after immunization.

MATERIALS AND METHODS

Experimental animals and induction of EAE: Nine-week-old female C57BL/6 mice were achieved from the Hallym University (Chuncheon, South Korea). Animals were housed in a conventional state and allowed free access to food and water. The experimental procedures were adhered to the guidelines (NIH Guide for the Care and Use of Laboratory Animals, NIH Publication). EAE was induced according to the protocol of previous study (Lee *et al.*, 2014). Mice (n=14) were subcutaneously immunized with 200µg of myelin oligodendrocyte glycoprotein (MOG₃₅₋₅₅) peptide (AnaSpec; Fremont, CA, USA) which were emulsified in complete Freund's adjuvant (CFA) (H37Ra; Difco Laboratories, Detroit, MI, USA) supplemented with Mycobacterium tuberculosis (4 mg/ml, Sigma-Aldrich; St. Louis, MO, USA). Mice were intraperitoneally injected with pertussis toxin (200ng, Sigma-Aldrich) on day 0 and 2 after immunization. Only CFA was injected in the control mice.

Clinical scoring and evaluation of behavior: Animals were daily checked for clinical scores with the following criteria - 0. healthy; 1. flaccid tail; 2. hind limb mild paralysis; 3. paralysis of both hind legs with residual mobility; 3.5. complete paralysis of both hind legs; 4. complete paralysis of both hind legs with front limb paralysis; 5. moribund or dead (Pöllinger *et al.*, 2009). The motor performance of EAE mice was assessed using rotarod (Med Associates, Inc., St Albans, VT, USA) by checking the latency time to fall (Tsunoda and Fujinami, 2010).

Histology: Mice (n=7) were deeply anesthetized by isoflurane (3% in oxygen), perfused trans-cardially with PBS and 10% neural buffered formaldehyde (NBF). The spinal cord was dissected and post-fixed with 10% NBF. Tissues were paraffin-embedded and sectioned at 6 µm with microtome. First, sections were stained with Luxol Fast Blue (LFB) (Sigma-Aldrich, St. Louis, MO, USA) to assess demyelination.

For immunohistochemistry (IHC) and double immunofluorescence staining, sections were processed according to the previous study (Chang *et al.*, 2012). Briefly, sections were incubated (overnight at 4°C) with rabbit anti-Nrf2 monoclonal antibody (1:200, NOVUS Biologicals, Littleton, CO, USA), and biotinylated-conjugated anti-rabbit antibody, horseradish peroxidase-conjugated streptavidin (1 hr at room temperature). IHC sections were visualized by diaminobenzidine tetrachloride (Vector Lab, Burlingame, CA, USA) and contrasted with hematoxylin counterstain. For double staining, sections were incubated with anti-Nrf2 monoclonal antibody (1:100), mouse anti-gial fibrillary acidic protein (GFAP) (1:200, Chemicon International, Temecula, CA), and mixture of both FITC-conjugated goat anti-mouse IgG (1:200, Jackson ImmunoResearch,

West Grove, PA) and Cy3-conjugated goat anti-rabbit IgG (1:200, Jackson ImmunoResearch). Counterstain was conducted with 4,6-diamidino 2-phenylindole (DAPI; Vector Lab) for nucleus. Images were taken using digital image analyzer (MCID; Inter-Focus Imaging Ltd., Cambridge, UK) and confocal microscope (LSM510 META NLO, Carl Zeiss, Germany). Numbers of Nrf2 immuno-reactivity positive (+) cells were counted in the spinal cord.

Western blotting: Mice (n=7) in each group were sacrificed and spinal cords were dissected and stored at -80°C until analysis. Tissues were processed according to the previous study (Chang *et al.*, 2012). After blotting, membranes were blocked with skim milk and incubated with Nrf-2 antibody and other antibodies (NQO1, HO1, Actin and secondary HRP-conjugated anti-IgG). Immuno-reactive proteins were detected using ECLplus Western Blot Detection Kit (GE Healthcare, Piscataway, NJ, USA). Bands density was quantitated using a density-metric Multi Gauge program (Fujifilm Corporation, Tokyo, Japan).

Statistical analysis: Data was analyzed using ANOVA followed by Duncan's multiple range as *post hoc* comparison test with SAS (SAS Institute Inc., version 9.2, Cary, NC, USA). A value of significant threshold was set as P<0.01. The analysis of motor performance of behavioral test adopted a Kruskal-Wallis nonparametric one-way analysis of variance and followed by a two-tailed Mann-Whitney U test. Statistically, a value of P<0.05 was significant.

RESULTS

Clinical and behavior scores in EAE model: Clinical scores of EAE induction were initially recorded at day 11 after MOG₃₅₋₅₅ immunization, and clinical signs progressed rapidly. In most mice in the experimental group, clinical scores were graded "5" 3 weeks after MOG₃₅₋₅₅ immunization (Fig. 1A). The staying period on the rotarod in the MOG₃₅₋₅₅ immunization group was significantly shorter than that of the control group. The latency time of immunized mice declined significantly beginning 1 week after MOG₃₅₋₅₅ immunization. After 3 weeks, none of the mice in the MOG₃₅₋₅₅ immunization group were able to remain on the rotating drum (Fig. 1B).

LFB staining and measurement of demyelination: Areas of delamination appeared in a multifocal fashion in the white matter of spinal cords of animals in the MOG₃₅₋₅₅ immunization group, while this phenomenon was not observed in the control group (Figs. 2A-F). The delamination areas showed significant time-dependent increases after MOG₃₅₋₅₅ immunization (Fig. 2G).

Nrf2 IHC in spinal cord: In the control group, Nrf2 immunoreactivity was detected in the grey and white matter of the spinal cord. The numbers of observed Nrf2⁺ cells in the spinal cord were similar to the control and 1 week after MOG₃₅₋₅₅ injection. Thereafter, the number of Nrf2⁺ cells in the spinal cord increased significantly compared to the controls 2 and 3 weeks after MOG₃₅₋₅₅ injection. Subsequently, 3 weeks after MOG₃₅₋₅₅ injection, the number of Nrf2⁺ cells in grey matter had decreased compared to that at 2 weeks (Fig. 3).




Fig. 1: Clinical score (A) and behavioral test (B). Clinical score was significantly increased on 11th day, after the induced immunization (MOG₃₅₋₅₅ peptide injection), most of the mice clinical score were "5" on 3rd week (A), results shown the daily mean clinical scores (n=21). Immunized mice latency time (second) was significantly declined after 1 week on rotarod test, the mice could not maintain on rotating drum after 3 weeks (B). Data are expressed as the mean ± S.E.M., and representative of experiments repeated three times.




Fig. 2: Detection of demyelination on the spinal cord with LFB staining (A-F). Demyelination has not been shown by the control group (A; x 100; B and C; x 200). However, MOG₃₅₋₅₅ peptide treatment group's demyelination was significantly increased with time dependently at 3 week (D; x 100; E and F; x 200). Demyelination area in the spinal cord is outlined by the dashed black line. Demyelination data was present with graph (G). Data are expressed as the mean±S.E.M. and representative, ANOVA, P<0.01, Duncan test. Meanwhile, demyelination was not detected in the grey matter of spinal cord.

Double immunofluorescence staining: Double staining was performed to differentiate the type of cells in the spinal cord expressing Nrf2 (Fig. 3). Large numbers of astrocyte in the white matter expressed Nrf2, while some neurons in grey matter showed Nrf2 positivity. We confirmed that, in the spinal cord, the Nrf2 expressing cells in white matter were astrocytes, while in the grey matter, Nrf2 was expressed in neurons (Fig. 3).

Western blot analysis: The results of western blotting revealed a pattern of Nrf2, HO1, and NQO1 protein expression in the spinal cord at 1, 2 and 3 weeks after MOG₃₅₋₅₅ immunization. As EAE progressed, the level of Nrf2 was significantly upregulated compared to the control at 2 and 3 weeks after immunization and peaking at 2 weeks. We also detected time-dependent changes in the Nrf2 target gene HO1, with increases during weeks 2 and 3 after immunization. In contrast, another Nrf2 target

gene, NQO1, remained unchanged 1 week after immunization and significantly decreased 2 and 3 weeks after MOG₃₅₋₅₅ immunization (Fig. 4).

DISCUSSION

This study was designed to investigate the morphometry of demyelination and the expression of Nrf2 in the EAE mouse spinal cord. LFB staining demonstrated multifocal time-dependent demyelination in the white matter of the spinal cord. Nrf2 upregulation was detected in astrocytes in the white matter and in neurons in the grey matter. The numbers of Nrf2⁺ astrocytes and neurons increased 2 weeks after MOG₃₅₋₅₅ immunization, and then decreased at 3 weeks. The protein expression levels of Nrf2 measured by western blotting were similar to the immunohistochemistry results, showing a peak at 2 weeks and a subsequent decrease 3 weeks post-MOG₃₅₋₅₅ immunization.




Fig. 3: Immunohistochemistry of Nrf2 in the spinal cord (A-F) and the positive cell number in gray matter (G) and white matter (H). Control (A and D), 2 weeks (B and E) and 3 weeks (C and F) after MOG₃₅₋₅₅ peptide treatment. Data are expressed as the mean±S.E.M. and representative, ANOVA, $P < 0.01$, Duncan's test. Arrows are marked neurons of Nrf2 immuno-reactivity. A-F: x 200. Double immunofluorescence staining of Nrf2 (red), astrocytes (GFAP, green) and neurons (blue) 3 week after immunizations (I x 100, J x 200). Nrf2 and astrocyte co-location was detected in white matter, and co-location with neuron was detected in grey matter of spinal cord. Nrf2 expressed cell is astrocyte in white matter and is neuron in grey matter of spinal cord. Arrow indicated Nrf2 expressed neurons in the grey matter, arrow head indicated Nrf2 expressed astrocytes in white matter.

Experimentally, MOG₃₅₋₅₅ immunization is a convenient animal model of EAE, which may correspond to human MS (Lee *et al.*, 2014). In this study, clinical grading scores increased 2 weeks after MOG₃₅₋₅₅ immunization, and behavioral disorders were recorded after 1 week. In the spinal cord, demyelinated white matter was observed 1 week after MOG₃₅₋₅₅ immunization, and demyelination progress paralleled the clinical symptoms and behavioral disorders. These phenomena were reported in our previous study and are known to be typical characteristics of MS (Lee *et al.*, 2014).

An inflammatory response and oxidative stress are detected in MS, and Nrf2 has been reported as being closely related to these inflammatory and oxidative stresses in the brain of MS patients (Gilgun-Sherki *et al.*, 2004; Pöllinger *et al.*, 2009; Johnson *et al.*, 2010; Rosenbaum *et al.*, 2012). In our EAE-modeled spinal cords, increased Nrf2 protein levels were again confirmed. In double-stained tissue, the Nrf2⁺ cells were astrocytes in the white matter and neurons in the grey matter. Nrf2⁺ neurons have been reported in the brains of EAE animal model (Linker *et al.*, 2011).




Fig. 4: Expression of Nrf2, NQO-1 and HO1 in the spinal cord. Total levels of Nrf2 and HO1 were increased from week 2 post-immunization of MOG₃₅₋₅₅, however, the level of NQO-1 decreased on 2 and 3 weeks. Results shown were mean±S.E.M and representative of experiments repeated three times. Each group compare with ANOVA, P<0.01. Duncan's test.

Additionally, the activation of Nrf2-ARE pathway was detected in the spinal cord of amyotrophic lateral sclerosis (ALS) model mice (Kraft *et al.*, 2007). Furthermore, Vargas *et al.*, reported that the astrocytes had strong Nrf2 expression and enhanced immune response in spinal cord (Vargas *et al.*, 2005).

The present results confirmed that, within 3 weeks of immunization, expression of Nrf2 is detectable in the neurons and astrocytes in the grey and white matter, respectively, of the spinal cord of EAE mice. Neurons also showed expression of Nrf2 in the grey matter of spinal cord, where the number of Nrf2⁺ cells was augmented at 2 and 3 weeks after immunizations in this study. Recently, Linker *et al.* reported that Nrf2 was expressed in the neurons and astrocytes of the spinal cord 41 and 74 days after EAE modeling, respectively, (Linker *et al.*, 2011). Accordingly, Nrf2 was reported to be expressed in the astrocytes and macrophages of spinal cord lesions in MS patients (van Horsen *et al.*, 2010). However, Nrf2 was also detected in the oligodendrocytes at the initial demyelination sites in MS patients (Licht-Mayer *et al.*, 2015). Interestingly, the number of Nrf2⁺ neurons paralleled the level of Nrf2 protein determined by western analysis, and grey matter of the spinal cord was unaffected by the inflammatory and oxidative attack of MOG₃₅₋₅₅ immunizations. This suggests that Nrf2 expression in neurons is more closely related to the impact on the protein level of Nrf2 in the spinal cord after EAE. Additionally, Johnson *et al.* reported that Nrf2 knockout exacerbated the severity of EAE, indicating that Nrf2 may participate in delaying the development of autoimmune disease (Johnson *et al.*, 2010).

To clearly define the meaning of Nrf2 upregulation in the previous and present studies, we focused on the fact that Nrf2 is an oxidative stress response mechanism initiating the expression of cytoprotective proteins. For instance, targets of Nrf2 include NQO1, glutamate-cysteine ligase catalytic (Gclc), HO1, glutathione S-transferase (GST), UDP-glucuronosyltransferase (UGT), and multi drug resistance-associated proteins (Mrps) (Solis *et al.*, 2002; Maher *et al.*, 2007; Wang and Dore,

2007; Yamamoto *et al.*, 2008; Jarmi and Agarwal, 2009; Zhao *et al.*, 2009).

The protein level of Nrf2 was upregulated in spinal cord tissue in the EAE model of MS and paralleled enhanced expression of the HO1 protein in the present study. Similarly, ginsenoside Rb1-induced protection of neural progenitors from oxidative stresses was mediated by upregulation of Nrf2 and its target HO1 (Ni *et al.*, 2014). The protective effects of HO1 are associated with attenuation of excessive nitric oxide production and cellular toxicity. Several earlier reports revealed HO1 involvement in acute lung injury sepsis, atherosclerosis, hypertension, and kidney injury (Yamamoto *et al.*, 2008; Jarmi and Agarwal, 2009). Our finding that HO1 is induced in the spinal cord in the EAE model indicates concomitant induction of HO1 in the Nrf2-dependent cytoprotective mechanism. NQO1 is another target gene of Nrf2 (Wang *et al.*, 2007) and catalyzes the reduction of quinines while preventing their participation in redox cycling. NQO1 is a broad-spectrum antioxidant that maintains antioxidative alpha-tocopherol and coenzyme Q10 in their active states by scavenging superoxide (Kraft *et al.*, 2004; Wang *et al.*, 2007). During autoimmune demyelination, NQO1 undergoes time-dependent changes different from Nrf2 and HO1 expression. In the present study, the NQO1 level in the spinal cord was significantly decreased 2 and 3 weeks after immunization. However, NQO1 expression was increased in the brain in the EAE animal model (Li *et al.*, 2013). Along with Nrf2-HO1, NQO1 was increased in the brain as a protective mechanism against lead toxicity (Ye *et al.*, 2016). To the contrary, Agúndez *et al.* reported that the NQO1 gene does not show a significant correlation in comparison between MS patients and healthy controls (Agúndez *et al.*, 2014). These inconsistent NQO1 results suggest the possibility of site-dependent differences in NQO1 expression or Nrf2-independent changes in NQO1 expression in the spinal cord. However, clarifying the correlation of NQO1 with Nrf2 in the spinal cord in this model will require additional intensive studies.

Conclusions: The present results indicate the involvement of Nrf2 signaling in regulation of cytoprotective genes in response to inflammatory mediators in MS pathology in spinal cords. Along with the induction of Nrf2, there are multidirectional changes in the expression of potent target genes including HO1 and NQO1. Expressional change of Nrf2 in the spinal cord of the EAE model suggests that future investigations in multiple sclerosis patients focusing on this molecule will open a window to increase understanding of the pathology of this disease.

Acknowledgements: This research was supported by the faculty research fund of Konkuk University in 2017 and by Basic Science Research Program through the National Research Foundation of Korea (NRF), the Ministry of Education (NRF-2017R1D1A1B03031998).

Authors contribution: HJ Tae, SM Nam, JH Lee and JH Lee conceived and conducted experiments, and collected the data. HJ Tae, SM Nam, JH Lee and JH Lee analyzed the data and wrote the manuscript. BJ Chang participated in designing and discussing the study. All authors approved the final version of the manuscript.

REFERENCES

- Agúndez JA, García-Martín E, Martínez C, *et al.*, 2014. NQO1 gene rs1800566 variant is not associated with risk for multiple sclerosis. *BMC Neurol* 14:87.
- Chang BJ, Jang BJ, Son TG, *et al.*, 2012. Ascorbic acid ameliorates oxidative damage induced by maternal low-level lead exposure in the hippocampus of rat pups during gestation and lactation. *Food Chem Toxicol* 50:104-8.
- Chen PC, Vargas MR, Pani AK, *et al.*, 2009. Nrf2-mediated neuroprotection in the MPTP mouse model of Parkinson's disease: Critical role for the astrocyte. *Proc Natl Acad Sci USA* 106:2933-8.
- Dendrou CA, Fugger L and Friese MA, 2015. Immunopathology of multiple sclerosis. *Nat Rev Immunol* 15:545-58.
- Gilgun-Sherki Y, Melamed E and Offen D, 2004. The role of oxidative stress in the pathogenesis of multiple sclerosis: the need for effective antioxidant therapy. *J Neurol* 251:261-8.
- Higashi C, Kawaji A, Tsuda N, *et al.*, 2017. The novel Nrf2 inducer TFM-735 ameliorates experimental autoimmune encephalomyelitis in mice. *Eur J Pharmacol* 802:76-84.
- Jarmi T and Agarwal A, 2009. Heme oxygenase and renal disease. *Curr Hypertens Rep* 11:56-62.
- Johnson DA, Amirahmadi S, Ward C, *et al.*, 2010. The absence of the pro-antioxidant transcription factor Nrf2 exacerbates experimental autoimmune encephalomyelitis. *Toxicol Sci* 114:237-46.
- Kraft AD, Johnson DA and Johnson JA, 2004. Nuclear factor E2-related factor 2-dependent antioxidant response element activation by tert-butylhydroquinone and sulforaphane occurring preferentially in astrocytes conditions neurons against oxidative insult. *J Neurosci* 24:1101-12.
- Kraft AD, Resch JM, Johnson DA, *et al.*, 2007. Activation of the Nrf2-ARE pathway in muscle and spinal cord during ALS-like pathology in mice expressing mutant SOD1. *Exp Neurol* 207:107-17.
- Lee JH, Halperin-Sheinfeld M, Baatar D, *et al.*, 2014. Tellurium compound AS101 ameliorates experimental autoimmune encephalomyelitis by VLA-4 inhibition and suppression of monocyte and T cell infiltration into the CNS. *Neuromolecular Med* 16:292-307.
- Li B, Cui W, Liu J, *et al.*, 2013. Sulforaphane ameliorates the development of experimental autoimmune encephalomyelitis by antagonizing oxidative stress and Th17-related inflammation in mice. *Exp Neurol* 250:239-49.
- Licht-Mayer S, Wimmer I, Traffehn S, *et al.*, 2015. Cell type-specific Nrf2 expression in multiple sclerosis lesions. *Acta Neuropathol* 130:263-77.
- Linker RA, Lee DH, Ryan S, *et al.*, 2011. Fumaric acid esters exert neuroprotective effects in neuroinflammation via activation of the Nrf2 antioxidant pathway. *Brain* 134:678-92.
- Maher JM, Dieter MZ, Aleksunes LM, *et al.*, 2007. Oxidative and electrophilic stress induces multidrug resistance-associated protein transporters via the nuclear factor-E2-related factor-2 transcriptional pathway. *Hepatology* 46:1597-610.
- Moon JH, Jung HW, Lee HC, *et al.*, 2015. A study of experimental autoimmune encephalitis in dogs as a disease model for canine necrotizing encephalitis. *J Vet Sci* 16:203-11.
- Ni N, Liu Q, Ren H, *et al.*, 2014. Ginsenoside Rb1 protects rat neural progenitor cells against oxidative injury. *Molecules* 19:3012-24.
- Pöllinger B, Krishnamoorthy G, Berer K, *et al.*, 2009. Spontaneous relapsing-remitting EAE in the SJL/J mouse: MOG-reactive transgenic T cells recruit endogenous MOG-specific B cells. *J Exp Med* 206:1303-16.
- Ransohoff RM, Hafler DA and Lucchinetti CF, 2015. Multiple sclerosis—a quiet revolution. *Nat Rev Neurol* 11:134-42.
- Rosenbaum S, Ringseis R, Hillen S, *et al.*, 2012. The stress signalling pathway nuclear factor E2-related factor 2 is activated in the liver of sows during lactation. *Acta Vet Scand* 54:59.
- Shih AY, Li P and Murphy TH, 2005. A small-molecule-inducible Nrf2-mediated antioxidant response provides effective prophylaxis against cerebral ischemia in vivo. *J Neurosci* 25:10321-35.
- Solis WA, Dalton TP, Dieter MZ, *et al.*, 2002. Glutamate-cysteine ligase modifier subunit: mouse Gclm gene structure and regulation by agents that cause oxidative stress. *Biochem Pharmacol* 63:1739-54.
- Tsujimura A, Matsuki M, Takao K, *et al.*, 2008. Mice lacking the kf-1 gene exhibit increased anxiety- but not despair-like behavior. *Front Behav Neurosci* 2:4.
- Tsunoda I and Fujinami RS, 2010. Neuropathogenesis of Theiler's murine encephalomyelitis virus infection, an animal model for multiple sclerosis. *J Neuroimmune Pharmacol* 5:355-69.
- van Horsen J, Drexhage JAR, Flor T, *et al.*, 2010. Nrf2 and DJ1 are consistently upregulated in inflammatory multiple sclerosis lesions. *Free Radic Biol Med* 49:1283-9.
- Vargas MR, Pehar M, Cassina P, *et al.*, 2005. Fibroblast growth factor-1 induces heme oxygenase-1 via nuclear factor erythroid 2-related factor 2 (Nrf2) in spinal cord astrocytes. *J Biol Chem* 280:25571-9.
- Wang J and Dore S, 2007. Heme oxygenase-1 exacerbates early brain injury after intracerebral haemorrhage. *Brain* 130:1643-52.
- Yamamoto T, Suzuki T, Kobayashi A, *et al.*, 2008. Physiological significance of reactive cysteine residues of Keap1 in determining Nrf2 activity. *Mol Cell Biol* 28:2758-70.
- Ye F, Li X, Li L, *et al.*, 2016. t-BHQ Provides Protection against Lead Neurotoxicity via Nrf2/HO-1 Pathway. *Oxid Med Cell Longev* 2016:2075915.
- Zhao F, Wu T, Lau A, *et al.*, 2009. Nrf2 promotes neuronal cell differentiation. *Free Radic Biol Med* 47:867-79.