



RESEARCH ARTICLE

Variation in the Genetic Effects of *ABCG2*, *Growth Hormone* and *Growth Hormone Receptor* Gene Polymorphisms on Milk Production Traits in Egyptian Native, Holstein and Hybrid Cattle Populations

AF El-Nahas^{1*}, WM Basiony², S El-Kassas³ and S Mahmoud⁴

¹Department of Animal Husbandry and Animal Wealth Development, Faculty of Veterinary Medicine, Alexandria University, Egypt; ²Department of Animal Wealth Development, Faculty of Veterinary Medicine, Kafrelsheikh University, Egypt; ³Animal, Poultry and Fish Breeding and Production, Department of Animal Wealth Development, Faculty of Veterinary Medicine, Kafrelsheikh University, Egypt; ⁴Department of Physiology, Faculty of Veterinary Medicine, Kafrelsheikh University, Egypt

*Corresponding author: abeer.elnahas@alexu.edu.eg

ARTICLE HISTORY (18-184)

Received: May 22, 2018
Revised: July 31, 2018
Accepted: August 01, 2018
Published online: September 27, 2018

Key words:

Cattle
Marker assisted selection
Milk traits
Polymorphism
RFLPs

ABSTRACT

The aim of the present study was to evaluate the effect of polymorphisms in the ATP-binding cassette subfamily G member 2 (*ABCG2*), *growth hormone (GH)* and *growth hormone receptor (GHR)* genes on 305-day yields and milk traits in native Egyptian cattle (Baladi), Holstein Friesian cattle and hybrid cattle using DNA sequencing and Restriction Fragment Length Polymorphisms (RFLPs). A total of 148 cows were selected from the three animal populations. The *ABCG2-HhaI-AA* genotype was significantly associated with an increased milk lactose percentage and reduced milk yields in Baladi cows. The *GH-MSP1*-heterozygous genotype (GH) showed significant associations with SNF, protein and lactose percentages in the hybrid cows. Fewer TG repeats were found in the 5' untranslated region of *GHR* in Baladi cows, which could be used for marker-assisted selection to protect against the lower milk yields observed in this population. Additionally, *GHR-MSP1-VL*, can be associated with higher SNF, fat, protein and lactose percentages in hybrid cows, as well as lower milk yields in Baladi cows. We concluded that the three cattle populations have their own genetic identities in relation to genetic markers, which can be used for marker-assisted selection to improve milk traits in these three cattle populations.

©2018 PVJ. All rights reserved

To Cite This Article: El-Nahas AF, Basiony WM, El-Kassas S and Mahmoud S, 2018. Variation in the genetic effects of *ABCG2*, *growth hormone* and *growth hormone receptor* gene polymorphisms on milk production traits in egyptian native, holstein and hybrid cattle populations. *Pak Vet J.* <http://dx.doi.org/10.29261/pakvetj/2018.089>

INTRODUCTION

Marked associations between milk yield and composition traits and many genes were recorded in several cattle breeds (Cohen-Zinder *et al.*, 2005; Komisarek *et al.*, 2011; Ibeagha-Awemu *et al.* 2016; Viale *et al.*, 2017). Among these genes, *ATP-binding cassette subfamily G member 2 (ABCG2)*, whose expression is significantly increased in the mammary gland during the lactation period (Farke *et al.*, 2008). *Growth hormone gene (GH)* plays a crucial role in the growth and development of the mammary gland, as well as the initiation and maintenance of milk secretion; the concentration of *GH* in animals controls the calving interval and the milk yield (Wall and McFadden, 2012). *GH* regulate many other genes that control milk

production, such as *insulin-like growth factor (IGF)* (Ge *et al.*, 2003). Furthermore, its action occurs through the activation of the growth hormone receptor (*GHR*), which has been identified as a strong positional and functional candidate gene that influences milk production (Maskur and Arman, 2014; Parihar, 2016). Mutations in the *GHR* gene alter *GH* functions by changing the protein's ability to bind to this receptor on target cells (Maskur and Arman 2014). Several reports documented the association between milk yield and composition in cattle and the polymorphism in *ABCG2* (Mousavizadeh *et al.* 2013; Ghombavani *et al.*, 2016), *GH* (Ge *et al.*, 2003; Kim *et al.*, 2004; Mullen *et al.*, 2010; Komisarek *et al.*, 2011; Mullen *et al.*, 2011), and *GHR* genes (Blott *et al.*, 2003; Maj *et al.*, 2005).

Different genetic variants are present at bovine milk trait loci, and considerable differences in allele

frequencies for these traits were observed among breeds (Cohen-Zinder *et al.*, 2005; Komisarek *et al.*, 2011; Alima *et al.*, 2013; Ibeagha-Awemu *et al.* 2016; Ozdemir *et al.*, 2018). Consequently, marker-assisted selection based on SNPs has been used for the selection of milk production traits (Hayes *et al.*, 2007).

Holstein Friesian, Egyptian Baladi (native) and hybrid cattle, which were established several decades ago, represent major cattle populations in Egypt. Under the use of different management systems, their the milk yield and composition are variable, allowing the production of a variety of milk products (Mousavizadeh *et al.*, 2013). Therefore, the objective of the present study is to examine milk composition traits (SNF, fat, protein, and lactose percentages) in relation to milk yields in Baladi, Holstein Friesian and hybrid cows and determine their potential associations with polymorphisms in the *ABCG2*, *GH* and *GHR* genes, as indicated by DNA sequencing and RFLPs in the three cattle populations.

MATERIALS AND METHODS

Animal resources and sampling: A total of 148 animals were selected from three animal populations in Egypt from different farms, including 48 Holstein Friesian, 68 hybrid and 32 Baladi cows. Blood and milk samples were collected from these animals; blood for DNA isolation, and milk for composition analysis, including the SNF, fat, protein and lactose percentages. The 305-day milk yield was also calculated for each animal based on farm records. All Animals were handled in accordance with the recommendations of the Committee on the Ethics of Animal Experiments of Kafrelsheikh University, Egypt.

DNA isolation, PCR amplification, and DNA sequencing: Genomic DNA was extracted from whole blood using the G-spin™ Total DNA Extraction Kit (Intron Biotechnology, South Korea) according to the manufacturer's protocol. PCR amplification of *ATP-binding cassette subfamily G member 2 (ABCG2)* (Mousavizadeh *et al.*, 2013), *growth hormone (GH)* (Zhou *et al.*, 2005) and the 5' untranslated region of the *growth hormone receptor (GHR)* gene were done. The primer sequences, annealing temperatures and amplified fragment sizes used during the amplification of each gene are listed in Table 1. The PCR reaction contained 12.5 µl of the PCR master mix (Thermo Scientific, Fermentas), 2 µl of the DNA, 0.5 µl of the forward and reverse primers (10 pmol/µl) and deionized water up to 25 µl. The PCR protocol was 95°C for 5 min, followed by 35 cycles of 95°C for 1 min, annealing at the temperature listed for each primer in Table 1 for 1 min and 72°C for 1 min, with a final extension at 72°C for 5 min. The PCR products were purified using the MEGA quick-spin total fragment DNA purification kit (Intron Biotechnology). The purified PCR products from selected samples for each gene were sent to the LGC Company (Germany) for forward DNA sequencing. The sequence data were analyzed using Chromas 1.45. Sequence comparisons were performed using the BLASTN program from the National Center for Biotechnology information website (<http://www.ncbi.nlm.nih.gov/>). The sequences were

aligned using CLUSTAL-W version 1.8 (18). The amino acid translation was achieved using MEGA version 6. The genbank accession numbers used in the comparison of the sequences of the studied genes are listed in Table 2.

Restriction fragment length polymorphisms (RFLPs):

The amplified PCR fragments were digested with the restriction enzymes: *RsaI*, *MSP1*, *HhaI*, *TaqI*, and *HaeIII* (Promega) in a total volume of 25 µl, which was prepared according to the manufacturer's protocol. *HhaI* produced a polymorphic band with *ABCG2*, while *MSP1* produced polymorphic bands in both *GH* and *GHR*. The cleaved fragments were detected via 3% agarose gel electrophoresis. Genotypes were generated for each gene based on the observed restriction enzyme cuts (Table 1).

Statistical Analysis: The phenotypic parameters, means and standard errors were calculated using the PROC MEANS procedure of the SPSS 22 software. Genotypic and allelic frequencies were calculated using the PopGene 32 software. The chi-square test (χ^2) was used to determine whether the populations were in Hardy-Weinberg equilibrium. The association of the identified SNP marker and the studied traits were investigated using the GLM procedures of SPSS version 22.

RESULTS

***ABCG2* gene sequencing, genotyping and its association with milk composition traits:** 100% sequence similarity was reported for the Holstein cows in *ABCG2* gene sequence compared with *B. Taurus*. Two novels repeated SNPs were observed in both Baladi and hybrid cows: 62437C>T and 62445C>G Table 2. The latter SNP was located at the recognition site of the *HhaI* enzyme, resulting in two genotypes (AA and AB), their genotype and allele frequencies are listed in Table 3. However, only one genotype (AA) was present in Holstein cattle. Additionally, two different genotypes were found in the examined cattle populations and were not significantly associated with milk composition traits, except for lactose percentage in the case of the Baladi cattle ($P<0.1$) (Table 4).

***GH* gene sequencing, genotyping and its association with milk composition traits:** The comparisons of *GH* sequences with *B. taurus* indicated the presence of three repeated SNPs in both hybrid and Holstein Friesian cows as shown in Table (2). Additionally, Baladi cows showed 100% similarity. *MspI* produced polymorphic genotypes for the *GH* genes in Baladi, hybrid, and Holstein Friesian cattle. The HH, GG and HG genotypes and the H and G alleles are listed in Table 3. In hybrid cattle, heterozygous genotypes (*GH*) showed a significant association with SNF, protein, and lactose percentages ($P<0.1$) (Table 5). Accordingly, the GG genotypes had lower percentages of SNF, lactose and protein than the GH and HH genotypes. Further, a non-significant association was reported for 305-day milk yields and fat% ($P>0.1$). However, this association was not observed in Holstein Friesian cows, in either homozygous or heterozygous genotypes.

Table 1: Primer sequences, their annealing temperature, PCR product size, restriction enzymes used and their generated genotypes

| Gene | Primer sequence Forward (5'-3') & Reverse (5'-3') | Annealing temperature (°C) | PCR product size | Restriction enzyme | Digestion product size (bp) |
|-------|---|-------------------------------|---------------------|-----------------------|--|
| ABCG2 | 5'- GTATTCACGAGACTGTCAGGG-3' 5'- GGCTTTATTCTGGCTGTTCC-3' | 60 | 240 | HhaI | (A / A) 240 (A / B) 240, 292, 128 (B / B) 292, 128 (G / G) 329 (G / H) 329, 224, 105 (H / H) 224, 105 |
| GH | 5'-CCCACGGGCAAGAATGAGGC-3' 5'-TGAGGAACTGCAGGGGCC | 65 | 329 | MspI | (V / V) 318 (V / L) 318, 292, 26 (L / L) 292, 26 |
| GHR | 5'-CTGGCGTATGGTCTTTGTCA-3' 5'-TGGTCTTGCTGCTTTCCTAT-3' | 55 | 318 | MspI | |

Table 2: SNP and Different Genotypes of ABCG2, GH and GHR genes in Baladi, Holstein and Hybrid cows

| | ABCG2 gene | | |
|---|---|---|--|
| | Baladi | Hybrid | Holstein |
| ABCG2 | 62437C>T | 62437C>T | - |
| Compared with Acc. no AJ871176.1 | 62445C>G | 62445C>G | - |
| Growth Hormone | | | |
| Comparison with gene bank | Compared with <i>Bos indicus</i> GH gene, ID: KT001497.1 100% similarity | Compared with <i>Bos taurus</i> GH, Ins T 1475 1476 G>C 1575 G>A 1576 A>G - | Acc. no JQ711182.1 - - 1575 G>A 1576 A>G 1601 C>T |
| Comparison with each other | 19 C 61 T 62 G 63 G 209 T | 19 G 61 C 62 T 63 C 209 C | 19 G 61 C 62 T 63 C 209 T |
| Growth Hormone Receptor | | | |
| No of TG microsatellite in each population | 10 | 14 | 16 |
| Sequence comparison of the three population | 9 SNPs | No SNPs | No SNPs |
| Compared with gene bank sequences | Compared with <i>Bos indicus</i> GHR ID: AF040955.1 562 (120833) C>G | Compared with <i>Bos taurus</i> GHR, ID: JQ711177.1 120836 G>A 120844 G>A | 120836 G>A 120844 G>A |
| | TG count 10 vs 10 | TG count 14 vs 17 | 16 vs 17 |

Table 3: Genotypic and allelic frequencies of assessed ABCG2, GH and GHR genes in Baladi, Holstein and Hybrid cows

| Breed | ABCG2 | | | | | | GH | | | | GHR | | | | | | | |
|-------------------|-------|------|------|-----|------|------|----|------|------|------|------|------|----|------|------|------|------|------|
| | n | AA | AB | BB | A | B | n | GG | GH | HH | G | H | n | VV | VL | LL | V | L |
| Baladi cows | 32 | 0.70 | 0.25 | 0.0 | 0.88 | 0.12 | 32 | 0.25 | 0.00 | 0.75 | 0.25 | 0.75 | 32 | 0.62 | 0.38 | 0.00 | 0.81 | 0.19 |
| Hybrid Cows | 68 | 0.84 | 0.16 | 0.0 | 0.92 | 0.08 | 68 | 0.05 | 0.16 | 0.79 | 0.16 | 0.84 | 68 | 0.32 | 0.68 | 0.00 | 0.66 | 0.34 |
| Holstein Friesian | 48 | 1.00 | 0.00 | 0.0 | 1.00 | 0.00 | 48 | 0.00 | 0.27 | 0.73 | 0.14 | 0.86 | 48 | 0.73 | 0.27 | 0.00 | 0.86 | 0.14 |

n: number.

GHR gene sequencing, genotyping and its association with milk composition traits: A comparison of the sequences of 5' untranslated region of *GHR* in the three cow populations revealed the presence of 10, 16 and 14 molecular variants of TG repeats in Baladi (*B. indicus*), Holstein (*B. indicus*) and hybrid cows, respectively.

Nine SNPs were detected in the Baladi cattle in comparison with the Holstein Friesian and hybrid cows. The PCR-RFLP-*MspI* analysis revealed two genotypes (VV and VL) in the Holstein Friesian, Egyptian Baladi and hybrid cows. The frequencies of these genotypes varied among the three examined cow populations (Table 3).

In Egyptian Baladi cows, *GHR-MspI* variant genotypes (Table 6) showed a marked association with measured milk composition parameters, including lactose, SNF, fat and protein percentages ($P < 0.1$). The heterozygous (VL) genotype had a higher content of SNF, lactose, fat and protein than the VV genotype. Notably, a non-significant association was found in the case of 305-day milk yields ($P > 0.1$). Additionally, a distinct, strong association with 305-day milk yields was observed in the hybrid cattle population ($P < 0.05$). The VL genotype was

associated with a higher milk yield than the VV genotype. However, a non-significant association was recorded for the milk composition parameters ($P > 0.1$). For Holstein cows, the *GHR* genotypes were not markedly associated with either milk yield or milk composition parameters ($P > 0.1$). Moreover, among these different genotypes, the V allele was predominant in the three studied cattle populations.

DISCUSSION

The potential use of genetic polymorphism, particular polymorphism of *ABCG2*, *GH*, and *GHR* genes in some cattle breeds as functional candidates for production traits have been documented in several reports (Lucy *et al.*, 1998; Zhou *et al.*, 2005; Hadi *et al.*, 2015; Ghombavani *et al.*, 2016). In this study we described the use of these markers with three cattle populations, Egyptian native (Baladi), Holstein Friesian and Hybrid cattle, to find out if these markers can be used with these different cattle population for selection and improvement of milk quality and milk yield.

Table 4: Association of different *ABCG2* genotypes with milk production characteristics in Baladi, Holstein and Hybrid cows

| Breed | genotype | AA | AB | P Value |
|---------------|-----------------------|----------------|-----------------|---------|
| Baladi cows | 305 d milk yield (kg) | 1804.58±229.83 | 1830±215.66 | 0.943 |
| | SNF (%) | 8.96±0.128 | 8.425±0.682 | 0.100 |
| | Fat (%) | 2.83±0.211 | 2.335±0.724 | 0.889 |
| | Protein (%) | 3.36±0.046 | 3.15±0.247 | 0.103 |
| | Lactose (%) | 4.97±0.072 | 4.595±0.378 | 0.09* |
| Hybrid cows | 305 d milk yield (kg) | 2354.22±56.545 | 2033.33±255.855 | 0.574 |
| | SNF (%) | 9.38±0.060 | 9.08±0.110 | 0.604 |
| | Fat (%) | 3.94±0.116 | 3.47±0.821 | 0.583 |
| | Protein (%) | 3.53±0.023 | 3.41±0.032 | 0.629 |
| | Lactose (%) | 5.09±0.032 | 4.95±0.068 | 0.604 |
| Holstein cows | 305 d milk yield (kg) | 8860±105.59 | - | - |
| | SNF (%) | 9.43±0.086 | - | - |
| | Fat (%) | 2.67±0.046 | - | - |
| | Protein (%) | 3.53±0.032 | - | - |
| | Lactose (%) | 5.15±0.047 | - | - |

Table 5: Association of different *GH* genotypes with milk production characteristics in Baladi, Holstein and Hybrid cows

| Breed | genotype | HH | GH | GG | P Value |
|--------------|-----------------------|------------------|-----------------|---------------|-----------|
| Baladi cows | 305 d milk yield (kg) | 1931.938±147.107 | - | 1448.75±323.1 | 0.652 |
| | SNF (%) | 8.86±0.96 | - | 9.15±0.88 | 0.751 |
| | Fat (%) | 3.12±0.296 | - | 3.60±0.156 | 0.118 |
| | Protein (%) | 3.315±0.040 | - | 3.435±0.040 | 0.684 |
| | Lactose (%) | 4.995±0.061 | - | 4.95±0.061 | 0.789 |
| Hybrid cows | 305 d milk yield (kg) | 2167.679±68.916 | 2668.75±32.678 | 2745±67.23 | 0.675 |
| | SNF (%) | 9.447±0.040 | 9.605±0.061 | 6.65±0.123 | 0.006 *** |
| | Fat (%) | 4.204±0.134 | 3.03±0.050 | 2.58±0.031 | 0.131 |
| | Protein (%) | 3.556±0.148 | 3.598±0.023 | 2.47±0.054 | 0.005 *** |
| | Lactose (%) | 5.129±0.229 | 5.238±0.032 | 3.65±0.002 | 0.007 *** |
| Friesian cow | 305 d milk yield (kg) | 8907.5±104.71 | 8733.33±429.080 | - | 0.888 |
| | SNF (%) | 9.76±0.123 | 9.21±0.045 | - | 0.653 |
| | Fat (%) | 2.481±0.085 | 2.36±0.230 | - | 0.128 |
| | Protein (%) | 3.65±0.017 | 3.12±0.072 | - | 0.630 |
| | Lactose (%) | 5.325±0.024 | 5.025±0.099 | - | 0.213 |

Table 6: Association of different *GHR* genotypes with milk production characteristics in Baladi, Holstein and Hybrid cows

| Breed | Genotype | VV | VL | P Value |
|---------------|-----------------------|-----------------|-----------------|---------|
| Baladi cows | 305 d milk yield (kg) | 2226.5±71.889 | 1118.33±58.697 | 0.223 |
| | SNF (%) | 8.77±0.071 | 9.17±0.338 | 0.06* |
| | Fat (%) | 2.804±0.420 | 3.36±0.502 | 0.05* |
| | Protein (%) | 3.295±0.306 | 3.433±0.119 | 0.07* |
| | Lactose (%) | 4.785±0.353 | 5.13±0.193 | 0.05* |
| Hybrid cows | 305 d milk yield (kg) | 2948±126.492 | 2005.962±52.470 | 0.02** |
| | SNF (%) | 9.505±0.077 | 9.254±0.060 | 0.481 |
| | Fat (%) | 4.03±0.407 | 3.8±0.109 | 0.507 |
| | Protein (%) | 3.576±0.326 | 3.476±0.022 | 0.530 |
| | Lactose (%) | 5.163±0.380 | 5.033±0.326 | 0.448 |
| Friesian cows | 305 d milk yield (kg) | 8448.75±225.451 | 9956.67±406.84 | 0.202 |
| | SNF% | 9.856±0.181 | 8.726±1.796 | 0.271 |
| | Fat (%) | 2.71±0.184 | 2.04±1.004 | 0.414 |
| | Protein (%) | 3.68±0.065 | 3.26±0.683 | 0.268 |
| | Lactose (%) | 5.376±0.502 | 4.762±0.969 | 0.273 |

In this study, the identified SNPs in *ABCG2* gene were few compared to those reported on gene bank. Mousavizadeh *et al.* (2013) demonstrated that non-synonymous nucleotide substitution in exon 14, which is associated with fat and protein percentages in Holstein cattle, was observed at only a 2% frequency. Additionally, Tantia *et al.*, (2006) indicated that the presence of fixed alleles of the *ABCG2* gene is responsible for higher milk fat yields and higher fat and protein percentages in Indian cattle (*B. indicus*). Additionally, Ibeagha-Awemu *et al.* (2016) observed low frequency SNPs in genes influencing cow milk traits using high density genome wide genotyping-by-sequencing. The genotypes of the *ABCG2* gene were significantly associated with the milk lactose % ($P<0.1$) in Egyptian Baladi cows (Table 4). Genotype AA

possessed a higher lactose percentage than genotype AB. However, a non-significant relationship was observed for the milk yield, as well as the protein, fat and SNF percentages ($P>0.1$). In addition, a non-significant association with milk yield and content parameters was found in hybrid and Holstein Friesian cattle.

These results are consistent with previous findings of (Ron *et al.*, 2006), they showed that allele A is more frequent than allele B in 23 out of 35 cattle breeds, including both *B. taurus* and *B. indicus* breeds. In addition, the predominance of the A allele is also proposed in the three cattle populations, suggesting that A is an ancestral allele. The significant association reported for lactose percentages in Egyptian Baladi cows indicates that the *ABCG2-HhaI-AA* genotype is associated with

higher lactose and lower milk yields than the AB genotype. This difference may be due to the negative correlation between milk yield (kg) and lactose (Rangel *et al.*, 2017). Thus, the *ABCG2-HhaI-AA* genotype could be incorporated into marker-assisted selection for culling of this genotype due to its association with lower milk yields.

GH-MSP1 polymorphism and its association with milk yield and milk production traits in dairy cattle have been documented in many reports (Wall and McFadden, 2012). The present results are consistent with those of a previous study (Zhou *et al.*, 2005) that reported an association between *GH-MspI*⁺ and more milk protein, as well as an association between heterozygous genotypes and higher milk fat content. Additionally, Falaki *et al.* (1996) found an association between the *GH-MspI* genotype and a high milk fat content. Consequently, the *GH-MSP1* polymorphism might be a good genetic tool for improving milk composition traits in hybrid cows, especially for SNF, protein and lactose percentages. These findings confirm previous findings in other dairy cattle populations (Blott *et al.*, 2003; Maj *et al.*, 2005).

The impact of *GH* on milk yield and composition is controlled by *GHR* which modulate its binding capacity to its target cells (Maskur and Arman, 2014). Molecular variants of TG-repeat and SNPs of 5'-untranslated region of *GHR* gene have the potential to serve as candidate genetic marker for selection in milk production traits (Muhagheh-Dolatbady *et al.*, 2013). Hale *et al.*, (2000) suggested that shorter TG alleles are common in *Bos indicus* cattle whereas longer 16- to 20-TG-repeat alleles predominate in *Bos taurus* breeds. Although we could not compare the amount of milk in three examined animal population, the great differences in their milk production are not mainly due to variation in environmental factors but may be associated with lower number of TG-*GHR* repeats.

Another important marker in our study was *GHR-MSP1* variant genotypes (VL and VV) in examined cattle populations. These genotypes were markedly associated with the variation in the milk composition contents of fat, protein, SNF and lactose % in Baladi cows. Heterozygous genotypes (VL) had higher SNF %, fat %, protein % and lactose % along with higher milk yield than homozygous genotypes (VV). This association possibly due to the short TG-*GHR* microsatellite in case of Baladi cows compared to Holstein Friesian and Hybrid cows. Besides, among these different genotypes, the allele H was reported to be predominant in the three studied cattle population. These findings provide confirmation of the previous findings in other dairy cattle population (Blott *et al.*, 2003; Maj *et al.*, 2005; Li *et al.*, 2013). Therefore, *GHR-MSP1* is tightly associated with the different milk composition characteristics in Baladi cattle.

Conclusions: The three cattle populations have their own genetic identities, which can be used to facilitate marker-assisted selection for each cattle population. The *ABCG2-HhaI-AA* genotype can be used for marker-assisted selection against this genotype due to its association with lower milk yields. The *GH-MSP1* polymorphism might be a good genetic tool for the improvement of milk composition traits in hybrid cows. The presence of fewer

TG-*GHR* repeats can be used in marker-assisted selection against lower milk yields, which is a common characteristic in *B. indicus* breeds. In contrast, a greater number of TG-*GHR* repeats and higher milk yields are obtained with *B. taurus* breeds (Holstein Friesian).

Authors contribution: AF conceived and designed study. WM executed the experiment. S. El analyzed the data. S. revised the manuscript. AF and S. El contributed in the intellectual & scholarly write up of the manuscript. All the authors approved the final version for submission.

REFERENCES

- Alima MA, Xiea Y, Fana Y, *et al.*, 2013. Genetic effects of *ABCG2* polymorphism on milk production traits in the Chinese Holstein cattle MA. *J Appl Anim Res* 41:333-8.
- Blott S, Kim JJ, Moio S, *et al.*, 2003. Molecular dissection of a quantitative trait locus: a phenylalanine-to-tyrosine substitution in the transmembrane domain of the bovine growth hormone receptor is associated with a major effect on milk yield and composition. *Genetics* 163:253-66.
- Cohen-Zinder M, Seroussi E, Larkin *et al.*, 2005. Identification of a missense mutation in the bovine *ABCG2* gene with a major effect on the QTL on chromosome 6 affecting milk yield and composition in Holstein cattle. *Genome Res* 15:936-44.
- Falaki M, Gengler N, Sneyers M, *et al.*, 1996. Relationships of polymorphisms for growth hormone and growth hormone receptor genes with milk production traits for Italian Holstein-Friesian bulls. *J Dairy Sci* 79:1446-53.
- Farke C, Meyer HHD, Bruckmaier RM, *et al.*, 2008. Differential expression of ABC transporters and their regulatory genes during lactation and dry period in bovine mammary tissue. *J Dairy Res* 75:406-14.
- Ge W, Davis M, Hines H, *et al.*, 2003. Association of single nucleotide polymorphisms in the growth hormone and growth hormone receptor genes with blood serum insulin-like growth factor I concentration and growth traits in Angus cattle. *J Anim Sci* 81:641-48.
- Ghobavani MS, Mahyari SA, Rostami, *et al.*, 2016. Effect of polymorphisms in the *ABCG2*, *LEPR* and *SCD1* genes on milk production traits in Holstein cows. *South Afr J Anim Sci* 46:196-203.
- Hale C, Herring W, Shibuya H, *et al.*, 2000. Decreased growth in angus steers with a short TG-microsatellite allele in the PI promoter of the growth hormone receptor gene. *J Anim Sci* 78:2099-104.
- Hadi Z, Atashi H, Dadpasand M, *et al.*, 2015. The relationship between growth hormone polymorphism and growth hormone receptor genes with milk yield and reproductive performance in Holstein dairy cows. *Iran J Vet Res* 16:244-8.
- Ibeagha-Awemu EM, Peters SO, Akwanji KA, *et al.*, 2016. High density genome wide genotyping-by-sequencing and association identifies common and low frequency SNPs, and novel candidate genes influencing cow milk traits *Eveline Scient Rep* 6:31109.
- Hayes B, Chamberlain A, McPartlan H, *et al.*, 2007. Accuracy of marker-assisted selection with single markers and marker haplotypes in cattle. *Genetics Res* 89:215-20.
- Kim NK, Seo YW, Kim GH, *et al.*, 2004. A previously unreported *Dral* polymorphism within the regulatory region of the bovine growth hormone gene and its association with growth traits in Korean Hanwoo cattle. *Anim Genet* 35:152-4.
- Komisarek J, Michalak A and Walendowska A, 2011. The effects of polymorphisms in *DGAT 1*, *GH* and *GHR* genes on reproduction and production traits in Jersey cows. *Anim Sci Papers Rep* 29:29-36.
- Li YD, Ma YN, He PJ, *et al.*, 2013. Effects of Growth Hormone (GH) and Growth Hormone Receptor (GHR) Gene Polymorphisms on Milk Production Traits in Holstein and Holstein x Chinese Yellow Cattle Crosses. *J Anim Vet Adv* 12:1210-16.
- Lucy M, Johnson G, Shibuya H, *et al.*, 1998. Rapid communication: Polymorphic (GT) n microsatellite in the bovine somatotropin gene promoter. *J Anim Sci* 76:2209.
- Maj A, Pareek C, Klauzińska M, *et al.*, 2005. Polymorphism of 5' region of the bovine growth hormone receptor gene. *J Anim Breed Genetics* 122:414-17.

- Maskur R and Arman C, 2014. Association of a novel single nucleotide polymorphism in growth hormone receptor gene with production traits in Bali cattle. *Italian J Anim Sci* 13:3461.
- Mousavizadeh S, Salehi A, Aminafshar M, *et al.*, 2013. Novel SNPs of the ABCG2 gene and their associations with milk production traits in Iranian Holstein bulls. *J Agr Sci Tech* 15:1145-51.
- Muhagheh-Dolatabady M, Habibiza J and Bahreini BM, 2013. Association of TG-repeats in the 5'-flanking region of bovine growth hormone receptor (GHR) gene with milk production traits and somatic cell count in Holstein cattle. *J Livestock Sci Technol* 1:29-34.
- Mullen M, Berry D, Howard D, *et al.*, 2010. Associations between novel single nucleotide polymorphisms in the *Bos taurus* growth hormone gene and performance traits in Holstein-Friesian dairy cattle. *J Dairy Sci* 93:5959-69.
- Mullen M, Lynch C, Waters S, *et al.*, 2011. Single nucleotide polymorphisms in the growth hormone and insulin-like growth factor-1 genes are associated with milk production, body condition score and fertility traits in dairy cows. *Genet Mol Res* 10:1819-30.
- Ozdemir M, Kopuzlu S, Topal M, *et al.*, 2018. Relationships between milk protein polymorphisms and production traits in cattle: a systematic review and meta-analysis. *Arch Anim Breed* 61:197-206.
- Parihar G, 2016. Genetic polymorphism in Receptor (GHR) gene in Sahiwaland Haryana Cattle. Msc Thesis. College of Veterinary Sciences & Animal Husbandry UP India.
- Rangel A, Zaros L, Lima T, *et al.*, 2017. Polymorphism in the Beta Casein Gene and analysis of milk characteristics in Gir and Guzerá dairy cattle. *Genetics Mol Res* GMR16.
- Ron M, Cohen-Zinder M, Peter C, *et al.*, 2006. A polymorphism in ABCG2 in *Bos indicus* and *Bos taurus* cattle breeds. *J Dairy Sci* 89:4921-23.
- Wall E and McFadden T, 2012. Triennial lactation symposium: a local affair: how the mammary gland adapts to changes in milking frequency. *J Anim Sci* 90:1695-707.
- Viale E, Tiezzi F, Maretto F, *et al.*, 2017. Association of candidate gene polymorphisms with milk technological traits, yield, composition, and somatic cell score in Italian Holstein-Friesian sires. *J Dairy Sci* 100:1-11.
- Zhou GL, Jin HG, Liu C, *et al.*, 2005. Association of genetic polymorphism in GH gene with milk production traits in Beijing Holstein cows. *J Biosci* 30:595-8.