



RESEARCH ARTICLE

Protective Effect of Probiotics in Combination with Vaccination on Antibody Response, Biochemical and Hematological Indices in Broilers

Noreen Sarwar ^{1*}, Atif Mehmood², Ali Sheraz² and Muhammad Noman²

¹University of Veterinary and Animal Sciences, Lahore; ²University of Lahore, Lahore, Pakistan

*Corresponding author: noreen.sarwar@uvas.edu.pk

ARTICLE HISTORY (18-473)

Received: December 13, 2018

Revised: December 20, 2018

Accepted: December 26, 2018

Published online: January 28, 2019

Key words:

Antibody response

IBDV vaccine

Multi strain probiotics

NDV vaccine

ABSTRACT

In this study, the effect of multi strain probiotics was studied on antibody response, blood chemistry and hematological parameters in combination with vaccination. A total of sixty day-old broiler chicks were equally divided into four groups: Control, Probiotic, Vaccine and Probiotic + Vaccine. Probiotics were offered at 2g/500ml of water for 5 days/week up to 6 weeks. NDV vaccine was given by eye drops at 0.025ml/bird at day 7. At 21st day NDV Lasota vaccine was offered in drinking distil water at 0.50ml /bird. IBDV vaccine was administered by drinking water at 10th, 22nd and 30th day of experiment. Blood samples were collected at 15, 30 and 45 days to measure antibody response, blood chemistry and hematological parameters. Antibody titer against NDV vaccine and IBD vaccine showed decline in control and probiotic groups. In case of NDV vaccine, there was no significant difference in Vaccine group (GMT=log₂^{7.33}) and Probiotic + Vaccine group (GMT=log₂^{7.66}) at day 45th of experiment. For IBDV vaccine, there was significant difference at 45th day in ELISA titer of vaccine group (2233) and Probiotic+Vaccine group (4206). Cholesterol and ALT showed significant decrease in PV group at 45th day. Significant increase in WBC count was observed in PV group. Level of significance was P≤0.05. Probiotics when administered along with vaccine showed improvement in antibody titer but no harmful increase or decrease in blood chemistry and hematological parameters was observed.

©2018 PVJ. All rights reserved

To Cite This Article: Sarwar N, Mehmood A, Sheraz A and Noman M, 2019. Protective effect of probiotics in combination with vaccination on antibody response, biochemical and hematological indices in broilers. Pak Vet J. <http://dx.doi.org/10.29261/pakvetj/2019.023>

INTRODUCTION

Poultry is the fastest growing sector all over the world. Effective and balanced diet is very important factor for cost effective poultry production. Several hormones and antibiotics are used as growth promoter but due to development of resistance against antibiotics and presence of their residues in feed, probiotics are safe and better choice as feed additives (Bidarkar *et al.*, 2014). Probiotics are emerging as alternatives prophylactics with additional benefits such as enhanced immune response, toxin binding and increased digestion (Pedroso *et al.* 2013, Jadhve *et al.*, 2015).

Probiotics are live microorganisms which when administered in adequate amounts confer a health benefit on the host. These are nontoxic and nonpathogenic microorganisms which when administered through digestive route produce health benefits. Probiotic level also effects on blood parameters, growth performance,

cecal microbiota, carcass traits and immune response of broiler (Pourakbari *et al.*, 2016). Probiotic species belonging to *Streptococcus*, *Enterococcus*, *Candida*, *Lactobacillus*, *Bacillus*, *Aspergillus*, *Saccharomyces* and *Bifidobacterium* have a very good effect on the variety of intestinal micro flora, growth performance, nutrient digestibility, immunomodulation, intestinal histological changes, certain haematobiochemical constrains, carcass characteristics and to improve sensory characters of broiler meat (Kabir, 2009; Pournazari *et al.*, 2017; Souza *et al.*, 2018).

Previously Kudair and Al- Hussary, 2010 observed vaccination against Newcastle disease (ND), Infectious bursal disease (IBD) and Infectious Bronchitis (IB) produces changes in the biochemistry of the serum. Moreover, it has resulted in the change of behavior of several enzymes including alkaline phosphatase (ALP), lactate dehydrogenase (LDH), aspartate aminotransferase (AST) and aminotransferase (ALT) in adding to levels of

glucose, proteins (globulin, albumin), triglycerides, cholesterol and lipids. Oral administration of *Lactobacillus*, *Bifidobacterium* and *Streptococcus* resulted in increase in antibody response against ND vaccine (Talazadeh *et al.*, 2016).

The present study was planned to investigate the combined effect of probiotics and vaccines on antibody response biochemical and hematological indices when offered along with NDV vaccine and IBDV vaccines.

MATERIALS AND METHODS

Experimental design: Sixty, day old broiler chicks were distributed into four groups; Control (C), Probiotic (P), Vaccination (V) and Vaccination + Probiotics (VP). The groups were housed separately, with adequate temperature, lighting, ventilation and feed.

Probiotics including *Bifidobacterium*, *Lactobacillus paracasei*, *Lactobacillus casei*, *Streptococcus thermophilus*, *Lactobacillus acidophilus* were offered at the ratio of 2g/500ml of water for 5 days/week up to 6 weeks. Vaccination against Newcastle disease (ND) and infectious bursal disease (IBDV) was carried out in vaccine group 3. At day 7, a dose of Newcastle disease vaccine (NDV) was given to the broilers by eye drops at the rate of 0.025ml/bird. At day 21 NDV (Lasota) was provided in drinking, distil water at the rate of 0.50ml/bird. Group V was vaccinated with infectious bursal disease (IBDV) (Lohmann Animal Health Heinz-Lohmann-Strasse 4 D-27472 Cuxhaven Cuxhaven ME D-27472 Germany) by drinking water at 10th, 22nd and 30th day. Probiotics were offer to V+P group as offered to P group. VP group was vaccinated at the same rate as V group was. No treatment was given to control group.

Sample collection: Blood samples were collected randomly from brachial wing veins of birds at day 15, 30 and 45. To determine specific antibody titers against NDV, serum samples were subjected to hemagglutination inhibition test (Choi *et al.*, 2013; Khurshid and Rhaman, 2018). IgG in sera against IBDV were detected by enzyme linked immunosorbent assay (Idexx® Bursa Ag capture test kit, Lyon, France) (Yurong *et al.*, 2005; Talebi *et al.*, 2008; Daodu *et al.*, 2018). Serum samples were subjected to biochemical analysis for Albumin, Uric acid, Cholesterol, ALT and AST. Blood samples were processed in hematology analyzer (Sysmex XP-100) for hematological parameters (HGB, HCT, WBC, RBC, MCH, MCHC) 15, 30 and 45 days.

Statistical analysis: All the data was subjected to analysis of variance using SPSS software and least significant design (LSD) test was used to determine the means of different treatments. The level of significance was $P \leq 0.05$ (Talebi *et al.*, 2008).

RESULTS

Antibody titer against NDV vaccine: Mean HI antibody titer of broiler chicks against NDV in control group showed a decline in maternal antibody titer. GMT value at day 15 was $\log_2^{2.66}$ while at day 45 it was $\log_2^{1.33}$. Probiotic group also showed decrease in GMT value from

2.66 at day15 to 1.66 at day 45. The decrease in GMT values in probiotic group was less steep than control group. In vaccinated group, highest GMT value was observed as $\log_2^{7.33}$ at day 45. Probiotic + vaccine group showed relatively increase GMT value i.e. $\log_2^{7.66}$ but statistically there was no significant difference ($P \leq 0.05$) (Fig. 1).

Antibody titer against IBDV vaccine: Maternal antibodies showed decline in control and probiotic groups. In control group ELISA titer was decreased from 553 to 215 at day 15th to day 45th. Probiotic group also showed decrease in GMT values from 580 to 285 at 15th to 45th day. Vaccine group showed the highest ELISA titer values as 2233 at 45th day. Probiotic + Vaccine group showed increase in ELISA titer that was 4207 day 45th. Significant increase in antibody titer was observed as compared to vaccine group ($P \leq 0.05$) (Fig 2).

Effect on Blood chemistry: At 15th and 30th day no significant difference was observed in uric acid, Albumin, AST, ALT and Cholesterol values in C, P, V and PV groups ($P \leq 0.05$). At 45th, uric acid values were significantly different in C, P and V groups while no significant difference was observed in P and PV group. No significant difference was observed in albumin values. ALT and cholesterol showed significant decrease in PV group. AST values were significantly different in V and PV groups only (Table 1).

Effect on hematological indices: At 15th and 30th day no significant difference was observed in WBC, RBC, HB, HCT, MCV, MCH and MCHC in all groups. At 45th day only WBC values showed significant difference while ($P \leq 0.05$). Highest value was observed in PV group followed by V, P and control groups. HB value in control group was significantly different from P, V and PV groups. Highest HB value (7.1) was observed in PV group. HCT values in P, V and PV groups were significantly different from C group but there was no significant difference in experimental groups. MCV, MCH and MCHC values showed no significant difference ($P \leq 0.05$) (Table 2).

DISCUSSION

Probiotics are biotherapeutic agents that enhance growth and immunity (Dhama *et al.*, 2011, Rajput *et al.* 2013). The use of probiotics for poultry is based on the knowledge that the gut flora is involved in resistance to enteric infections including *E. coli*, *Salmonella* and *Campylobacter* (Stern *et al.*, 2001). In present study *Bifidobacterium*, *Lactobacillus paracasei*, *Lactobacillus casei*, *Streptococcus thermophilus*, *Lactobacillus acidophilus* were used as probiotics. *Enterococcus*, *Bacillus*, *E. coli*, *Lactobacillus*, *Streptococcus*, *Bifidobacterium*, *Lactococcus*, *Pediococcus* species, *Saccharomyces boulardii* have been used as a probiotics (Mountzouris *et al.*, 2007, Rajput *et al.*, 2013; Souza *et al.*, 2018). *Lactobacillus acidophilus* and *Bifidobacterium bifidum*, along with *Streptococcus faecalis* showed considerable increase in growth and immunity in the chicken (Hamid *et al.*, 2006)

Table 1: Effect of Probiotics and Vaccines on Blood chemistry of chicks in different groups at different days

Groups	Uric Acid (mg/dl)	Albumin (g/dl)	ALT (IU/L)	AST (I/L)	Cholesterol (mg/dl)
At 15 th day					
C	3.46±0.1 ^a	8.1±0.5 ^a	1.23±0.1 ^a	193±3.1 ^a	136±7.2 ^a
P	3.73±0.7 ^a	8.26±1.8 ^a	1.56±0.4 ^a	188±5.1 ^a	127±11.1 ^a
V	3.8±0.9 ^a	7.06±0.1 ^a	0.13±0.05 ^a	183±16.09 ^a	104±7.54 ^a
PV	4.2±0.5 ^a	7.96±0.2 ^a	1.13±0.5 ^b	186±15.0 ^a	138±25.2 ^a
At 30 th day					
C	5.5±0.6 ^a	9.1±1.0 ^a	0.96±0.0 ^a	187±5.6 ^a	221±2.5 ^a
P	4.7±0.7 ^a ^b	9.06±1.8 ^a	1.76±0.5 ^a	185±6.8 ^a	192±17.5 ^a
V	3.8±0.3 ^b	8±0.8 ^a	1.43±0.4 ^a	182±20.2 ^a	182±37.01 ^a
PV	5.03±0.4 ^a	8.63±0.6 ^a	0.2±0.1 ^a	155±31.7 ^a	174±12.0 ^a
At 45 th day					
C	6.7±0.2 ^b	7.4±0.5 ^a	6.6±0.3 ^b	295±04.0 ^a	228±05.5 ^b
P	5.25±0.1 ^a	9.85±1.7 ^a	6.4±0.5 ^a	293±2.0 ^a	231±8.0 ^a
V	7.03±0.1 ^c	9.7±0.4 ^a	9.1±0.2 ^c	295±2.0 ^b	294±03.78 ^c
PV	3.9±0.6 ^a	9.2±0.1 ^a	2.76±0.1 ^b	137±5.0 ^a	199±1.52 ^b

Values (mean±SD) in each column having different superscript differ significantly ($P \leq 0.05$); C = Control; P= Probiotics; V= Vaccines; PV = Probiotic+ Vaccines.

Table 2: Effect of Probiotics and vaccines on Hematological parameters of chicks in different Groups at different days

Groups	WBC	RBC	HB	HCT	MCV	MCH	MCHC
At 15 th day							
C	223±2.0 ^d	1.98±0.61 ^a	6.0±0.0 ^b	20.4±0.51 ^a	128.1±1.04 ^a	38.5±0.51 ^b	28.6±1.52 ^a
P	256±3.0 ^b	2.17±0.1 ^a	9.93±0.1 ^a	29.3±1.1 ^b	129.9±0.05 ^a	41.46±1.2 ^{ab}	31.9±1.6 ^a
V	249±1.52 ^c	2.46±0.29 ^a	9.0±1.0 ^a	29.3±1.1 ^b	128.1±2.2 ^a	41.8±0.72 ^a	30.8±1.75 ^a
PV	229±1.5 ^a	2.2±0.1 ^a	9.06±0.9 ^a	26.16±0.7 ^a	129.5±3.5 ^a	41.9±1.7 ^a	29.7±2.6 ^a
At 30 th day							
C	126±0.5 ^a	2.63±0.1 ^a	10.2±0.7 ^a	31.05±2.4 ^{ab}	116±1.0 ^a	37±0.5 ^a	32.6±0.4 ^a
P	126.5±0.1 ^a	2.77±0.1 ^a	10.3±0.1 ^a	31.8±0.6 ^a	113±3.0 ^a	37±0.1 ^a	32.5±0.4 ^a
V	127±0.94 ^a	2.63±0.2 ^a	9.3±0.1 ^a	27.5±0.5 ^b	110.5±0.5 ^a	38.2±0.3 ^a	33.7±0.7 ^a
PV	126.5±0.73 ^a	2.55±0.05 ^a	11.2±0.3 ^a	33.4±1.7 ^a	117±2.0 ^a	39.5±1.5 ^a	33±0.7 ^a
At 45 th day							
C	122±1.4 ^a	3.36±0.4 ^c	6.3±0.7 ^c	12.11±0.9 ^b	10.6±0.7 ^b	53.8±0.4 ^a	52.8±1.5 ^a
P	141.3±2.3 ^b	4.8±0.3 ^b	9.83±0.3 ^{ab}	26.9±1.8 ^a	12.4±0.4 ^{ab}	50.7±4.0 ^{ab}	52.9±1.7 ^a
V	156.3±2.5 ^c	5.4±1.0 ^b	8.8±0.1 ^b	24.4±0.4 ^a	13.3±0.4 ^{ab}	45.0±4.0 ^{ab}	52.1±0.8 ^a
PV	171±3.0 ^d	7.1±0.3 ^a	10.8±0.7 ^a	23.86±1.5 ^a	15.7±2.9 ^a	41.67±0.9 ^b	68.1±7.4 ^a

Values (mean±SD) in each column having different superscript differ significantly ($P \leq 0.05$); C = Control; P= Probiotics; V= Vaccines; PV = Probiotic+ Vaccines.

In present study antibody response against NDV vaccine in control group showed a decrease in maternal antibodies. In P group decrease was less steep might be because probiotic reduced the degradation of antibodies. Bioactive peptides formed during lactic acid fermentation could contribute to the immunomodulatory effects of probiotic bacteria (Leblanc *et al.*, 2004) and T cell or B cell mediated immune response can be produced by the interaction between pathogens and host cells (Haghighi *et al.*, 2006). PV group showed increase in titer as compared to V group but difference was not significant. Talebi *et al.*, 2008 reported similar results that probiotics and vaccination when given simultaneously produced better antibody response. The increase in antibody titer in PV group was might be because of immunomodulation of different subsets of immune system cells. The results are against the findings of Talazadeh *et al.*, 2016 that aqua blend probiotics showed significant increase in antibody titer against NDV vaccine.

Antibody titer against IBDV vaccine showed significant increase in Probiotic + Vaccine group as compared to vaccine group. Immunostimulation by probiotics depends upon number and type of probiotics. The findings were in accordance with that of Talebi *et al.*, 2008. Talazadeh *et al.*, 2016 observed significant increase in antibody titer against NDV vaccine but no significant increase in antibody titer against Avian influenza virus vaccine. It might be due to orally administered probiotics can enhance the systemic antibodies in case of some antigens (Haghighi *et al.*, 2005).

Hematological indices like WBC, RBC and HB values showed increase at day 45th of experiment in probiotic+ vaccine group. While HCT, MCHC and MCH did not showed any notable change. HCT did not show notable changes while others showed significant increase in parameters matched with control. Owsibo *et al.*, 2013 observed decrease in lymphocytes, HGB and RBC values after 6 weeks of probiotic treatment.

At 45th day the significant decrease in values of uric acid, ALT, AST and cholesterol was observed in Probiotic+Vaccine group. No significant change was observed in albumin values. The results agreed with Fahimeh work that cholesterol decreased by probiotics and vaccine action in the treatments having no significant effect on albumin in serum and significant decrease effect was observed in uric acid. The highest level of uric acid was observed in control group while low values of uric acid were observed in experimental batch (Hashemzadeh *et al.* 2010). Similarly, Pourakbari *et al.*, 2016 observed decrease in cholesterol and triglyceride level. Owsibo found that cholesterol level reduced significantly and no significant difference was observed in albumin. Owsibo did not observed significant difference in ALT, AST, Uric acid, Albumin.

Our result for cholesterol disagreed with Islam who reported that cholesterol level was increased (Islam *et al.* 2004). The decrease in cholesterol level might be because of hydrolysis of bile and decrease in cholesterol producing enzyme.




Fig. 1: Mean HI antibody titer against NDV vaccine in response to probiotics and/or vaccine in broiler chicks.




Fig. 2: Mean ELISA antibody titer against IBDV vaccine in response to probiotics and/or vaccine in broiler chicks.

Conclusions: It is concluded that as Probiotic+Vaccinated group showed increase in GMT value and showed better immune response and no harmful effect was observed on hematological and biochemical indices so probiotic may be used in feed of broiler. Better combination of probiotic strains might be needed to enhance the immune response and performance of broiler chicks.

Authors contribution: NS conceived the idea, designed the study and was involved in data analysis and write up. AM, AS and MN executed the experiment.

REFERENCES

Bidarkar VK, Swain PS, Ray S *et al.*, 2014. Probiotics: potential alternative to antibiotics in ruminant feeding. *Trends Vet Anim Sci* 1:1-4.

Choi KS, Lee EK, Jeon WJ, *et al.*, 2010. Antigenic and immunogenic investigation of the virulence motif of the Newcastle disease virus fusion protein. *J Vet Sci* 11:205-11.

Daodu OB, Oludairo OO, Aiyedun JO, *et al.*, 2018. Assessment of antibody assay methods in determination of prevalence of

infectious bursal disease among local chickens and guinea fowls in Kwara state, North Central Nigeria. *Veterinary World* 11:1183-7.

Dhama K, *et al.*, 2011. Applications of probiotics in poultry: Enhancing immunity and beneficial effects on production performances and health: A review. *J Immunol Immunopathol* 13:1-19.

Haghighi HR, Gong J, Gyles CL, *et al.*, 2005. Modulation of antibody-mediated immune response by probiotics. *Clin Diag Lab Immunol* 12:1387-92.

Haghighi HR, Gong J, Gyles CL, *et al.*, 2006. Probiotics stimulate production of natural antibodies in chickens. *Clin Vaccine Immunol* 13:975-80.

Hamid RH, Gong J, Gyles CL, *et al.*, 2006. Probiotics stimulate production of natural antibodies in chickens. *Clin Vaccine Immunol* 13:975-80.

Hashemzadeh F, Rahimi S, Torshizi MAK, *et al.*, 2013. Effects of Probiotics and Antibiotic Supplementation on Serum Biochemistry and Intestinal Microflora in Broiler Chicks. *Intl J Agri Crop Sci* 5:2394-8.

Jadhav K, Sharma KS, Katoch S, *et al.*, 2015. Probiotics in poultry broiler feed: A review. *J Anim Nutr Physiol* 1:4-16.

Kabir SML, Rahman MM, Rahman MB, *et al.*, 2004. The dynamic of probiotics on growth performance and immune response in broiler. *Int J Poultry Sci* 3:361-4.

Kabir SM, 2009. The role of probiotics in the poultry industry. *Int J Mol Sci*. 10:3531-46.

Kudair IM and Al-Hussary NAJ, 2010. Effect of vaccination on some biochemical parameters in broiler chickens. *Iraqi J Vet Sci* 24:59-64.

Leblanc J, Fliss I, Matar C, *et al.*, 2004. Induction of a humoral immune response following an Escherichia coli O157:H7 infection with an immunomodulatory peptidic fraction derived from Lactobacillus helveticus-fermented milk. *Clin Diag Lab Immunol* 11:1171-81.

Mountzouris KC, Tsirtsikos P, Kalamara E, *et al.*, 2007. Evaluation of the efficacy of a probiotic containing *Lactobacillus*, *Bifidobacterium*, *Enterococcus*, and *Pediococcus* strains in promoting broiler performance and modulating caecal microflora composition and metabolic activities. *Poult Sci* 86:309-17.

Owosibo AO, Odetola OM, Odunsi OO, *et al.*, 2013. Growth, haematology and serum biochemistry of broilers fed probiotics based diets. *Afric J* 8:5076-81.

Pedroso AA, Hurley-Bacon AL, Zedek AS, *et al.*, 2013. Can probiotics improve the environmental microbiome and resistome of commercial poultry production? *Int J Environ Res Public Health* 10:4534-59.

Pourakbari M, Seidavi A, Asadpour L, 2016. Probiotic level effects on growth performance, carcass traits, blood parameters, cecal microbiota, and immune response of broiler. *Ann Bra Academy Sci* 88:1011-21.

Pournazari M, Qotbi AAA, Seidavi A, Corazzin M. Probiotics, probiotics and thyme (*Thymus vulgaris*) for broilers: performance, carcass traits and blood variables. *Rev Colomb Cienc Pec* 2017; 30:3-10.

Rajput IR, *et al.*, 2013. Effect of *Saccharomyces boulardii* and *Bacillus subtilis* B10 on intestinal ultrastructure modulation and mucosal immunity development mechanism in broiler chickens. *Poult Sci* 92:956-65.

Khurshid S and Rahman IU, 2018. Serological status of newcastle disease virus in live broiler birds of Mansehra Khyberpakhtunkhwa, Pakistan. *Adv Biotech Micro* 11:555810.

Souza LFAD, Araújo DN, Stefani LM *et al.*, 2018. Probiotics on performance, intestinal morphology and carcass characteristics of broiler chickens raised with lower or higher environmental challenge. *Austral J Vet Sci* 50, 35-41.

Stern NJ, Cox NA, Bailey JS, *et al.*, 2001. Comparison of mucosal competitive exclusion and competitive exclusion treatment to reduce *Salmonella* and *Campylobacter* spp. colonization in broiler chickens. *Poult Sci* 80:156-60.

Talazadeh F, Mayahi F and Zeinali, 2016. The effect of Aqua blend Avian probiotic © Including Lactobacillus, Streptococcus and Bifidobacterium on systemic antibody response against Newcastle and Influenza disease vaccine in broiler chicken. *Int J Enteric Pathog* 4(2): e35689.

Talebi A, *et al.*, 2008. Effects of a multi-strain probiotic (Primalac) on performance and antibody responses to Newcastle disease virus and infectious bursal disease virus vaccination in broiler chickens. *Avian Pathol* 37:509-12.

Yurong Y, Ruiping S, Shimin S., 2005. Effect of probiotics on intestinal mucosal immunity and ultrastructure of cecal tonsils of chickens. *Arch. Anim Nutr* 59:237-46.