



SHORT COMMUNICATION

Efficacy of Three Different Peste des Petits Ruminants (PPR) Vaccines and Determination of Appropriate Age of PPR Vaccination in Lambs

Hamid Irshad¹, Aman Ullah¹, Aitezaz Ahsan¹, Aamer Bin Zahur¹, Muhammad Jamil Basra², Muhammad Sajid Azhar² and Muhammad Afzal³ and Muhammad Jahangir¹

¹Animal Health Research Laboratories, Animal Sciences Institute, National Agricultural Research Centre (NARC), 44000, Park Road, Islamabad; ²Livestock Experimental Station Rakh Ghulaman, Bhakkar; ³Food and Agriculture Organization of United Nations, Islamabad, Pakistan

*Corresponding author: hamidirshad@hotmail.com

ARTICLE HISTORY (19-011)

Received: January 08, 2019
Revised: March 08, 2019
Accepted: March 20, 2019
Published online: April 23, 2019

Key words:

Lambs
Pakistan
Peste des petits ruminants
Vaccination

ABSTRACT

The study evaluated the efficacy of three different Peste des petits ruminants (PPR) vaccines and assessed the appropriate age of vaccination in lambs born without maternal antibodies at Livestock Experimental Station, Rakh Ghulaman. The lambs (n=94) were divided into ten groups. The lambs in Group A1 (1-<2 month), A2 (2-3 months) and A3 (>3 months) were vaccinated with PPR vaccine from company A and the lambs in Group B1 (1-<2 month), B2 (2-3 months) and B3 (>3 months) were vaccinated with PPR vaccine from company B. However, PPR vaccine from company C was given to four groups of lambs (Group C1, C2, C3 and C4). Serum samples were collected from experimental animals at monthly intervals for five months. Sera were analyzed using c-ELISA. The results indicated that most appropriate age of vaccination against PPR in lambs without maternal antibodies is 2-3 months with any of three vaccines. However, vaccine C may also be used at the age of < 1 month to get protection against PPR.

©2019 PVJ. All rights reserved

To Cite This Article: Irshad H, Ullah A, Ahsan A, Zahur AB, Basra MJ, Azhar MS, Afzal M and Jahangir M, 2019. Efficacy of three different Peste des petits ruminants (PPR) vaccines and determination of appropriate age of PPR vaccination in lambs. Pak Vet J. <http://dx.doi.org/10.29261/pakvetj/2019.063>

INTRODUCTION

Peste des petits ruminants (PPR) is an acute, highly contagious viral disease of sheep and goats caused by morbillivirus of family paramyxoviridae. It is characterized by high fever, erosive lesions in the mouth, muco-purulent discharges from eyes and mouth, diarrhea, pneumonia and death. Morbidity and mortality rates may vary up to 100% and 90% respectively (Wasee Ullah *et al.*, 2016).

Since first outbreak of PPR in Pakistan (Amjad *et al.*, 1996) many PPR outbreaks have been reported from different parts of the country (Abubakar *et al.*, 2018). During last decade the number of PPR outbreaks has increased at alarming rate affecting sheep and goats of areas where disease has never been reported (Abubakar *et al.*, 2015; Abubakar *et al.*, 2018).

Vaccination is considered an effective control measure against PPR in small ruminants. In Pakistan homologous PPR vaccine (Nigerian strain 75/1) which provides protection for at least three years is available for immunization of small ruminants (Zahur *et al.*, 2014).

High morbidity and mortality due to PPR has been reported in young animals. Studies have shown that maternal antibodies against PPR persist for six months but fell below protection level four months after birth (Bodjo *et al.*, 2006). Previous studies (Bodjo *et al.*, 2006; Balamurugan *et al.*, 2012) estimated the age of PPR vaccination in lambs/kids having maternal antibodies. In Pakistan PPR vaccine coverage in small ruminants is low. The kids and lambs born to unvaccinated animals are without maternal antibodies therefore their susceptibility to disease is very high. A study was therefore, carried out to see the efficacy of three different PPR vaccines and also to assess the appropriate age of PPR vaccination in lambs born without maternal antibodies.

MATERIALS AND METHODS

The study was carried out in lambs of Thalli breed kept at Livestock Experimental Station (LES), Rakh Ghulaman. The ethical approval of the study was obtained from Animal Welfare Committee of the National Agricultural Research Center (NARC), Islamabad. All the

lambs (n=94) were sero-screened for PPR virus (PPRV) antibodies using competitive ELISA (c-ELISA) (ID Screen®PPR Competition, ID.vet, Montpellier, France) before vaccination. The PPR vaccine from three different companies A, B and C were used for the experiment. Each dose of PPR vaccine from company A, B and C contained PPR virus Nigerian strain 75/1 not less than 2.5 TCID₅₀. The names of the companies are kept anonymous to avoid any trade conflict or competing financial conflict of interest. The lambs were divided into 10 groups. The lambs in Group A1, A2 and A3 were vaccinated with PPR vaccine from company A and the lambs in Group B1, B2 and B3 were vaccinated with PPR vaccine from company B. However, PPR vaccine from company C was given to four groups of lambs (Group C1, C2, C3 and C4). All the lambs were given recommended dose (1 ml each) of PPR vaccine through subcutaneous route (Table 1). Blood samples were collected from experimental animals at monthly interval by jugular vein puncture and transported to the Animal health Laboratories, NARC, Islamabad under cold conditions. Sera were harvested from clotted blood samples by centrifugation (Sigma, Germany) at 1500×g for 10 minutes. Serum samples were stored at -20°C till further analysis. The serum samples were analysed for the presence of PPRV haemagglutinin (H) antibodies using c-ELISA with a kit (ID Screen®PPR Competition, ID vet, Montpellier, France) according to manufacturer's instructions. The results of c-ELISA were expressed in terms of percent inhibition (PI). The PI values were calculated using the formula:

$$PI = OD_{\text{sample}} / OD_{\text{NC}}$$

Where OD_{sample} = Optical Density of sample
 OD_{NC} = Optical Density of Negative control

Samples showing PI_≤50 were considered as positive and responsive to PPR vaccine while the sample showing PI_>50 were considered as negative and non-responsive to PPR vaccine.

RESULTS AND DISCUSSION

The sero-screening of lambs before vaccination indicated that 13 lambs were positive for PPRV antibodies. Five of these lambs were present in Group A1, 2 in Group A2, 3 in Group B1 and one each in Groups B2, C1 and C2. The lambs positive for maternal antibodies against PPRV were differentiated into two groups on the basis of age of vaccination for analysis purpose. The PI values of lambs with maternal antibodies against PPRV vaccinated at 1-<2 months of age reduced from 25 to 55 one month after vaccination. These PI values remain negative (PI=82) for PPRV till the end of experiment. The PI values of lambs with maternal antibodies against PPRV vaccinated at 2-3 months of age also became negative (PI=52) after two month of vaccination and remain negative (PI=70) till the end of the experiment (Fig. 1). Maternal immunity plays a significant role in protecting young animals against infectious diseases (Pravieux *et al.*, 2007). However, studies carried out in human and animal models showed negative impact of maternal immunity on vaccine specific humoral response (Rowe *et al.*, 2004).

This study also showed that PI values in the lambs born with maternal antibodies against PPR decreased rapidly when vaccinated against PPR at 1-<2 months or 2-3 months of age making lambs susceptible to PPR. Previous studies have indicated that maternal antibodies against PPR persist for 3-4 months of age (Bodjo *et al.*, 2006) therefore, the suitable age of vaccination for young animals having maternal antibodies is 3-4 months after birth.

The PI values of the lambs in Group A1 (PI values ranges from 72-59) and B1 (PI values ranges from 83-85) remained above 50 even after vaccination indicating that vaccination of lambs at the age of 1-<2 months with vaccine from company A or B did not give protective titres against PPR. In contrast to that PI values of lambs in Group C1 (PI values ranges from 90-33) and C2 (PI values ranges from 79-42) which were vaccinated with vaccine from company C at the age of <1 month and 1-<2 months respectively decreased below 50 two to three months post vaccination. These values remained below 50 till the end of the experiment indicating that vaccine C provided protective titres against PPR. The vaccines used in this experiment from company A, B or C were prepared using Nigerian strain but the type of stabilizers was different in all the vaccines. Studies have indicated that immune response of the vaccine also varies with change of stabilizer and number of viruses in the vaccine (Riyesh *et al.*, 2011). The PI values of the lambs without maternal antibodies vaccinated at the age of 2-3 months and >3 months with vaccine A, B or C decreased below 50 one month post-vaccination and remained below 50 till the end of the experiment (Fig. 1). Vaccine safety and efficacy are biggest challenges in providing protective immunity against infections at early age. Our study indicated that none of three PPR vaccines produced any negative effect on health of lambs even if these vaccines were administered at the age of <1 month. The results also indicated that lambs without maternal antibodies weather vaccinated with vaccine A, B or C gave PI values below 50 if vaccinated at the age of 2-3 months or >3 months. Based on these findings the most appropriate age for vaccination against PPR is 2-3 months. The study also indicated that lambs without maternal antibodies against PPR may also be vaccinated at the age of <1 month with vaccine C to get protection against PPR. However, this experiment was carried out in limited number of lambs of a single breed on single experimental farm therefore further studies are required in large number of animals from different breeds to get better understanding of appropriate age of PPR vaccination in small ruminants of Pakistan.

Acknowledgements: This study was supported by Agricultural Linkages Program, Pakistan Agricultural Research Council (“Development of models for the control of Peste des petits ruminants (PPR) in Pakistan and PPR vaccine using local isolates”) and Food and Agriculture Organization of the United Nations (“Progressive control of Peste des petits ruminants in Pakistan” GCP/PAK/127/USA). The authors are also thankful to Livestock and Dairy Development Department, Punjab and administration of Livestock Experimental Station, Rakh Ghulamman for their assistance.




Fig. 1: Mean percent inhibition (PI) values of Peste des petits ruminants (PPR) c-ELISA in lambs vaccinated with commercially available PPR vaccines from three different companies A, B and C. Each dose of PPR vaccine from company A, B and C contained PPR virus Nigerian strain 75/1 not less than 2.5 TCID₅₀.

Table 1: Lambs (n=94) kept at Livestock Experimental Station (LES), Rakh Ghulaman were vaccinated with Peste des petits (PPR) vaccine from three different companies A, B and C. Each dose of PPR vaccine from company A, B and C contained PPR virus Nigerian strain 75/1 not less than 2.5 TCID₅₀. Serum samples collected from experimental animals were analysed using c-ELISA for presence of PPR virus antibodies

S. No.	Group	PPR vaccine	Age at vaccination	No of lambs
1	A1	A	1-<2 month	9
2	A2		2-3 months	9
3	A3		>3 months	9
4	B1	B	1-<2 month	10
5	B2		2-3 months	10
6	B3		>3 months	10
7	C1	C	<1 month	8
8	C2		1-<2 month	11
9	C3		2-3 months	9
10	C4		>3 months	9

Authors contribution: HI: Planning of study, collection of blood samples and data, analysis of data and writing of manuscript. AU: Collection of blood samples and data. AA: Collection of blood samples, analysis of samples and assistance in write up of manuscript. MJ: Collection of blood samples and analysis of samples. ABZ: Planning of study and analysis of data. MJB: Maintenance of experimental animals and collection of blood samples from experimental animals. MSA: Maintenance of experimental animals and collection of blood samples from experimental animals. MA: Assisted in planning of study and write up of manuscript.

REFERENCES

- Abubakar M, Zahur AB, Naeem K, *et al.*, 2018. Field and molecular epidemiology of peste des petits ruminants in Pakistan. *Pak J Zool* 50:559-66.
- Abubakar M, Irfan M, Manzoor S, *et al.*, 2015. Peste des petits ruminants in Pakistan; past, present and future perspectives. *J Anim Sci Technol* 57:1-8.
- Amjad H, Forsyth M, Barrett T, *et al.*, 1996. Peste des petits ruminants in goats in Pakistan. *Vet Rec* 139:118-9.
- Balamurugan V, Sen A, Venkatesan G, *et al.*, 2012. Study on passive immunity: time of vaccination in kids born to goats vaccinated against Peste des petits ruminants. *Virologia Sin* 27:228-33.
- Bodjo SC, Couacy-Hymann E, Koffi MY, *et al.*, 2006. Assessment of the duration of maternal antibodies specific to the homologous peste des petits ruminant vaccine "Nigeria 75/1" in Djallonké lambs. *Biokemistri* 18:99-103.
- Pravieux J, Poulet H, Charreyre C, *et al.*, 2007. Protection of newborn animals through maternal immunization. *J Comp Path* 137:32-4.
- Riyesh T, Balamurugan V, Sen A, *et al.*, 2011. Evaluation of efficacy of stabilizers on the thermostability of live attenuated thermo-adapted Peste des petits ruminants vaccines. *Virologia Sin* 26:324.
- Rowe J, Poolman J, Macaubas C, *et al.*, 2004. Enhancement of vaccine-specific cellular immunity in infants by passively acquired maternal antibody. *Vaccine* 22:3986-92.
- Wasee Ullah R, Zahur AB, Latif A, *et al.*, 2016. Detection of peste des petits ruminants viral RNA in fecal samples of goats after an outbreak in Punjab Province of Pakistan: a longitudinal study. *Biomed Res Int*:1-8.
- Zahur AB, Irshad H, Ullah A, *et al.*, 2014. Peste des Petits Ruminants vaccine (Nigerian Strain 75/1) confers protection for at least 3 years in sheep and goats. *J Biosci Med* 2:2.