



RESEARCH ARTICLE

Molecular Characterization and Phylogenetic Analysis of Babesia Species Isolated from Domestic Cattle

Rimsha Farooq¹, Mian Abdul Hafeez*¹, Muhammad Oneeb¹, Azhar Rafique³, Kamran Ashraf¹, Faiza Aslam⁴, Nauman Rauf¹, Kiran Khalid¹, Farrukh Bilal¹, Sammina Mahmood⁵ and Adeel Sattar²

¹Department of Parasitology; ²Department of Pharmacology and Toxicology, University of Veterinary and Animal Sciences, Lahore, Pakistan; ³Department of Zoology Government College University, Faisalabad, Pakistan; ⁴Livestock and Dairy Development Department, Punjab, Pakistan; ⁵Department of Botany, Division of Science and Technology, University of Education, Lahore, Pakistan

*Corresponding author: abdul.hafeez@uvas.edu.pk

ARTICLE HISTORY (19-237)

Received: June 15, 2019
Revised: September 25, 2019
Accepted: January 06, 2020
Published online: January 07, 2020

Key words:

Babesia species
Molecular characterization
Phylogenetic analysis
RT-PCR

ABSTRACT

Babesiosis is a key tick born disease caused by the protozoal parasites belonging to the genus Babesia, cosmopolitan in nature and infecting enormous range of large ruminants. Study was intended to determine the presence of local isolates of Babesia species (*B. bovis*, *B. bigemina*) their molecular characterization and Phylogenetic Relationship in cattle (district Narowal) Punjab. Blood samples were collected from 200 suspected cattle, from seven villages of selected district. DNA was extracted by using DNA zole, Gene-all Kit, manual method and subjected to PCR for amplification using specific RLB and Universal Primers. Quantification of protozoan parasite was performed by RT PCR using Solis biodine kit. A total of 36 out of 200 animal samples, were found positive by microscopy. PCR positive samples (17) were amplified and bands of strength 520-bp and 800-bp with specific RLB and for universal primers respectively were obtained for Babesia Species. These 17 samples were sent for sequencing and by using Clustal W; Bioedit software with Neighbor-joining method phylogenetic tree was established. Current study reported the presence of *B. bigemina* i.e. 18% (conventional method) in large ruminants of district Narowal, Pakistan, and confirmation by (molecular characterization) PCR (8.5%) and phylogenetic relationship. Phylogenetic results showed that our study has homology with *Babesia bigemina* strain.

©2020 PVJ. All rights reserved

To Cite This Article: Farooq R, Hafeez MA, Oneeb M, Rafique A, Ashraf K, Aslam F, Rauf N, Khalid K, Bilal F, Mahmood S and Sattar A, 2020. Molecular characterization and phylogenetic analysis of Babesia species isolated from domestic cattle. Pak Vet J. <http://dx.doi.org/10.29261/pakvetj/2020.003>

INTRODUCTION

The major constraints of livestock industry are tick borne diseases. In tick transmitted diseases, the imperative one is the tick fever, caused by intra-erythrocytic parasite of genus Babesia (Apicomplexa: piroplasmadia: babesidae). Tick fever is most ubiquitous disease in its nature. Substantially backlog like reduced reproductive fertility and even deaths observed due to these tick borne diseases in livestock population (Vannier and Krause, 2009). Babesia after Trypanosome is the second most dangerous and prevalent haemo-parasite for mammalian host (Barandika *et al.*, 2006). Babesia species are classified according to their hosts, i.e. Cattle (*B. bovis*, *B. bigemina*, *B. divergens*, *B. major*) Buffalo (*B. bovis* and *B. bigemina*) etc. (Uilenberg, 2006). High temperature and anemia are the common clinical signs of babesiosis (Chisu

et al., 2019). Hemoglobinuria is a typical sign not commonly observed in other protozoan diseases. Other signs include anorexia, yellow mucous membrane, nasal and ocular discharge and reduction in milk yield in lactating animals (Muraleedharan, 2015). About 500 million cattle are at risk throughout the globe (Chaudhry *et al.*, 2010; Oscar and Cristian 2018). Tick borne diseases cause 18.6 billion losses and deficiency of 3 billion pieces of hide in cattle per year (Terkawi *et al.*, 2012). Economic losses of production by ticks are about 13-19 billion per year globally (De Castro and Newson, 1993; Oscar and Cristian 2018). Diminazine aceturate and imidocorb dipropionate are used in the treatment of Babesiosis (Mosqueda *et al.*, 2012).

Focus on the mitochondrial Cytochrome c Oxidase Subunit I (COXI) gene in the present study was due to its nature of being the most conservative protein-coding

genes in the mitochondrial genome of animals. COXI exhibit fast mutation rate and easily distinguished between closely related species (Hong *et al.*, 2019), preferred for the evolutionary time depths. Internal Transcribed Spacer primers reveal highest degree of variation along its genomic sequences via unequal crossing over and gene conversion (Eui *et al.*, 2016).

The molecular characterization and phylogenetic analysis was opted as such for the confirmation of parasites as by (Oliveira-Sequeira *et al.*, 2005; Rajabi *et al.*, 2017; Barbara *et al.*, 2017; Jafarbeklo *et al.*, 2018; Nasr *et al.*, 2018; Pardeep *et al.*, 2019). For prevention and immunization of calves, vaccines are used mostly imported ones. It is imperative to get idea about the prevalent strain and their genetic evolutionary relationship to be taken as candidate vaccinal antigen (local isolates). The purpose of present study is to know the genetic makeup of prevalent strains of Babesia with parasite of other areas to estimate the extent of disease caused by different strains.

MATERIALS AND METHODS

Sampling and identification: A total number of 200 blood samples were randomly collected from suspected cattle in district of Narowal. Sampling was performed on the basis of number of animals present in each station. Blood was collected from jugular vein into EDTA-coated vacutainers and transported. Staining was performed by thin and thick blood smear.

DNA Extraction and PCR: The preliminary screened samples were subjected to DNA extraction. For comparative study DNA extraction was performed by three different methods (DNAzol BD, Kit & Manual method) as per manufacturer's instructions.

Kit method: GeneAll Exgene kit was used for the purification of total DNA from blood samples. DNA was extracted by using kit by following the directions.

Manual method: DNA was extracted by manual method by using lysis buffer, TEN buffer and PCI buffer. The concentration of DNA was estimated by using the Nano Drop (Mtshali *et al.*, 2013).

Two sets of primers were selected for the amplification of DNA sample. One set of primers 18S ribosomal DNA specific for *Babesia* species termed as RLB primers of product size 520-bp was used for amplification. Second set of primers was universal (COXI) i.e. 1202 and 400 primers which amplified a product of 800-bp size (Table 1).

PCR and conditions: To confirm the *B. bigemina* and *B. bovis* PCR was executed by using 25- μ l reaction mixture, having 2 μ l of extracted DNA, 2.5 μ M of each primer and 15 μ l of Dream Taq Green PCR Master Mix. The PCR reactions were performed in an automated Thermal Cycler (Bio Rad, USA). Each cycle consisted of denaturation @ 94°C for 1 min., an annealing step of 1 min. at 55°C (with a bit modification) followed by 40 cycles and extension @ 72°C for 1.5 min. Preceding's the final extension step at 72°C for 10 min. (Gubbles *et al.*, 1999).

The condition for universal primer was initial denaturation consist of 96°C for 10 minute followed by 94°C for 30 second annealing at 48°C for 30 second followed by 35 cycles extension at 72°C for 90 second and final extension at 72°C for 10 minute. Agarose gel 1.5% stained with Biotium GelRed Acid Stain was used to analyze the PCR-generated amplicons by electrophoresis. The gel electrophoresis was operated at 113 Volts for 35 minutes. DNA Ladder was also used as Molecular weight marker (Mtshali *et al.*, 2013). A 100 kb molecular weight marker was run along with samples.

Sequencing: The confirmed PCR product was sent to laboratory Advance Bioscience International First base Malaysia for sequencing. Nucleotide sequences were calibrated and arranged by using Bioedit software. The consensus sequences which were obtained by using Bioedit software, was trimmed manually to the equivalent length for the construction of phylogenetic tree. The other software Mega-7 and neighbor-joining methods was used for the construction of phylogenetic tree (Criado-Fornelio *et al.*, 2003). The Basic Local Alignment of BLAST Search Tool was used to check the sequence similarity between our data and in database. Sequence homology helped to identify the putative gene. Sequences were checked on NCBI website by using URL Genbank. <http://www.ncbi.nlm.nih.gov/nucleotide>.

RT-PCR: The Solis biodine kit was used for Real time PCR. Each cycle consisted of denaturation @ 94°C for 1 min., an annealing step of 1 min. at 55°C (with a bit modification) followed by 40 cycles and extension @ 72°C for 1.5 min. Preceding's the final extension step at 72°C for 10 min. The results for quantification of parasites were recorded in terms CT values.

Statistical analysis: The data pertaining prevalence was imperiled to chi-square test and univariate analysis for analyses using Statistical Package for Social Sciences (SPSS) version 20. P-value less than 0.05 were considered significant.

RESULTS

Samples of cattle (200 in number) were processed for the identification of Babesia under microscope after staining procedure. Thin blood smear was stained with Giemsa and observed under oil immersion lens of microscope i.e. 40x. Intra erythrocytic bodies were found in heavy concentration with Giemsa Stain. Blood sample was also stained with field stain B and intra-erythrocytic bodies were observed. These intra-erythrocytic bodies were found inside the red blood cells and these were horse shoe shape.

Out of 200 samples 36 were found positive i.e. they have intra erythrocytic bodies. Percentage of infection was 18% by microscopic examination. The PCR detected 17 samples positive (Table 2). PCR reactions targeting the 18S ribosomal DNA (RLB-Internal Transcribed Spacer Primers) of Babesia *bigemina* species was tested for its ability to amplify the target Babesia species with 100bp ladder. Exact 520 bp amplified product was retrieved. The amplified DNA used (Fig. 1) in PCR was of *Babesia*.




Fig. 1: PCR reactions targeting the RLB Primers of *Babesia bigemina* species that infect cattle & buffalo. Primer pair was tested for its ability to amplify the target *Babesia* species with 100bp ladder in lane #1. Lane#2 Positive Control, Lane #3 Negative Control, Lane #4, 5 & 7 Samples. Exact 520 bp amplified product was found.




Fig. 2: PCR reactions targeting the Mitochondrial Cytochrome c Oxidase subunit I (COI) loci of *Babesia bigemina* species that infect cattle & buffalo. Primer pair was tested for its ability to amplify the target *Babesia* species with 100bp ladder in Lane #1. Lane # 2, 3, 4 & 5 samples. Exact 800 bp amplified product was instituted.

Babesia sp. WA1 isolate CA5 18S ribosomal RNA gene, partial sequence
Sequence ID: [AY027815.1](#) Length: 1742 Number of Matches: 1

Score	Expect	Identities	Gaps	Strand
520 bits(281)	2e-143	373/415(90%)	16/415(3%)	Plus/Plus
Query 1	GGTGCCAGCAGCCGCGTAATTCAGCTCCAATAGCGTATATTAATAATGTTGCAGTTAA	60		
Sbjct 551	GGTGCCAGCAGCCGCGTAATTCAGCTCCAATAGCGTATATTAATAATGTTGCAGTTAA	610		
Query 61	AAAGCTCGTAGTTGAATTTCTGCTGCAATTG-C-TT-GTGTCCCTCTGGGGTCT-G-T-GC	114		
Sbjct 611	AAAGCTCGTAGTTGAATTTCTGCTGCAATTG-C-TT-GTGTCCCTCTGGGGTCT-G-T-GC	669		
Query 115	AT-GTGGCTTTTTTCGGACGG-AGTT-T-C-TTTGTCTGAATGTTTACTTTGAGAAAAAT	169		
Sbjct 670	CTGGTGGCTTACCCTGCGCGGTGTTCTCCATTTG-C-CAGT-TTTACTTTGAGAAAAAT	726		
Query 170	AGAGTGCTCAAAGCAGGCTTTTCGCCTTGAATAGTTAGCATGGAAATAAATAAGTAGGACT	229		
Sbjct 727	AGAGTGTTCAAGCAGGCTTTTCGCCTTGAATAGTTAGCATGGAAATAAATAAGTAGGACT	786		
Query 230	TTGGTTCTATTTTGTGGTTTAGGTACCAAAGTAATGGTTAATAGGAACAGTTGGGGGC	289		
Sbjct 787	TTGGTTCTATTTTGTGGTTTAGGTACCAAAGTAATGGTTAATAGGAACAGTTGGGGGC	845		
Query 290	ATTCTGATTTAACTGTCAGAGGTGAAATTCCTAGATTTGTTAAAGACGAACTACTGCGAA	349		
Sbjct 846	ATTCTGATTTAACTGTCAGAGGTGAAATTCCTAGATTTGTTAAAGACGAACTACTGCGAA	905		
Query 350	AGCATTTGCCAAGGATGTTTTTCATTAATCAAGAACAAAAGTTAGGGGATCGAAGA	404		
Sbjct 906	AGCATTTGCCAAGGATGTTTTTCATTAATCAAGAACAAAAGTTAGGGGATCGAAGA	960		

Fig. 3: Partial Sequence of *Babesia* Species WA-1 isolate from Narowal (18-S Ribosomal RNA gene) Partial sequences of *Babesia* species WA1 isolates from Narowal.




Fig. 4: Phylogenetic analyses sequences obtained from PCR positive sample. The Phylogenetic Tree reflecting the exact position of *Babesia bigemina* present the District Narowal, Pakistan.

Table 1: Two sets of primers were selected for the amplification of DNA sample. One set of primers 18s ribosomal DNA specific for *Babesia* species (RLB primers). Second set of primers was universal (COXI) i.e. 1202 and 400

Specie	Primer name	Primer sequence	Target region	Product size
Babesia	RLB(Forward)	5'- GAGGTAGTGACAAGAAATAACAATA-3'	18S ribosomal	520bp
	RLB(Reverse)	5'-TCTTCGATCCCCCTAACTTTC-3'		
Babesia	400(Forward)	5'-GGDTCAGGTRTTGGTTGGAC-3'	COXI specific	800bp
	1202(Reverse)	5'-CCAAKRAYHGACCAAGAGATA-3'		

(Ogedengbe *et al.*, 2011; (Gubbels *et al.*, 1999).




Fig. 5: Graphical representation of Real time PCR for *Babesia bigemina* as of X-axis number of cycles and on Y-axis Fluorescence. The Ct value for *Babesia bigemina* samples showed fluorescence of sample R (purple) 22.6 and sample RM (yellow) 22.14.

Table 2: Percentage of Positive and Neagtive Samples by Micrometry & Molecular characterization of *Babesia bigemina* (Chi square)

	Microscopy	PCR
Positive	36	17
Negative	164	183
Total	200	200

P value=9.13E-060.000009 (Chi square was used to get the percentage).

Table 3: Estimates of net base composition bias disparity between sequences of babesia species prevalent in narowal and other areas of Pakistan

	SS2	SS3	SS4	SS7	Ikhlash_ Darman Bara- pur.	Bara- mangh
SS2_RLBP						
SS3_RLBP	0.000					
SS4_RLBP	0.000	0.000				
SS7_RLBP	0.000	0.000	0.000			
Ikhlash_pur_Pak.	0.000	0.000	0.000	0.000		
Baramangh_Pak.	0.000	0.000	0.000	0.000	0.080	
Shakargarh_Pak.	0.000	0.000	0.000	0.000	0.000	0.028
Noorkot_Pak.	0.811	0.811	1.255	0.722	0.448	0.189
Ikhlash_Poor_Pak.	0.000	0.000	0.038	0.000	0.000	0.009
Bhajina_Pak.	0.000	0.000	0.000	0.000	0.000	0.009
Shakargarh_Pak.	4.692	4.692	5.175	4.821	0.000	5.259
Narowal	0.425	0.425	0.608	0.325	2.637	0.368
SS5-RLPB	0.000	0.000	0.000	0.000	0.000	0.000

The Mitochondrial Cytochrome c Oxidase subunit I (COI) loci with universal Primers (1202 Reverse and 400 Forward) of *Babesia bigemina* species that infect cattle was experienced to amplify the target *Babesia* species. Precise 800 bp amplified product was obtained (Fig. 2).

Sequencing and phylogenetic: The tree (Fig. 4) showed a similar ancestor relationship of *Babesia* species with other areas as found in district Narowal. The sequences of the present study reflected to be distributed in same cluster of *Babesia bigemina* as well as same clade. But the constructed phylogenetic tree revealed that one sequence (*B. bigemina* Nar/Pak-8 Fig. 3) clustered in a different sub clade, denoted to be genetically different from other isolates collected from Narowal.

Disparity index per sites in comparison with area of present study for all sequences is presented (Table 3). The Zero value reflected the no difference while the values greater than zero indicate the higher difference in base composition. Evolutionary analyses were conducted in MsEGA7.

RT-PCR: The Ct value for *Babesia bigemina* was calculated by real time PCR and the samples showed fluorescence of sample R (purple) 22.6 and sample RM (yellow) 22.14 (Fig. 5). The NTC was used as threshold and two samples were amplified; named as R and Rm. AT cycle no. 20 the reaction shows fluorescence. This shows that it starts amplification at cycle no. 20 and the maximum shows at ct value 22.55 and 22.14.

DISCUSSION

Tick borne diseases impose considerable constraints on cattle and buffalo health and economic development in temperate areas of the world (Barandika *et al.*, 2006). Annually 500 million dollars economic losses are observed because of tick fever and most losses occur due to *B. bovis* infection (Ramos *et al.*, 2012). These samples were screened for the phylogenetic analysis of *B. bovis* and *B. bigemina*. Although the distribution of *B. bovis* and *B. bigemina* was found all over the country but a notable observation was made during the screening of blood samples of district Narowal i.e. the distribution of *B. bigemina* was more than 18% i.e. in 200 suspected blood samples 36 were found as positive by microscopy which is in coherence with a previous study in which prevalence rate of *B. bigemina* was 20.66%. The high prevalence rate was found in low age group and in females. (Saad *et al.*, 2015; Khan *et al.*, 2016)

Another study showed a high prevalence rate of *B. bigemina* in which 24 samples were found positive out of 100 for *B. bovis* and *B. bigemina* by microscopy (Saad-Roy *et al.*, 2015) which is in accordance with present study. The prevalence rate of *B. bigemina* around 19% and of *B. bovis* was 11% by Ahmad *et al.*, 2014 is also in agreement with current findings. The prevalence rate in calves less than one year was higher than aged calves. As for as sex wise prevalence was concerned, female calves were more prone to *Babesia* than male calves (Ahmad *et al.*, 2014; Abdelrasol *et al.*, 2017).

Out of 36 microscopy samples only 17 samples got amplified. PCR reactions targeting the 18S ribosomal DNA of *Babesia bigemina* species was tested and exact 520 bp amplified product was retrieved which is in coherence with Gubbels *et al.*, 1999.

The Mitochondrial Cytochrome c Oxidase subunit I (COXI) loci with universal Primers of *Babesia bigemina* species was experienced and 800 bp amplified product was instituted as that in case of Ogedengbe *et al.*, 2011; Hong *et al.*, 2019.

These were sent for sequencing and by using Bioedit software extract the sequencing files. Mega 7 is software by align the sequences and Clustal W consensus sequences were formed. Jafarbeklo *et al.*, 2018; Nasr *et al.*, 2018; Pradeep *et al.*, 2019. Sequences confirmed *B. bigemina* by using Neighbour-joining, Maximum likelihood method and Boot stripping method a phylogenetic tree was constructed.

The sequences of the present study reflected to be distributed in same cluster of *Babesia bigemina* as well as same clade. But the constructed phylogenetic tree revealed that one sequence (*B. bigemina* Nar/Pak-8) clustered in a different sub clade, denoted to be genetically different from other isolates collected from Narowal. This tree was as similar as constructed within a study that conducted in South Africa in 2013 (Mtshali *et al.*, 2013; Hajeel and Al-Fatlawi *et al.*, 2019). The tree is drawn to scale, with branch lengths measured in the number of substitutions per site. The analysis involved 15 nucleotide sequences (Mtshali *et al.*, 2013). All positions containing gaps and missing data were eliminated. There was a total of 212 positions in the final dataset. Evolutionary analyses were conducted in MEGA7. This tree showed a similar ancestor relationship that found in district Narowal. This study was planned in first time in Narowal district and their evolutionary relationship in form of phylogenetic tree was constructed first time in Narowal.

In present study two samples of Babesial DNA was operated at same condition at which conventional PCR gave results. Same RLB primers were used. The Ct value at which it started to produce fluorescence was 22.41 and 22.31 respectively. The results were in comparison with the study carried out in 2007 by Chulmin Kim in National Research Centre for Protozoan disease and showed fluorescence at same Ct value (Kim *et al.*, 2007; Barbara *et al.*, 2017).

Conclusions: The presence of *B. bigemina* i.e. 18% (conventional method) in large ruminants of district Narowal, Pakistan, and their molecular characterization by PCR (8.5%) and phylogenetic relationship. Phylogenetic results showed that our study has homology with *Babesia bigemina* strain.

Authors contribution: Conceived and designed the experiments: RF, MA Hafeez. Performed the experiments: RF, KK, FA, AR, SM & FB: Analyzed the data: KA, MO & NR: Contributed reagents/materials/ analysis tools: Wrote the paper: MAH & RF.

REFERENCES

- Abdel R, Saeed KH, Salih MTH, *et al.*, 2017. Seroprevalence and molecular detection of babesia species in cattle in Baghdad city. *Bas J Vet Res* 16:54-64.
- Ahmad I, Khwaja A, Shams S, *et al.*, 2014. Detection of babesiosis and identification of associated ticks in cattle. *Inter J Bioassays* 3:3195-9.
- Barbara A, Qurollo, Nikole RA, *et al.*, 2017. Improved molecular detection of babesia infections in animals using a novel quantitative real-time PCR diagnostic assay targeting mitochondrial DNA. *Parasit Vect* 10: 128.
- Barandika J, Berriatua E, Barral M, *et al.*, 2006. Risk factors associated with ixodid tick species distributions in the Basque region in Spain. *Med Vet Ento* 20:177-88.
- Chaudhry Z, Suleman M, Younus M, *et al.*, 2010. Molecular detection of *Babesia bigemina* and *Babesia bovis* in crossbred carrier cattle through PCR. *Pak J Zool* 42:201-4.
- Chisu V, Alberti A, Zobba R, *et al.*, 2019. Molecular characterization and phylogenetic analysis of *Babesia* and *Theileria* spp. in ticks from domestic and wild hosts in Sardinia. *Acta Trop* 196:60-65.
- Criado-Fornelio A, Martinez-Marcos A, Buling-Sarana A, *et al.*, 2003. Molecular studies on *Babesia*, *Theileria* and Hepatozoon in southern Europe: Part II. Phylogenetic analysis and evolutionary history. *Vet Parasitol* 114:173-94.
- De Castro J and Newson R, 1993. Host resistance in cattle tick control. *Parasitol Today*. 9: 13-17.
- Eui JD and Paek S, *et al.*, 2016. Application of Partial internal transcribed spacer sequences for the discrimination of *Artemisia capillaris* from Other artemisia species. *Evid-Based Compl Alt Med* <http://dx.doi.org/10.1155/2016/7043436>
- Gubbels J, De Vos A, Van der Weide M, *et al.*, 1999. Simultaneous Detection of Bovine *Theileria* and *Babesia* Species by Reverse Line Blot Hybridization. *J Clin Micro* 37:1782-9.
- Hajeel MS and Al-Fatlawi MAA, 2019. Sequencing-based phylogenetic study of *Babesia* spp. detected in tick tissues in Al-Diwaniyah province, Iraq. *Iraqi J Vet Sci* 33:9-12.
- Hong SE and Kim SY, *et al.*, 2018. Detection and characterization of an emerging type of *Babesia* sp. similar to *Babesia motasi* for the first case of human babesiosis and ticks in Korea. *Emerg Microb Infect* 8: 869-78.
- Jafarbekloo A, Ramzougouyan MR, Shirian S, *et al.*, 2018. Molecular Characterization and Phylogenetic Analysis of *Theileria* spp. and *Babesia* spp. Isolated from various ticks in Southeastern and Northwestern Regions of Iran. *Vect Borne and Zoon Dis* 18:595-600.
- Khan A, Abid Ur Rehman MH, Khan AZ, *et al.*, 2016. Burden of babesiosis among domestic cattle of southern Khyber Pakhtunkhwa, Pakistan. *Pak J Entomol Zool Stud* 4:305-7.
- Kim C, Iseki H, Herbas MS, *et al.*, 2007. Development of TaqMan-based real-time PCR assays for diagnostic detection of *Babesia bovis* and *Babesia bigemina*. *American J Tropical Med Hyg* 77:837-41.
- Mtshali MS and Mtshali PS, 2013. Molecular diagnosis and phylogenetic analysis of *Babesia bigemina* and *Babesia bovis* hemoparasites from cattle in South Africa. *BMC Vet Res* 9:154-60.
- Mosqueda J, Olvera-Ramirez A, Aguilar-Tipacamú G, *et al.*, 2012. Current advances in detection and treatment of babesiosis. *Current Med Chem* 19:1504-18.
- Muraleedharan K, 2015. Babesia and babesiosis in livestock of Karnataka state, India-an overview. *Vet Res Int* 3:81-8.
- Nasr M, Soad MM, Wael MG, *et al.*, 2018. Molecular Detection of *Babesia Bigemina* and *Babesia Bovis* In Cattle In Behaira Governorate. *Europ J Pharm Med Res* 5:441-6.
- Ogedengbe JD, Hanner RH and Barta JR, 2011. DNA barcoding identifies *Eimeria* species and contributes to the phylogenetics of coccidian parasites (*Eimeriorina*, *Apicomplexa*, *Alveolata*). *Inter J Parasitol* 41:843-50.
- Oliveira-Sequeira T, Oliveira Mds, Araujo JJ, *et al.*, 2005. PCR-based detection of *Babesia bovis* and *Babesia bigemina* in their natural host *Boophilus microplus* and cattle. *Inter J Parasitol* 35:105-111.
- Oscar JBH and Cristian GR, 2018. Economic and Health Impact of the Ticks in Production Animals, Ticks and Tick-Borne Pathogens, Muhammad Abubakar and Piyumali K. Perera, IntechOpen, DOI: 10.5772/intechopen.81167. Available from: <https://www.intechopen.com/books/ticks-and-tick-borne-pathogens/economic-and-health-impact-of-the-ticks-in-production-animals>
- Pradeep RK, Nimisha M, Sruthi MK, *et al.*, 2019. Molecular Characterization of South Indian Field Isolates of Bovine *Babesia* spp. and *Anaplasma* spp. *Parasitol Res* 118:617-630.
- Rajabi S, Esmaeilnejad B and Tavassoli M, 2017. A molecular study on *Babesia* spp. in cattle and ticks in West-Azerbaijan Province, Iran *Vet Res Forum* 8:299-306.
- Ramos CA, Araújo FR, Alves LC, *et al.*, 2012. Genetic conservation of potentially immunogenic proteins among Brazilian isolates of *Babesia bovis*. *Vet Parasitol* 187:548-52.
- Saad F, Khaisroon M, Khan K, *et al.*, 2015. Prevalence and molecular detection of babesiosis in the slaughter animals of Peshawar Khyber Pakhtunkhwa Pakistan. *Int J Curr Microbiol App Sci* 4:1030-6.
- Saad-Roy C, Shuai Z and van den Driessche P, 2015. Models of bovine babesiosis including juvenile cattle. *Bulletin of Math Biol* 77:514-7.
- Terkawi MA, Alhasan H, Huyen NX, *et al.*, 2012. Molecular and serological prevalence of *Babesia bovis* and *Babesia bigemina* in cattle from central region of Syria. *Vet Parasitol* 187:307-11.
- Uilenberg G, 2006. Babesia-a historical overview. *Vet Parasit* 138:3-10.
- Vannier E and Krause PJ, 2009. Update on babesiosis. *Interdiscip Perspect Infect Dis* doi:10.1155/2009/984568.