

## RESEARCH ARTICLE

### Evaluating the Bactericidal Activity of Various Disinfectants against *Pseudomonas aeruginosa* Contamination in Broiler Chicken Hatcheries

Hazem M. Ibrahim<sup>1</sup>, Heba M. Salem<sup>2,3</sup>, Soha A. Alamoudi<sup>4</sup>, Nawal Al-Hoshani<sup>5</sup>, Abdullah M. Alkahtani<sup>6</sup>, Naheda M. Alshammari<sup>7</sup>, Lamaia R. Altarjami<sup>8</sup>, Eman A. Beyari<sup>7</sup>, Mohamed T. El-Saadony<sup>9\*</sup>, and Hanan S. Khalefa<sup>10</sup>

<sup>1</sup>Veterinary Serum and Vaccine Research Institute, Agricultural Research Center, Egypt; <sup>2</sup>Poultry Diseases Department, Faculty of Veterinary, Medicine Cairo University, Giza, 12211, Egypt; <sup>3</sup>Department of Diseases of Birds, Rabbits, Fish & their Care & Wildlife, School of Veterinary Medicine, Badr University in Cairo (BUC), Badr City, Cairo, Egypt; <sup>4</sup>Biological Sciences Department, College of Science & Arts, King Abdulaziz University, Rabigh 21911, Saudi Arabia; <sup>5</sup>Department of Biology, College of Science, Princess Nourah bint Abdulrahman University, P.O. Box 84428, Riyadh 11671, Saudi Arabia; <sup>6</sup>Department of Microbiology & Clinical Parasitology College of Medicine, King Khalid University, Abha 61413, Saudi Arabia; <sup>7</sup>Department of Biological Sciences, Microbiology, Faculty of Science, King Abdulaziz University, Jeddah 21589, Saudi Arabia; <sup>8</sup>Department of Chemistry, College of Science & Arts, King Abdulaziz University, Rabigh 21911, Saudi Arabia; <sup>9</sup>Department of Agricultural Microbiology, Faculty of Agriculture, Zagazig University, Zagazig 44511, Egypt; <sup>10</sup>Department of Veterinary Hygiene and Management, Faculty of Veterinary Medicine, Cairo University, Giza, 12211, Egypt

\*Corresponding author: m.talaatsadony@gmail.com

#### ARTICLE HISTORY (24-171)

Received: March 19, 2024

Revised: May 31, 2024

Accepted: June 9, 2024

Published online: July 09, 2024

#### Key words:

Disinfectants

Broiler chicken

*Pseudomonas aeruginosa*

Glutaraldehyde

Hatchery

#### ABSTRACT

Hatcheries are hubs for incoming eggs and progeny flock output, making them a crucial component of the poultry production chain. This study involved performing quantitative microbiological tests in a commercial hatchery with numerous compartments, including an egg handling room, cold storeroom, setter room, hatcher room, and chick production hall. There were 150 air samples and 180 surface swabs collected in the incubator before and after disinfection over ten visits, in addition to 250 yolk sac and organ samples taken from late-dead embryos. As a result, surface swabbing could detect higher contamination levels than open-plate methods in all hatchery visits, mainly in handling eggs, cold storage, and hatcher halls. This study examines the presence of *Pseudomonas aeruginosa* strains in hatchery environments and dead embryos. Biochemical and PCR testing were used to identify *P. aeruginosa* using 16SrDNA primers at 956bp and the *tox A* gene at 396bp. In hatchery environmental samples, the incidence rate was 10.7%, and in dead embryos, it was 10%; therefore, maintaining good hygiene, especially in hatcheries, is essential for *Pseudomonas* species control. Subsequently, in this study, a virulent strain of *P. aeruginosa* was subjected to an *in vitro* test with 10 disinfectants from six chemical groups. Iodine compounds with phosphoric acids, per-acetic acid, sodium dichloroisocyanurate, and quaternary ammonium compounds (QACs) with glutaraldehyde compounds showed 100% microbial reduction even in the presence of organic matter with exposure times of 30 min. It was concluded that the most effective and cost-effective way to prevent and control infections is to use a disinfectant with sufficient concentration and exposure time in hatcheries.

**To Cite This Article:** Ibrahim HM, Salem HM, Alamoudi SA, Al-Hoshani N, Alkahtani AM, Alshammari NM, Altarjami LR, Beyari EA, El-Saadony MT and Khalefa HS, 2024. Evaluating the bactericidal activity of various disinfectants against *Pseudomonas aeruginosa* contamination in broiler chicken hatcheries. Pak Vet J, 44(3): 683-690. <http://dx.doi.org/10.29261/pakvetj/2024.203>

#### INTRODUCTION

A hatchery is an essential component of poultry production and provides an opportunity to segregate between generations of birds microbiologically; otherwise, control infectious diseases spread via the application of proper biosecurity measures (Abd El-Hack

*et al.*, 2023; Yousef *et al.*, 2023). Unfortunately, rather than acting as a firebreak for infectious agents, hatcheries may enable the spread and multiplication of infections among generations of poultry (Wales and Davies, 2020).

Microorganisms can adversely affect hatchability and cause embryonic death in a poultry hatchery environment (Kumar *et al.*, 2012). A hatchery is vulnerable to pathogens

that may be carried on or within eggs, on staff, on objects like trolleys and trays, or by air (Mcmullin, 2009). A commercial broiler hatchery was evaluated for hatchery hygiene using a surface swap and an open-plate method; despite the application of various sanitation measures, epidemiological investigations have revealed significant variances in microbial populations over time (Shehata *et al.*, 2019; Sallam *et al.*, 2023). Different bacteria were recovered from dead shell embryos that contaminate hatcheries (Bakheet and Torra, 2020); a particular hatchery-born pathogen is *P. aeruginosa*, known as gram-negative, non-spore-forming rods, motile, and fruity scent is produced because of the production of watery soluble green pigments (Elshafiee *et al.*, 2022; Salem *et al.*, 2024).

Chickens and embryos become infected with *P. aeruginosa*, causing respiratory complications, septicemia, and even death (Shahat *et al.*, 2019). *P. aeruginosa* virulence factors cause microbial invasion, multiplication, dissemination, tissue destruction, and septicemia, with high mortality rates in humans and animals (Poursina *et al.*, 2023). *P. aeruginosa* exotoxin A, a virulence factor that inhibits the biosynthesis of proteins at the polypeptide chain elongation factor 2, can cause severe damage to tissues and organs (Aljebory, 2018). According to Walker *et al.* (2002) and Marouf *et al.* (2023), *Pseudomonas* species are more germicide-resistant than other bacteria. In addition, the concentrations of disinfectants effective against *Salmonella* and *Staphylococcus* species are lower than those effective against *Pseudomonas* species (Gehan, 2009; Beier *et al.*, 2021a, b; Rabie *et al.*, 2023).

Further, appropriate hygienic measures like regular cleaning with wide-spectrum disinfectants are vital to eliminate pathogenic organisms from farms and hinder the re-infection of farm workers and animals (Davies and Wales, 2019). Thus, hatchery sanitation programs require safe and effective disinfectants that inhibit microorganism growth and maintain the hatchability of the eggs treated with them (McElreath, 2019). Additionally, convenient disinfectants that minimize sanitization time and are easy to use are needed (Gehan, 2009).

This study aimed to determine the sanitary situations in a poultry hatchery by bacteriological investigation of hatchery environmental samples during production (before and after disinfection) so they can identify the sources of bacterial contamination and implement appropriate preventive measures. In addition, an *in vitro* trial was conducted to evaluate the effectiveness of 10 disinfectants of different chemical groups at various contact times of 10, 30, and 60 minutes against a virulent strain of *Pseudomonas aeruginosa* isolated from poultry hatcheries were also investigated.

## MATERIALS AND METHODS

**Study area:** The Institutional Animal Care and Use Committee Vet ethically approves this work. CU. IACUC with code "Vet CU 08072023693". The current study was adopted in a broiler hatchery in the El Beheira governorate. A total of 40 incubators and 21 hatcher (multistage) were available, and each incubator held 115200 chicken eggs (Cobb). Each hatcher can hold approximately 38400 eggs. The relative humidity and

temperature were 25% and 36.5°C, respectively, for 18 days in the setting period, while the relative humidity and temperature of the hatcher were 30% and 36.0°C, respectively, for the rest of the three days.

**Sampling location and time:** Samples were taken on ten different morning dates as hatching chicks were processed after the hatchery had been cleaned and disinfected. During each visit, air samples and surface swabs were collected from egg handling rooms, cold storerooms, setter rooms, hatcher rooms, chick production halls, and organ and yolk sac swabs collected after hatching from the dead in-shell embryos. One hundred fifty air samples (inlet of airflow, Air conditioner, air ducts, and fans) were collected for bacterial load detection, and 180 swabs were collected from the surfaces of equipment and facilities (wall, floor, Trolley, Box). The hatchery was checked ten times from September 2020 to October 2021. The first, second, third, fifth, and seventh investigations were conducted before cleaning and disinfection in September 2020, December 2020, January 2021, April 2021, and June 2021, respectively. The fourth, sixth, eighth, ninth, and tenth examinations were performed immediately post cleaning and disinfection (after disinfection) of the hatchery post the elimination of chicks in March 2021, April 2021, July 2021, September 2021, and October 2021, respectively. Also, 200 swabs from yolk sacs of late dead in-shell embryos were taken in sterile plastic containers, preserved in ice boxes, and rapidly transferred to the laboratory for further examinations, and 50 samples were collected from hatched dead chicks (obtained from these visits) at the first two weeks of age.

**Examination for the bacterial load of air, the surface of equipment & facilities in the studied hatchery:** For air samples, the open-plate method was used; plate count agar (Difco, Detroit, MI) was used for aerobic bacteria. The prepared agar plates were subjected to be uncovered for 10 min at a meter height from the floor surface (Berrang *et al.*, 1995). Once the plates were submitted to the lab, they were kept for 24hrs. at 37°C. Microbial loads were expressed as colony forming units (CFU) per 10cm diameter plate, after which they were scored for bacterial multiplication following Table 1.

Surface samples were collected using microbiological swabs. Sterile cotton swabs were rubbed in sterile saline solution on a surface of 16cm<sup>2</sup> using sterile metal templates. The samples were transported in tubes containing 10 mL of sterile saline solution and refrigerated in the lab. A sterile saline solution was used to prepare ten-fold dilutions. Aerobic bacteria were quantified using aliquots of 0.1 mL from the primary dilution and decimal dilutions to plate count agar (Hi Media, India) (Lazarov *et al.*, 2018).

**Bacteriological isolation and identification of *P. aeruginosa*:** A total of 108 samples were collected from hatchery environment (50 aerial samples were tested using an open-plate method, and 58 samples from hatchery surfaces). Also, 200 swabs from late dead in shell embryos and 50 pooled liver, heart blood, and yolk sac samples from the investigated hatchery were collected for further bacteriological examination. Samples were taken

**Table 1:** Interpretation reference scored for bacterial growth inside poultry hatchery

Colony Count	Score	Grade
0-5	0	None or very slight contamination (considered excellent)
6-15	1	Slight contamination (considered good)
16-30	2	Moderate contamination (borderline acceptable)
31-50	3	Significant contamination (poor)
>50	4	heavy contamination (unacceptable)
TNTC	5	very heavy contamination (unacceptable)

from the different hatchery rooms and labeled along the whole hatchery's processing pathway, from the egg handling room to the chick production hall.

Samples were quickly transported to the lab in ice boxes and kept in sterile plastic containers. Then the swab samples were put in nutrient broth and kept at 37°C for 24 hrs., then sub-cultured onto selective agar (MacConkey and *Pseudomonas* agar base medium with C-N supplement) and incubated at 37°C for 24 hrs. In addition, subculture onto a nutrient agar plate to observe the *P. aeruginosa* pigmentation (Elshafiee *et al.*, 2022). The supposed colonies were exposed to biochemical examinations such as catalase, oxidase, gelatin liquefaction, methyl red, arginine hydrolysis, indole, and urease (Cheesbrough, 2006).

**Molecular identification of *P. aeruginosa*:** The DNA was extracted using the QIA amp DNA Mini Kit (Qiagen, Germany) from all biochemically positive *P. aeruginosa* (n=36; 11 environmental samples and 25 dead embryos) maintained overnight in TSB broth. Using primers from Metabion, Germany, PCR was used to identify virulence-associated genes (*toxA*) and *Pseudomonas* species-specific genes (16S rDNA). The reaction volume (25µl) contained 12.5µl of 2x premix Emerald Amp GT PCR master mix (Takara), forward and reversed primer (20 p mol; 1µl each), PCR grade water (4.5µl) and template DNA (6µl). The reaction circumstances were summarized in Table 2 and the PCR products were inserted in electrophoresis in 1.5% agarose gel (AB gene). A 100bp DNA Ladder (Qiagen, USA) determines the fragment sizes. The gel picture was taken via a documentation system; then, the data was kept via computer software.

**In vitro evaluation of disinfectants against *P. aeruginosa*:** Using a modified version of the quantitative suspension test established by the British Standard Institute (BSI, 2019), the effect of disinfectants on *P. aeruginosa* strains was evaluated. The manufacturer's instructions state that the disinfectants were made in sterile distilled water (Table 3). *P. aeruginosa* was cultured in TSB broth overnight to produce bacterial suspension, which was then turbidity-adjusted to 0.5 MacFarland standards. In sterile test tubes, ten microliters of disinfectant were mixed with 0.1 mL of adjusted bacterial broth in each tube after being individually evaluated. Control tubes contain only diluted bacterial suspensions. At room temperature (25°C), each disinfectant was incubated with a bacterial suspension for various contact durations of 10, 30, and 60 min. The disinfectant was neutralized using 8 mL of neutralizing broth (Sigma) containing 0.5% sodium thiosulphate. Similar tests were conducted again using organic matter

(2% yeast extract). Following a neutralizing period, triplicate subcultures from each set of tubes were placed on cetrimide agar plates, and the plates were left at 37°C overnight to assess the microbial proliferation. The equation  $\log \text{reduction} = \log_{10} (\text{initial CFU/ml}) - \log_{10} (\text{final CFU/ml})$  was used to compute the logarithmic reduction of the growth in each solution and the control.

**Statistical analysis:** The statistical analysis was performed utilizing SPSS software and one-way ANOVA (Park, 2009). All examined means (treatments) were compared using the LSD test, ensuring significance at a probability of  $P < 0.05$ .

## RESULTS

**Microbiological examination of air and surface samples:** Table 4 shows the results of aerobic bacterial air pollution. The mean count from the before disinfection investigation (first, second, third, fifth, and seventh) sampling times, which collected samples during egg incubation, was higher than that of hatchery sampling sites after cleaning and disinfection. The higher mean bacterial count isolation was from the air duct of setter hall 1, measuring; (410 CFU/Ø Petri-dish) whereas in the other sampling sites, hatchery hall 2, vaccine preparation room, and boxes washing room, it was minimal, TBC about (200 CFU/Ø petri-dish). The average bacterial score for every visit was calculated, and we observed that before disinfection, hatchers' visits ranged from slight to moderate contamination. In contrast, the other after-disinfection hatcher's visits show non-contamination (score 0-1), as shown in Fig. 1.

Table 5 shows the surface contamination caused by aerobic bacteria on the equipment and facilities. The most contaminated sites in post-disinfection hatcher's visits were the floor and eggshell in the egg handling room of more than ( $10^3$  CFU /16 cm<sup>2</sup>) and the eggshell and floor of the cold storeroom, followed by the floor of the setter and floor of the hatchery and box in the chick production hall. Regardless of the hatchery's cleanliness and sanitization, except the egg handling room, aerobic bacteria after disinfection hatcher's visits were moderate, measuring less than (100 CFU/16 cm<sup>2</sup>) at all sampling times, floor of cold storeroom, floor of setter, and box in chick production hall measuring ( $1.26 \times 10^3$ ,  $14.68 \times 10^2$ ,  $14.68 \times 10^2$ , and  $0.810 \times 10^3$  CFU/16 cm<sup>2</sup>) respectively. The remaining sampling sites were contaminated but to a relatively low level, measuring around 10 CFU/16 cm<sup>2</sup> for aerobic bacteria. The microbial contamination of the chick processing room in the two hatchery investigation sets was higher than in the other sampling sites. Surface swabbing detected a higher degree of microbial contamination than the open-plate method in the investigated hatchery along different sampling times, as shown in Fig. 2.

***P. aeruginosa* isolation from hatcheries and molecular detection:** The prevalence of *P. aeruginosa* in environmental samples was 10.7%, with eleven isolates found out of 103 total samples, as shown in Table 6. Morphologically, *P. aeruginosa* appears as pale colonies of non-lactose fermenter on MacConkey agar. The

**Table 2:** Primer sequences for detection of *Pseudomonas* and specific Exotoxin A gene (toxA) of *P. aeruginosa*

Target	gene	Oligonucleotide sequence 5'-3'	Annealing temp. (°C)	Fragment size (bp)	References
<i>Pseudomonas</i> spp.	16S rDNA	GACGGGTGAGTAATGCCTA CACTGGTGTTCCTTCCTATA	20 sec. at 54°C	618	Spilker et al. (2004)
<i>P. aeruginosa</i>	16S rDNA	GGGGGATCTTCGGACCTCA TCCTTAGAGTGCCCCACCCG	20 secs at 54°C	956	
<i>P. aeruginosa</i> (Virulence gene)	toxA	GACAACGCCCTCAGCATCACCAGC CGCTGGCCCATTCGCTCCAGCGCT	55°C 45 sec.-	396	Matar et al. (2002)

**Table 3:** The disinfectants used for in vitro testing efficacy against virulent strains of *P. aeruginosa* isolated from the hatchery. were classified according to their active compounds and concentration recommendations

Active compound	Disinfectant	Producer country	Composition	Concentration
Aldehyde & quaternary ammonium compounds	A	Hungary	Glutaldehyde 150 gm.-QACs 100gm- Draymarin Brilliant Blue 0.4gm- Nrinazine 0.3gm-Azorobin 0.3gm	0.5%
	B	England	Glutaldehyde 15% - QACs 10%	0.5%
Acidic compound	C	England	Phosphoric acid 10%- Sulfonic acid 30% - chlorinated phenols 40%	0.4%
	D	Egypt	Orthophosphoric acid 60% - Formic acid 10%	1%
Iodine	E	England	Iodine 5% - Phosphoric acid 14% - Alcohol ethoxylate 24%	0.25%
	F	USA	1.75% titratable iodine	0.4%
Peracetic acid-hydrogen peroxide	G	Belgium	The stabilized mixture of Peracetic acid-hydrogen peroxide -organic acids- wetting agents -belong term stabilizer	2%
Ethoxylated Alcohol	H	Egypt	Sodium Hydroxide N-oxide amine Ethoxylated Alcohol	1.7%
Sodium	I	Ireland.	2.5 gm DiChloro Iso Cyanurates 62% in form of tablet	1/15 l
Dichloroisocyanurate 15	J	England	Potassium Persulfate 50% + sodium dichloroisocyanurate NaDCC 2.5%	0.5%

**Table 4:** Mean±SE of bacterial contamination (CFU/ Ø Petri dish) of air samples using the open plate method in the studied poultry hatchery at different sampling times (before disinfection, and h) and sampling sites from different hatchery's process steps

Sampling site		Before disinfection						After disinfection					
		1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	5 <sup>th</sup>	7 <sup>th</sup>	*Mean ±SE	4 <sup>th</sup>	6 <sup>th</sup>	8 <sup>th</sup>	9 <sup>th</sup>	10 <sup>th</sup>	*Mean ±SE
1	Air conditioner of Egg handling hall	3	42	11	1	14	14.2±7.36	1	2	0	2	0	1±0.45
2	Fan of preheating room	4	31	0	9	5	9.8±5.49	3	7	3	2	2	3.4±0.93
3	Air duct of Inovo hall	2	16	2	40	0	12±7.56	3	1	4	25	1	6.8±4.59
4	Air duct of setter hall 1	TNTC	TNTC	30	3	21	410.8±240.58	6	5	0	0	3	2.8±1.24
5	Air duct of setter hall 2	4	9	0	0	4	3.4±1.66	15	18	2	0	0	7±3.92
6	Setter 30 right side	1	0	2	4	8	3±1.41	0	21	6	10	4	8.2±3.95
7	Air duct of hatcher hall 1	42	3	4	0	4	10.6±7.88	7	1	1	0	1	2±1.26
8	Air duct of hatcher hall 2	4	4	TNTC	2	1	202.2±199.45	6	6	3	1	1	3.4±1.12
9	Air duct of hatcher hall 3	1	0	0	23	4	5.6±4.41	0	2	9	50	1	12.4±9.53
10	Hatcher	9	20	7	3	43	16.4±7.22	3	6	12	6	1	5.6±1.86
11	Air duct of Chick production hall	12	19	50	40	11	26.4±7.88	30	20	14	3	6	14.6±4.87
12	Vaccine storeroom	12	10	13	9	3	9.4±1.75	9	4	3	16	8	8±2.30
13	Vaccine preparation room (1 Day old)	50	TNTC	14	4	10	215.6±196.26	0	9	4	10	18	8.2±3.04
14	Air conditioner of cold storeroom	0	0	0	0	2	0.4±0.4	0	0	1	1	1	0.6±0.24
15	Boxes washing room	TNTC	40	45	4	4	218.6±195.54	3	10	4	1	0	3.6±1.75
*P value		0.03											

Cfu = colony forming unit, TNTC= Too Numerous to Count &gt;300 CFU, SE= standard error, \* P value &lt;0.05 is significant between the 2 means values

**Table 5:** Mean ±SE of bacterial contamination (CFU/16 cm<sup>2</sup>) of walls, floor, and other surfaces in the poultry hatchery was investigated at different sampling times (before disinfection and After disinfection) and sampling sites based on the different hatchery process stages


Sampling site		Before disinfection						After disinfection					
		1 <sup>th</sup>	2 <sup>th</sup>	3 <sup>th</sup>	5 <sup>th</sup>	7 <sup>th</sup>	Mean×10 <sup>2</sup> ±SE ×10 <sup>2</sup>	4 <sup>th</sup>	6 <sup>th</sup>	8 <sup>th</sup>	9 <sup>th</sup>	10 <sup>th</sup>	Mean ×10 <sup>2</sup> ±SE ×10 <sup>2</sup>
Subunit/Surface samples													
Egg handling room	Floor	20	20	3×10 <sup>3</sup>	—	5×10 <sup>2</sup>	13±7.4	100	0	—	—	0	33.3±33.3
	Wall	30	0	0	—	12	10.5±7.09	0	10	120	—	0	32.5±29.26
	Eggshell	3.2×10 <sup>2</sup>	4.2×10 <sup>2</sup>	40×10 <sup>2</sup>	—	6.4	13.5±8.87	—	30×10 <sup>2</sup>	1.2×10 <sup>2</sup>	—	0	10.33±9.83
Cold storeroom	Floor	33×10 <sup>2</sup>	30×10 <sup>2</sup>	40	300	9×10 <sup>2</sup>	14.7±6.44	2.9×10 <sup>3</sup>	1.1×10 <sup>3</sup>	0	0.2×10 <sup>3</sup>	10	8.4±5.69
	Wall	0	100	0	80	10	38 ±21.54	0	30	90	10	0	26± 16.91
	Eggshell	90	31× 10 <sup>2</sup>	—	10.2×10 <sup>2</sup>	±3.9×10 <sup>2</sup>	11.2± 6.54	20	6× 10 <sup>2</sup>	10×10 <sup>2</sup>	10	0	3.26 ±2.03
Setter hall	Floor of setter hall	60	20× 10 <sup>2</sup>	—	100	50	5.2±4.82	5× 10 <sup>2</sup>	7× 10 <sup>2</sup>	—	—	0	4 ±2.08
	Wall of Setter Hall	0	0	—	0	6× 10 <sup>2</sup>	150 ±150	0	10	—	—	0	3.3 ±3.3
	Trolley	10	—	—	3.8× 10 <sup>2</sup>	±3.6×10 <sup>2</sup>	250±120.14	300	20×10 <sup>2</sup>	—	20	0	5.80±4.78
Hatcher halls	Floor of hatcher 5	80× 10 <sup>2</sup>	55× 10 <sup>2</sup>	80×10 <sup>2</sup>	6.8× 10 <sup>2</sup>	150	5.9±120.69	20	0	4.2×10 <sup>2</sup>	20	0	92± 82.12
	Floor of hatcher 18	50	2× 10 <sup>3</sup>	100	0.8× 10 <sup>3</sup>	0.39×10 <sup>3</sup>	7±3.6	10	10	10×10 <sup>2</sup>	10	0	2.55±248.34
Chick production hall	Floor	6× 10 <sup>2</sup>	210	8× 10 <sup>2</sup>	4× 10 <sup>2</sup>	4.5× 10 <sup>2</sup>	4.9± 99.07	10	0	40	50	0	20 ±10.49
	Wall	10	10	0	100	0	24 ± 19.13	—	230	80	3.2×10 <sup>2</sup>	0	157.5±72.15
	Box	2.5× 10 <sup>3</sup>	0.45×10 <sup>3</sup>	1.4× 10 <sup>3</sup>	0.29×10 <sup>3</sup>	0.31×10 <sup>3</sup>	8±3.2	—	10	20×10 <sup>2</sup>	130	0	5.35 ±4.89
Roll of Counting machine		7.5× 10 <sup>2</sup>	—	10	9× 10 <sup>2</sup>	—	5.5±275.10	0	—	—	0	0	0±0

Cfu= colony forming unit, SE= standard error, \* P value &lt;0.05 is significant between the 2 means values.


organism is characterized by its greenish diffusible pigment and fruity smell. Biochemically, it shows positive results for oxidase, catalase, urea, citrate utilization, and gelatin hydrolysis but negative for indole, methyl red, and

Voges Proskauer.

The highest contamination of *P. aeruginosa* was found on surfaces in the hatchery (16.9%), particularly in the cold storeroom and exhaust setter. Air samples



**Fig. 1:** Total mean score of aerial bacterial contamination of poultry hatchery in different sampling times.



**Fig. 2:** Mean  $\pm$  SE of bacterial contamination in poultry hatchery environment (from air samples and surfaces) at different sampling times.

**Table 6:** Incidence of identified *P. aeruginosa* isolated along the study period from the from the investigated hatcher environmental samples (air and surface; n=103)

Nature of sample		Before disinfection		After disinfection		Total
		No. of samples	No. of positive (%)	No. of samples	No. of positive (%)	
Air samples	Air conditioner of Egg handling hall	5	0(0)	5	0(0)	2/50 (4%)
	Hatcher air samples	5	0(0)	5	1(20)	
	Setter air samples	5	0(0)	5	0(0)	
	Air duct of Chick production hall	5	0(0)	5	0(0)	
	Box storeroom	5	0(0)	5	1(20)	
Surfaces	The floor of the egg handling room	4	0(0)	-	-	9/53 (16.98%)
Swabs	cold storeroom	-	-	3	1(33.3)	
	The floor of the setter hall	5	1(20)	5	0(0)	
	Exhaust setter	3	0(0)	3	1(33.3)	
	Floor of hatcher	5	2(40)	5	1(20)	
	floor of Chick production hall	5	2(40)	5	0(0)	11/103 (10.67%)
	Vacuum of Chick production hall	5	0(0)	5	1(20)	
Total						

The chi-square statistic between environmental samples (air and water) is 4.54, and the *p*-value is .033, Significant at (*P*<0.05).

**Table 7:** The incidence of *P. aeruginosa* isolated from chicken in Elbehira governorate hatchery

Type of samples	Site of collection	No examined samples	No positive isolates	%
Dead In Shell Embryos (swabs from liver, heart blood, and yolk sacs)	Before disinfection hatchery investigations	100	7	7
	After disinfection hatchery investigations	100	8	8
Young chicks (1-2 weeks of chick incubation)	At farm level	50	10	20
Total	-	250	25	10

**Table 8:** The Efficacy of ten different disinfectants belong to six different chemical groups at various contact times of 10, 30, and 60 minutes against a virulent strain of *Pseudomonas aeruginosa* (titer of  $1.5 \times 10^8$ /ml) isolated from studied poultry hatchery


The used disinfectants			Presence organic matter						Absence organic matter					
			10 min		30 min		60 min		10 min		30 min		60 min	
			Count	R %	Count	R %	Count	R %	Count	R %	Count	R %	Count	R %
1	A	0.5%	-	100	-	100	-	100	-	100	-	100	-	100
2	B	0.5%	3×10 <sup>6</sup>	98	-	100	-	100	5.3×10 <sup>5</sup>	99.6	-	100	-	100
3	C	0.4%	2×10 <sup>6</sup>	98.6	-	100	-	100	-	100	-	100	-	100
4	D	1%	-	100	-	100	-	100	-	100	-	100	-	100
5	E	0.25%	-	100	-	100	-	100	-	100	-	100	-	100
6	F	0.4%	9×10 <sup>6</sup>	94	3.5×10 <sup>6</sup>	97.6	-	100	-	100	-	100	-	100
7	G	2%	-	100	-	100	-	100	-	100	-	100	-	100
8	H	1.7%	3×10 <sup>7</sup>	80	22×10 <sup>6</sup>	85.33	45×10 <sup>5</sup>	97	21×10 <sup>6</sup>	86	12×10 <sup>6</sup>	92	2×10 <sup>6</sup>	98.66
9	I	1/15 l	-	100	-	100	-	100	-	100	-	100	-	100
10	J	0.5%	-	100	-	100	-	100	-	100	-	100	-	100

R %: reduction percent

taken after cleaning and disinfection had lower contamination levels (4%). *P. aeruginosa* is not found in earlier processing rooms, such as the egg handling and setter rooms. Out of 250 samples taken from all hatchery visits (200 from dead in-shell embryos and 50 from freshly dead birds), only 25 samples were *P. aeruginosa* positive (10%); this information is presented in Table 7. The freshly dead birds showed postmortem lesions that appeared in shape with varied degrees of congestion, omphalitis, severe pneumonic lungs, air sacculitis, kidney lesions with distended

ureters with ureates, enteritis, and unabsorbed yolk sac (in some cases greenish discoloration of yolk sac has been noticed) (Fig. 3).

**In vitro efficacy of disinfectants on *P. aeruginosa*:** As indicated in Table 8, the outcomes demonstrated that all disinfectants were effective against the tested *P. aeruginosa* isolates. In the presence of organic matter, the disinfectants C and B showed 100% reduction after 30 min, disinfectant (F) showed 100% at 60 min, while disinfectant (H) failed to eliminate contamination post 60



**Fig. 3:** The postmortem examination of freshly dead chicks (obtained from the investigated hatcheries) showed A: arrow refers to unabsorbed yolk sac with greenish discoloration with distended intestine with a mild degree of enteritis; B: arrows refer to pathological changes in the kidney with distended ureters with ureates; C: arrows refer to severe degree of lung congestion and pneumonia

min. While in the absence of organic matter, Except for disinfectant (B); show 100% reduction after 30 and 60 minutes, and disinfectant (H) which resulted in 98.6% reduction after 60 minutes, the rest disinfectants demonstrated superior effectiveness in removing the microbes (100% reduction after 10 min).

## DISCUSSION

Different pathogens transmitted due to hatcheries contamination, they resulted in financial losses in poultry industry (Abd El-Hack *et al.*, 2022a, b; El-Saadony *et al.*, 2022; Khalifa *et al.*, 2023). Several authors have suggested different ways to maintain biosecurity and hygiene in hatcheries (Bennett, 2017).

According to the results, surface swabbing identified a higher level of microbial contamination in the examined hatchery than the open-plate method. Compared to other frequently used rooms, the air duct of the setter hall had the highest level of aerobic bacterial contamination (410 CFU), followed by the air duct of the hatcher hall (202 CFU). Additionally, there are high counts of >200 CFU in the vaccine preparation and box washing rooms. The active hatcher had high concentrations of coliform, fungi, and aerobic bacteria in the air, measuring over 300 CFU /63.6 cm<sup>2</sup> (Kim and Kim, 2010).

The egg sorting room had a moderate level of contamination. However, this was not the case in other spaces, like the setter room, candling-transfer room, and chick counting room, where contamination was minimal and measured less than 10 CFU /63.6 cm<sup>2</sup> for aerobic bacteria. According to Fig. 1, the second hatchery's investigation time had the highest mean score for bacterial contamination (1.48), while the fourth hatchery's investigation time had the lowest (0.23) also, the chick processing room had microbial contamination levels over 100 CFU/16 cm<sup>2</sup>, the highest of all the areas sampled. According to the current study, surfaces had higher bacterial count than samples collected from air. The results concur with McElreath's (2019) discovery that although the open-plate method is quick, cheap, and easy to use, it can only identify living microorganisms. In contrast, Kim and Kim (2010) discovered that the levels of bacterial contamination on equipment and facilities were similar to those in the air. However, there were higher levels on surfaces in certain areas of the hatchery. *P. aeruginosa* is a serious hatchery-borne disease that can infect and colonize fertilized and

embryonated eggs, causing in-shell death to embryos and chicks after hatching (Dinev *et al.*, 2013).

Among the most toxic virulent factors of pathogenic *P. aeruginosa* is exotoxin A, which inhibits eukaryotic protein synthesis and promotes tissue necrosis (Eman *et al.*, 2017). *P. aeruginosa* was isolated from environmental samples at a rate of 10.67%, with only two out of 50 air samples and 9 out of 53 surface samples being positive. Similarly, to Gehan, (2009), in current study it was found that some bacterial strains could not be isolated through the open-plate method but could be detected by surface swabs. The incidence of *P. aeruginosa* surface isolation differs significantly between samples from air and surface swabs.

The isolation of *P. aeruginosa* virulence strains from the environment is associated with the isolation from dead embryos, as environmental pollution causes a significant issue in chicken hatcheries as it affects recently hatched chickens, resulting in an elevated rate of embryo death (Dinev *et al.*, 2013). The incidence of *P. aeruginosa* isolates was 7.5% from the yolk sac of dead in-shell embryos and 20% from slow chicken. The current results are like those of Bakheet *et al.* (2017) and Shahat *et al.* (2019), who isolated *P. aeruginosa* from chicks with 18.6% and 20% incidence rates, respectively. Also, Hassan, (2013) recovered *P. aeruginosa* from freshly dead broiler chickens and one-day-old chicks in incidences of 25.3 and 10%, respectively.

Elsayed *et al.* (2016) isolated *P. aeruginosa* with a percentage of 22.9% and a high isolation rate from dead-in-shell embryos yolk sac (52%). Still, the liver samples 2- 40 days old, diseased, and freshly dead were (12%). These findings were validated by Kebede, (2010), who demonstrated that experimentally infection with *P. aeruginosa* during the hatching period from the environment or by infiltrating the eggshell of the embryo was the major cause of the high mortality rate in unhatched chicken and early chicks, which led to death.

In this study, the observed postmortem lesions in the freshly dead birds were omphalitis septicemic, including congestion of subcutaneous tissues and muscles with increased size and congestion of the parenchymatous organs and in some freshly dead birds, unabsorbed yolk sac, enteritis, air sacculitis, and pneumonia were recorded. These findings concur with Walker *et al.* (2002) recorded in freshly dead SPF chicks due to *P. aeruginosa* experimental infection. Hatcheries are vulnerable to infectious agents that may enter on or within eggs, on



staff, on objects like trolleys and trays, or as airborne contaminants (Marouf *et al.*, 2023; Saad *et al.*, 2023; Elsayed *et al.*, 2014).

The current research showed that compounds containing glutaraldehyde, such as disinfectant A, efficiently eliminated *P. aeruginosa* even with organic matter. According to Jiang *et al.* (2018), QACs eliminated 1% of the bacteria after 30 & 60 minutes, and disinfectant B demonstrated effectiveness after 30 minutes. The results supported earlier studies' results (Gehan, 2009). When no organic matter is present, the phenolic compound disinfectant C effectively eliminates germs within 10 minutes. It takes 30 minutes to be completely effective if organic matter is present. These findings align with those made by McLaren *et al.* (2011), who found that phenolic disinfectants tend to be consistent in efficacy.

Even in organic matter, *P. aeruginosa* was quickly eradicated by disinfectant (D). This result is in line with the efficiency of organic acids as bactericidal agents, which are preferable to antibiotics because they do not result in the emergence of bacterial strains resistant to the drugs (Novickij *et al.*, 2019). Gram-negative bacteria, including *Pseudomonas* species, have long been known to be resistant to low pH. According to the current study, disinfectant (E) is more efficient than disinfectant (F) because it achieved a 100% reduction after only 10 minutes. Hydrogen peroxide and PAA are combined to create a disinfectant (G), effective against microbes even when organic matter is present. This concurs with the suggestion of Rodgers *et al.* (2001) to use H<sub>2</sub>O<sub>2</sub> as a disinfectant in hatcheries. However, PAA was not recommended as the preferred sanitizing agent for chicken processing equipment by Rossoni and Gaylarde (2000).

A disinfectant (H) with a 0.5% concentration is specifically used to clean eggshells. However, in the current study, even after 60 minutes of contact time, both with and without organic matter, this concentration could not wholly eradicate *P. aeruginosa*. Only 98% and 97% of *P. aeruginosa* were reduced, respectively. This study discovered that with and without organic matter, sodium di-chloroisocyanurate compounds effectively eliminated microbes. The findings support earlier research that demonstrated sodium hypochlorite successfully lowered *P. aeruginosa* counts (Krause *et al.*, 2019). This contrasts with a prior study by Gharieb *et al.* (2022), which discovered that 3% sodium hypochlorite could only reduce the microbe count by 89% even after 60 minutes of contact with organic matter.

Overall, the current findings supported those of Moustafa *et al.* (2009), who claimed that QAC, glutaraldehyde, and per-acetic acid had demonstrated their efficacy in preventing the contamination of poultry hatcheries and could be used as effective formaldehyde substitutes. In a lab setting, Glutaraldehyde 1% and hydrogen peroxide 3% were effective in a short time on specifically *P. aeruginosa* in the absence and presence of organic matter compared to other disinfectants, according to Gharieb *et al.* (2022).

**Conclusions:** Based on virulent genes and isolation from the hatchery environment, *P. aeruginosa* was a critical pathogen causing the deaths of newly hatched chicks during the first two weeks of their rearing age. Therefore,

maintaining good hygiene, especially in hatcheries, is essential for the control of *Pseudomonas* spp. Iodine compound with phosphoric acids (disinfectant, E), per-acetic acid (disinfectant, G), sodium dichloroisocyanurate (disinfectant I; disinfectant J), and QAC with glutaraldehyde compound (disinfectant, A) all show *in vitro* 100% microbial reduction even in the presence of organic matter with exposure times of 30 min. Further studies are recommended to evaluate these disinfectants in hatcheries in the field.

**Acknowledgements:** The authors gratefully acknowledge Princess Nourah bint Abdulrahman University Researchers Supporting Project number (PNURSP2024R437), Princess Nourah bint Abdulrahman University, Riyadh, Saudi Arabia. The authors extend their appreciation to the Deanship of Scientific Research at King Khalid University for funding this work through large group Research Project under grant number RGP2/530/44.

**Competing interests:** The authors declare that they have no competing interests.

**Authors contributions:** Conceptualization, HMI, HMS, SAA, EAB, MTES, and HSK, formal analysis, NAH, AMA, NMA, and LRA, investigation, HMI, HMS, SAA, EAB, MTES, and HSK, data curation, NAH, AMA, NMA, and LRA, writing original draft preparation, HMI, HMS, SAA, EAB, MTES, and HSK, writing final manuscript and editing, NAH, AMA, NMA, and LRA, visualization and methodology, HMI, HMS, SAA, EAB, MTES, HSK, NAH, AMA, NMA, and LRA. All authors have read and agreed to the published version of the manuscript.

## REFERENCES

- Abd El-Hack ME, El-Saadony MT, Salem HM, *et al.*, 2022a. Alternatives to antibiotics for organic poultry production: types, modes of action and impacts on bird's health and production. *Poult Sci* 101(4): 101696.
- Abd El-Hack ME, El-Saadony MT, Saad AM, *et al.*, 2022b. Essential oils and their nanoemulsions as green alternatives to antibiotics in poultry nutrition: a comprehensive review. *Poult Sci* 101: 101584.
- Abd El-Hack ME, Salem HM, Khafaga AF, *et al.*, 2023. Impacts of polyphenols on laying hens' productivity and egg quality: A review. *J Anim Physiol Anim Nutr* 107: 928-947.
- Aljebory IS, 2018. PCR detection of some virulence genes of *Pseudomonas aeruginosa* in Kirkuk city, Iraq. *J Pharm Sci Res* 10(5): 1068-1071.
- Bakheet AA, Naglaa MA, Sayed HA, *et al.*, 2017. Soad AN. Detection of Disinfectant resistant aerobic bacteria in unhatched chicken eggs. *Benha Vet Med J* 32(2): 248-259.
- Bakheet AA and Torra DE, 2020. Detection of *Pseudomonas aeruginosa* in dead chicken embryo with reference to pathological changes and virulence genes. *Alexandria J Vet Sci* 65 (1): 81-89.
- Bennett B, 2017. The importance of biosecurity in the modern-day hatchery. *Int Hatchery Practice* 31: 21-23.
- Berrang E, Cox NA and Bailey JS, 1995. Measuring air-borne microbial contamination of broiler hatching cabinets. *J Appl Poult Res* 4: 83-87.
- Beier RC, Byrd JA, Andrews K, *et al.*, 2021a. Disinfectant and antimicrobial susceptibility studies of the foodborne pathogen *Campylobacter jejuni* isolated from the litter of broiler chicken houses. *Poult Sci* 100(2): 1024-1033.
- Beier RC, Andrews K, Hume ME, *et al.*, 2021b. Disinfectant and antimicrobial susceptibility studies of *Staphylococcus aureus* strains and ST398-MRSA and ST5-MRSA strains from swine mandibular lymph node tissue, commercial pork sausage meat and swine feces. *Microorganisms* 9(11): 2401.

- British Standard Institute (BSI. BS EN 1276), 2019. Chemical disinfectants and antiseptics-Quantitative suspension test for the evaluation of bactericidal activity of chemical disinfectants and antiseptics used in food, industrial, domestic and institutional areas-Test method and requirements (phase 2, step 1). London, UK: British Standards Institute.
- Cheesbrough M, 2006. District laboratory practice in tropical countries., 2<sup>nd</sup>, pp. 178–187. Cambridge: Cambridge University Press.
- Davies R, and Wales A, 2019. Antimicrobial resistance on farms: A review including biosecurity and the potential role of disinfectants in resistance selection. *Comp Rev Food Sci. Food Saf* 18(3): 753-774.
- Dinev I, Denev S and Beev G, 2013. Clinical and morphological studies on spontaneous cases of *Pseudomonas aeruginosa* infections in birds. *Pak Vet J* 33: 398-400.
- El-Saadony MT, Salem HM, El-Tahan AM, et al., 2022. The control of poultry salmonellosis using organic agents: an updated overview. *Poult Sci* 101(4): 101716.
- Elsayed MSA, Ammar AM, Al Shehri ZS, et al., 2016. Virulence repertoire of *Pseudomonas aeruginosa* from some poultry farms with detection of resistance to various antimicrobials and plant extracts. *Cell Mol Biol* 62(1): 1-5.
- Elsayed MM, El-Basrey YF, El-Baz AH, et al., 2014. Ecological prevalence, genetic diversity, and multidrug resistance of *Salmonella enteritidis* recovered from broiler and layer chicken farms. *Poult Sci* 103(2): 103320.
- Elshafie EA, Khalefa HS, Al-Atfeehy NM, et al., 2022. Biofilms and efflux pump regulatory gene (mexR) in multidrug-resistant *Pseudomonas aeruginosa* isolated from migratory birds in Egypt. *Vet World* 15(10): 2425.
- Eman MF, Heba R, Bakheet AA, et al., 2017. Advanced studies on *Pseudomonas aeruginosa* infection in chicken. *Anim Health Res* 5(4): 207-217.
- Gehan ZM, 2009. A new approach to evaluate the hygienic condition of commercial hatcheries. *Int J Poult Sci* 8 : 1047-1051.
- Gharieb R, Saad M, Khedr M, et al., 2022. Occurrence, virulence, carbapenem resistance, susceptibility to disinfectants and public health hazard of *Pseudomonas aeruginosa* isolated from animals, humans and environment in intensive farms. *J Appl Microbiol* 132: 256-267.
- Hassan HM, 2013. M. V. Sc. Thesis, Faculty of Veterinary Medicine, Beni-Suef University; Characterization of *Pseudomonas aeruginosa* Isolated from Different Pathological Lesions in Chickens.
- Jiang L, Li M, Tang J, et al., 2018. Effect of different disinfectants on bacterial aerosol diversity in poultry houses. *Front Microbiol* 9: 2113.
- Kebede F, 2010. *Pseudomonas* infection in chickens. *J Vet Med Anim Heal* 2:55-58.
- Khalifa WH, Sallam MG, Kamel NN, et al., 2023. Using in-ovo injection of olive pulp extract and vitamin C to improve hatchability, post hatch growth performance, carcass traits and some biochemical blood analysis in broiler chickens. *Int J Vet Sci* 12(3): 353-359.
- Kim JH and Kim KS, 2010. Hatchery hygiene evaluation by microbiological examination of hatchery samples, *Poult Sci* 89: 1389-1398.
- Krause KM, Haglund CM, Hebner C, et al., 2019. Potent LpxC inhibitors with in vitro activity against multidrug-resistant *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 63(11): 10-1128.
- Kumar MC, Farms EO, Pomeroy BS, et al., 2012. College of Veterinary Medicine University of Minnesota. Blankenship, Leroy, ed. Colonization Control of Human Bacterial Enteropathogens in Poultry. Academic Press, 243.
- Lazarov I, Zhelev G, Lytzkanov M, et al., 2018. Dynamics of microbial contamination in a poultry hatchery. *Arch Vet Med* 11(1): 37-44.
- Marouf S, Li X, Salem HM, et al., 2023. Molecular detection of multidrug-resistant *Pseudomonas aeruginosa* of different avian sources with pathogenicity testing and in vitro evaluation of antibacterial efficacy of silver nanoparticles against multidrug-resistant *P. aeruginosa*. *Poult Sci* 102(10) : 102995.
- Matar GM, Ramlawi F, Hijazi N, et al., 2002. Transcription levels of *Pseudomonas aeruginosa* exotoxin A gene and severity of symptoms in patients with otitis externa. *Curr Microbiol* 45:350-4.
- McElreath JS, 2019. Detection and Reduction of Microbial Load in Poultry Facilities (Doctoral dissertation, University of Georgia).
- McLaren I, Wales A, Breslin M, et al., 2011. Evaluation of commonly used farm disinfectants in wet and dry models of *Salmonella* farm contamination. *Avian Pathol* 40: 33-42.
- Mcmullin PF, 2009. Hygiene and microbiological control in hatcheries. *Avian Biol Res* 2: 93–97.
- Moustafa GZ, Amer AW, ELSabagh IM, et al., 2009. In vitro Efficacy comparisons of disinfectants used in the commercial poultry farms. *Int J Poult Sci* 8 (3) : 237-241.
- Novickij V, Lastauskienė E, Staigvila G, et al., 2019. Low concentrations of acetic and formic acids enhance the inactivation of *Staphylococcus aureus* and *Pseudomonas aeruginosa* with pulsed electric fields. *BMC Microbiol* 19 : 1-7.
- Park HM, 2009. Comparing group means: t-tests and one-way ANOVA using Stata, SAS, R, and SPSS.
- Poursina S, Ahmadi M, Fazeli F, et al., 2023. Assessment of virulence factors and antimicrobial resistance among the *Pseudomonas aeruginosa* strains isolated from animal meat and carcass samples. *Vet Med Sci* 9: 315-325.
- Rabie NS, Fedawy HS, Sedeek DM, et al., 2023. Isolation and serological identification of current salmonella species recovered from broiler chickens in Egypt. *Int J Vet Sci* 12(2): 230-235.
- Rodgers JD, McCullagh JJ, McNamee PT, et al., 2001. An investigation into the efficacy of hatchery disinfectants against strains of *Staphylococcus aureus* associated with poultry industry. *Vet Microbiol* 82: 131-140.
- Rossoni EM and Gaylarde CC, 2000. Comparison of sodium hypochlorite and peracetic acid as sanitizing agents for stainless steel food processing surfaces using epifluorescence microscopy. *Int J Food Microbiol* 61: 81-85.
- Saad N, El-Abasy MA, El-Khayat F, et al., 2023. Efficacy of chitosan nanoparticles as a natural antibacterial agent against pathogenic bacteria causing omphalitis in poultry. *Pak Vet J* 43(3) : 573-578.
- Salem M, Younis G, Sadat A, et al., 2024. Dissemination of mcr-I and  $\beta$ -lactamase genes among *Pseudomonas aeruginosa*: molecular characterization of MDR strains in broiler chicks and dead-in-shell chicks infections. *Ann Clin Microbiol Antimicrobe* 23(1) : 1-19.
- Sallam MG, Samy A, Yassein SA, et al., 2023. Influence of in-ovo feeding bovine serum albumin or L-glutamine to Japanese quails on hatchability, performance of hatched chicks, antioxidant activity, lymphoid organs, and some blood biochemical parameters. *Int J Vet Sci* 12(2): 212-217.
- Shahat HS, Mohamed H, Al-Azeem A, et al., 2019. Molecular detection of some virulence genes in *Pseudomonas aeruginosa* isolated from chicken embryos and broilers with regard to disinfectant resistance. *SVU-Int J Vet Sci* 2: 52-70.
- Shehata AA, Basiouni S, Elrazek AA, et al., 2019. Characterization of *Salmonella enterica* isolated from poultry hatcheries and commercial broiler chickens. *Pak Vet J* 39 : 515-520.
- Spilker T, Coenye T, Vandamme P, et al., 2004. PCR based assay for differentiation of *Pseudomonas aeruginosa* from other *Pseudomonas* species recovered from cystic fibrosis patients. *J Clin Microbiol* 42: 2074-2079.
- Wales A and Davies R, 2020. Review of hatchery transmission of bacteria with focus on *Salmonella*, chick pathogens and antimicrobial resistance. *Poult Sci* 76 : 517-536.
- Walker SE, Sander JE, Cline JL, et al., 2002. Characterization of *Pseudomonas aeruginosa* isolates associated with mortality in broiler chicks. *Avian Dis* 46: 1045-1050.
- Yousef HM, Hashad ME, Osman KM, et al., 2023. Surveillance of *Escherichia coli* in different types of chicken and duck hatcheries: one health outlook. *Poult Sci* 102:103108.