



RESEARCH ARTICLE

Morphological Description of Fatal Sarcocystosis in Cattle with Implication of their Immunological and Histopathological Alteration

Safia M.A. Bahshwan¹, Mina A. Almayouf², Hayat S. Al-Rashidi², Amal M. Alzahrani³, Samar Ahmad Khan⁴, Haifaa A. Mahjoub¹, Mada M. AL-Qurashi^{1*}, Nawal Al-Hoshani⁵, Mohammed A. Alqahtani⁶ and Marwa M. Attia⁷

¹Biological Sciences Department, College of Science and Arts, King Abdulaziz University, Rabigh 21911, Saudi Arabia;

²Department of Biology, College of Science, Qassim University, Buraydah 51452, Saudi Arabia; ³Department of Biology, Faculty of Sciences, Al-Baha University, Al-Baha 65779, Saudi Arabia.; ⁴Department of Biology, College of Science, University of Jeddah, Jeddah 21589, Saudi Arabia; ⁵Department of Biology, College of Science, Princess Nourah bint Abdulrahman University, P.O. Box 84428, Riyadh 11671, Saudi Arabia; ⁶Department of Biology, College of Science, King Khalid University Abha 61413, Saudi Arabia; ⁷Department of Parasitology, Faculty of Veterinary Medicine, Cairo University, Giza 12211, Egypt

*Corresponding author: mmalqurashi@kau.edu.sa

ARTICLE HISTORY (24-539)

Received: September 1, 2024
Revised: October 31, 2024
Accepted: November 6, 2024
Published online: November 16, 2024

Key words:

Cattle
Macrocysts
Cytokines
Histopathological examination
Morphological Features
Sarcocystis fusiformis

ABSTRACT

Sarcocystis (*S.*) is a cyst-forming parasite with over 200 species in the phylum Apicomplexa. This study aimed to investigate the prevalence of *S. fusiformis* in cattle and examine its impact on the animals. A total of 200 cattle carcasses were examined for the presence of *S. fusiformis* in various tissues, including the esophagus, heart, tongue, and skeletal muscles, to assess the extent of infection. The study also focused on analyzing the associated immunological and histopathological changes in infected animals. Infected tissues, including the esophagus, hearts, and muscles, were subjected to light microscopy, transmission electron microscopy (TEM), and histopathological examination. A 10% prevalence of *S. fusiformis* was found among the examined cattle. Toluidine blue-stained sections revealed that the sarcocyst walls ranged from 1.8 to 6.5µm in thickness and displayed branching villar protrusions. These protrusions were observed to have homogeneous, papillomatous forms with a width of 100 nm. TEM analysis revealed filamentous tubular structures within the sarcocyst walls, some reaching lengths up to 6µm. These tubules ran parallel to the long axis of the villar projections and lacked granules. Furthermore, the expression levels of various cytokines in the infected tissues were significantly elevated, including tumor necrosis factor-alpha (TNF-α), interleukin-1β (IL-1β), and interferon-gamma (IFN-γ). Histopathological examination revealed tissue changes, with a small number of mononuclear inflammatory cells present in the affected organs, indicating an immune response to the parasite. This study provides valuable insights into the unique characteristics of *S. fusiformis* sarcocysts, helping to distinguish them from other *Sarcocystis* species found in both domestic and wild ruminants. Further research is recommended to explore potential control measures and better understand the broader implications of *S. fusiformis* infections in livestock.

To Cite This Article: Bahshwan SMA, Almayouf MA, Al-Rashidi HS, Alzahrani AM, Khan SA, Mahjoub HA, AL-Qurashi MM, Al-Hoshani N, Alqahtani MM and Attia MM, 2024. Morphological description of fatal sarcocystosis in cattle with implication of their immunological and histopathological alteration. Pak Vet J. <http://dx.doi.org/10.29261/pakvetj/2024.280>

INTRODUCTION

Globally, there are approximately one billion beef cattle. However, compared to pigs and chickens, they are less efficient at converting feed into meat. This means that producing beef requires more resources and potentially has a greater environmental impact (Terry *et al.*, 2020;

Tenu *et al.*, 2023). These measurements, however, do not account for the fact that beef cattle provide high-quality protein from animal foods that are unsuitable for other species of livestock (Thomas *et al.*, 2021). Principal areas of emphasis for methods to enhance the productivity of beef cattle include the structure and content of feed, host genetics, the functioning of the rumen and pulmonary

microbiomes, and breeding and operational management (Lovarelli *et al.*, 2020; Thornton *et al.*, 2022). Strategies for improving beef production must consider not only the changing environmental conditions in which cattle are raised but also the overall health and immunity of the herd (Attia *et al.*, 2022a). Factors, i.e., climate change, resource availability, and disease resistance, need to be taken into account to ensure sustainable and efficient beef production (Kinley *et al.*, 2020; Palange and Dhattrak, 2021). Around 45% of the worldwide protein supply for humans is attributed to meat and milk derived from cattle and buffalo (Mottet *et al.*, 2017; Cordeiro *et al.*, 2022).

Parasitism in livestock production poses a significant threat, leading to various clinical signs, mortality, reduced production, increased condemnation rates, and economic losses (Soliman *et al.*, 2022a; Strydom *et al.*, 2023; Soliman *et al.*, 2024). *Sarcocystosis* is a major parasitic disease affecting livestock, with significant economic consequences. Infected cattle can experience severe health issues, including miscarriage, reduced milk production, neurological problems, and weight loss, potentially leading to death (Ahmed *et al.*, 2016; Badr *et al.*, 2024).

According to Castro-Forero *et al.* (2022), the majority of *Sarcocystis* (*S.*) species follow a life cycle featuring two hosts: herbivores as host intermediates and carnivores as ultimate hosts. *Sarcocystis* species are transmitted to the intermediate host by ingesting sporocysts, oocysts, or both expelled from the definitive host's feces (Fig. 1). The parasite has a short duration of infective schizogony before encysting in tissues to generate sarcocysts (Mousa *et al.*, 2016; Dubey and Rosenthal, 2023). Dogs are crucial in spreading this infection since they reside close to buffalo herds (Wielgus *et al.*, 2021). Feeding dogs the raw meat of killed animals and the high frequency of interaction between animals and domestic or stray dogs provide ideal conditions for the parasite, *S. cruzi* maintenance (Mendoza Roldan and Otranto, 2023; Pal *et al.*, 2023)

Sarcocysts are embedded in the tissues of intermediate hosts, which the definitive host consumes and then gets infected. The infected body parts with macrocysts can only be identified by inspection after slaughtering the infected animals at slaughterhouses (Constable *et al.*, 2016; Valizadeh and Ebrahimi, 2022).

The discovery of the *Sarcocystis* life cycle in 1972 initially led scientists to believe that only one species, *S. fusiformis*, infected both cattle and water buffalo. However, further research revealed that cattle harbor three distinct species: *S. hirsuta*, *S. hominis*, and *S. cruzi*. The *S. fusiformis* was then used solely to identify the species responsible for forming larger cysts (Mousa *et al.*, 2016; Dubey *et al.*, 2023).

Compared to cattle, the life cycles of *Sarcocystis* species that infect buffalo are poorly understood, especially the stages involving schizont and sarcocyst formation. Four species, i.e., *S. fusiformis*, *S. buffalonis*, *S. levinei*, and *S. dubeyi*, have been definitively identified in water buffalo (Dubey *et al.*, 2014; Ras, 2021). The largest sarcocysts in the genus are those of *S. fusiformis* (Barakat *et al.*, 2023; Menshawy *et al.*, 2023). Additionally, the study of El-Mahdi *et al.* (2023) investigated the life cycle and disease-causing potential of *S. cruzi* by examining infected cattle in El-Kharga, Egypt, and then studying

how the parasite develops and affects dogs, its final host. This study identified diverse forms of *S. cruzi* oocysts in cattle from Egypt's New Valley Governorate. Genetic analysis confirmed these parasites were indeed *S. cruzi*, highlighting this species as the most common in the region's cattle population. A high prevalence (76%) of *Sarcocystis* infection was found in mustelids, with six species typically associated with canids identified for the first time in these animals. *S. arieticanis*, *S. bertrami*, *S. capracanis*, and *S. morae* were the most common, each infecting around 40% of the examined mustelids (Attia *et al.*, 2022b; Šneideris *et al.*, 2024; Al-Nabati *et al.*, 2024).

Researchers have extensively studied the structure of the *S. fusiformis* sporocyst wall, which is branching and contains microtubules, as observed in transmission electron microscopy (TEM) studies (Gjerde *et al.*, 2020; El-Morsey and Abdo, 2024). While different studies have reported varying details about *S. fusiformis* merozoites, both light microscopy and TEM have been used to describe these structures. The current study aims to provide a detailed description of *Sarcocystis* cysts, their contents, and the histopathological changes observed in affected organs, alongside genetic analysis of the implicated tissues, such as muscles and hearts.

MATERIALS AND METHODS

Sampling: A post-mortem inspection was conducted on 200 female cattle (6 years old) to detect *S. fusiformis* in the 60 esophagi, 30 hearts, and 60 tongues. The parasite was found in 10% of the animals, and the affected tissues were then sent to a parasitology laboratory, where the white cyst was separated from the skeletal muscles and preserved in two tubes: one with 10% formalin and the other with 2.5% glutaraldehyde. The diseased tissues were subjected to histopathological analysis (using the formalin-preserved tissues), while the remaining tissues were stored in a refrigerator for gene expression studies.

Parasitological examination: The sarcocyst was carefully removed from the surrounding muscle tissue and placed on a clean slide. After air drying, the sarcocyst was fixed with methanol, dehydrated, and stained with Giemsa for 30min. The slide was then rinsed with water, allowed to dry, and examined under an Olympus CX40 optical microscope (Motamedi *et al.*, 2011).

Histopathological examination: Skeletal muscle samples showing visible white cysts were embedded in paraffin wax to create a solid block for cutting. Thin sections (3-4µm thick) of the muscle tissue were then cut from the paraffin block using a microtome. The paraffin was removed from these sections, and they were stained with hematoxylin and eosin (H&E) to highlight the cellular structures for examination under a microscope (Bancroft and Gamble, 2008). Olympus optical microscope supplied with a digital camera was used to inspect the tissue sections (Tokyo, Japan).

Ultrathin preparation: The isolated *Sarcocystis* cysts from skeletal muscles were prepared for transmission electron microscope (TEM). Briefly, the fixed samples were dehydrated using a series of graded ethanol, fixed in




Fig. 1: Life cycle of *Sarcocystis fusiformis* between final host and intermediate host.

a 1% osmium tetroxide solution in Millonig's buffer at a pH of 7.4, penetrated with a mixture of epoxy resin and propylene oxide, encased in absolute resin, and polymerized overnight at 60°C. Ultrathin resin slices of macrocysts were contrasted using lead citrate and uranyl acetate agents; they were then investigated in an 80 kV-operated TEM (JEOL, JEM-1400 TEM) according to (Dubey *et al.*, 2015).

Inflammatory markers: Muscles (masseter muscles), esophagus, and heart parts from *S. fusiformis*-infected cattle were collected in the same manner as those from five young, uninfected cattle calves (negative controls), and stored aseptically in a refrigerator.

DNA extraction: Infected muscle tissue was digested, and the resulting sediment was washed three times with phosphate-buffered saline (PBS) to collect the zoites. These were stored in conical tubes at -20°C (Motamedi *et al.*, 2011). Genomic DNA was then extracted from the positive samples using a phenol-chloroform method (Köchl *et al.*, 2005; Salem *et al.*, 2022a) and stored at -20°C.

RNA isolation: The RNA obtained from 100 mg of tissue under examination was isolated using the total RNA kit (Ambion, Applied Biosystems) following the manufacturer's instructions. The Nano-Drop (Thermo Scientific, USA) was used to quantify and assess the purity of the RNA. Specifically, 500ng mRNA was produced using DNase I amplification grade (Invitrogen) according to the directions provided by the manufacturer. A reverse transcription reaction was conducted on the treated RNA using the High-Capacity cDNA Archive Kit from Applied Biosystems (Yu *et al.*, 2021; Saad *et al.*, 2017).

Quantitative real-time PCR protocol: The PCR program consisted of an initial denaturation step at 94°C for 5 minutes, followed by 30 cycles of denaturation at 94°C for 2 minutes, annealing at 57°C for 30 seconds, Extension at 72°C for 2 minutes, and A final extension step at 72°C for 5 minutes completed the amplification (Hamidinejat *et al.*, 2014; Attia *et al.*, 2022c; Soliman *et al.*, 2022b).

Specific PCR primers were designed to amplify genes for important immune molecules in cattle to study the immune response to the parasite, tumor necrosis factor-

alpha (TNF- α), interleukin-1 β -8, and interferon-gamma (IFN- δ). These primer sequences were based on information from the Gene Bank (Table 1).

Table 1: List of primers used for amplification of pro-inflammatory genes.

Gene	Primers Sequence (F/R)
TNF- α	F- CAAGTAACAAGCCGGTAGCC/ R- TGGAAGACCCCTCCCTGGTA
IL-1 β	F-CTAGCCCATGTGTGCTGAAG/ R- CCTTACTTGGCTCTTCACC
IFN- γ	F- ATAACCAGGTCATTCAAAGG/ -R- AT TCTGACTTCTCTCCGCT
IL-8	F-ATGACTTCCAAGCTGGCTGTT/R- CATGGATCTTGCTTCTCAGC
Beta-actin	F- CGCACCACTGGCATTGTGCAT/ R- TCCAAGGCGACGTAGCAGAG

The gene for beta-actin, a protein involved in cell structure, was used as a reference gene to normalize the expression levels of the immune genes. These genes were analyzed in a separate set of cDNA samples obtained from five parasite-free animals (Yu *et al.*, 2021).

Statistical analysis: The triplicate data are presented as mean \pm SE. The means of the groups (positive and negative PCR-testing cattle) were then statistically compared using the t-test on independent samples. Statistical significance was determined at a p-value of less than 0.05 using SPSS v 26 (IBM, Chicago, IL, USA).

RESULTS

Parasites incidence during macroscopic and microscopic examination: Twenty out of 200 (20/200) examined female cattle (5 years old) were positive for *S. fusiformis* with a prevalence of 10%. *S. fusiformis* cysts were white, ranging from 4–12 mm long and 3–6 mm wide, and they were simple to separate from the host tissue. The cyst contains bradyzoites, which are usually fusiform (broad in the center and tapered ends); some appear to have blunt ends (Fig. 2); the nucleus either centrally or toward one end, and the other end contains granules.

Histopathological examination of the cysts reveals that the macroscopic cysts were detected mostly in skeletal muscles of cattle which indicated the presence of thick-walled sarcocyst containing bradyzoites embedded in muscle with mild tissue reaction as interstitial edema and few eosinophilic inflammatory cells (Fig. 3a&b), other cases showed the presence of large one with thick-walled sarcocyst with severe tissue reaction in the form of eosinophilic inflammatory cells (Fig. 3c &d; Fig. 4), hyaline degeneration of muscle fibers (Fig. 3e), The interstitial tissue was infiltrated mostly by eosinophilic inflammatory cells (Fig. 3f). In some cases, there was also significant infiltration of mononuclear inflammatory cells and muscle degeneration with a large cyst wall (Fig. 3g).

Semi thin ultrastructure of *S. fusiformis*: *Sarcocystis* cyst was composed of a wall with different layers, several cells of metrocytes, and a different compartment. This compartment is separated by septa, filled with bradyzoites in the periphery, leaving the center free from bradyzoites. The bradyzoites appear as a banana shape with an anterior point and a broad posterior (Fig. 5).




Fig. 2: Photomicrograph of cattle *S. fusiformis* showing the banana shape bradyzoites with tapering ends (one end pointed and the other is broad). H&E $\times 40$.




Fig. 3: Photomicrograph of cattle skeletal muscles showing (a) thick-walled sarcocyst containing bradyzoites (long arrow) and interstitial edema (short arrow) (H&EX400), (b) showing interstitial edema and few mononuclear inflammatory cells infiltration (arrow) (H&EX400). (c) large cyst with thick-walled sarcocyst which embedded in muscles (arrows) (H&EX400). (d) showing infiltration with mononuclear inflammatory cells (short arrow) (H&EX400). (e) Hyaline degeneration of skeletal muscle fibers (long arrows) and interstitial mononuclear inflammatory cells infiltration with thick large cyst wall (H&EX400). (f) Infiltration of the interstitial tissue of muscle fibers with esinophils (H&EX400). (g) thick-walled sarcocyst with muscle degeneration (H&EX400).

Ultrathin structure of the *S. fusiformis*: The wall consisted of a layer with a high electron density, measuring up to 50 nm in thickness, which surrounded the outermost parasitophorous vacuolar membrane (ppm)

of the sporocyst wall. Metrocytes had a comparatively lower electron density than bradyzoites. The particles were spherical to elongated in form and were 7–15 μ m in length (Fig. 6). The structures comprised a nucleus,

endoplasmic reticulum, lipid droplets, fissures, and a few numbers to many amylopectin granules. Fig. 5 shows that bradyzoites obtained from 5 *Sarcocysts* were $15.5 (11.5-16.5) \mu\text{m}$ in length and $2.5-4.0\mu\text{m}$ in width when sliced longitudinally.

Transcript levels of the pro-inflammatory cytokines:

The analysis of IL-1 β levels in *S. fusiformis*-infected cattle revealed (8.5 ± 1) levels in muscles, while 9.7 ± 1 in esophagus and 10.5 ± 1 in hearts, which were greater than those of healthy cattle (3.5 ± 0.50 ; Fig. 7). However, IFN- γ levels in *S. fusiformis* -infected cattle were 7 ± 1 in muscles, while in esophagus, 7.5 ± 1.5 and 9.6 ± 1.49 in hearts, which were higher than those in healthy cattle (3 ± 0.55 ; Fig. 8). The analysis of TNF- α levels in *S. fusiformis* -infected cattle revealed (9 ± 1) levels in muscles, while 10.5 ± 1 in esophagus, and 10.0 ± 1 in hearts, which were greater than those of the healthy cattle (3 ± 0.50 ; Fig. 9). The analysis of IL-8 levels in *S. fusiformis* -infected cattle revealed 10 ± 1 in muscles, 12 ± 1 in esophagus, and 95 ± 1

in hearts, which were higher than in healthy cattle (3.5 ± 0.59 ; Fig. 10).

DISCUSSION

In this study, the prevalence of *Sarcocystis* infection in cattle was 10% (Sarafraz *et al.*, 2020). Dubey *et al.* (2023) detected sarcocysts of *S. dubeyi* in 11 (30%) of 35 cattle. Mohamed *et al.* (2020) investigated the prevalence of *Sarcocystis* infection in imported cattle at the Duhok abattoir and evaluated the effectiveness of traditional detection methods. Muscle samples (esophagus, heart, and diaphragm) were collected from 150 cattle from three different countries. A high percentage of the samples were infected with *Sarcocystis*: 94% of the esophagus, 92% of the diaphragm, and 41.3% of heart samples. This study highlights the common occurrence of *Sarcocystis* in imported cattle and emphasizes the need for further epidemiological investigations to assess the extent of the infection within the country.




Fig. 4: Photomicrograph of cattle skeletal muscles showing (a) thick-walled sarcocyst containing bradyzoites (long arrow) (H&EX400), (b) showing degenerated muscle fibers with the wall of the cyst and banana-shaped bradyzoites (H&EX400). (c) large cyst showing the compartment of the cyst (H&EX400). (d) Higher magnification of the cyst in the center shows the trabeculae, which differentiate the cyst into compartments; (short arrow) (H&EX400).




Fig. 5: Semi-thin section of *Sarcocystis fusiformis* showing the cyst wall; metrocyte and merozoites. **A:** *Sarcocystis* cyst was composed of a wall (w) with different layers, several cells of metrocytes, and a different compartment. **B:** compartment in the cyst separated by septa, which filled with merozoites. **C:** The metrocyte and compartment of the cyst are closely packed with merozoites. **D:** the center of the cyst appears nearly free from merozoites.




Fig. 6: TEM study of *Sarcocystis* cyst showing several bradyzoites which were globular to oblong in shape; **A:** They contained a nucleus (N), endoplasmic reticulum, lipid droplets (ld), a few to several amylopectin granules (ap) and golgi apparatus (GA).




Fig. 7: Transcript levels of IL1- β in different organs infected with *S. fusiformis* in cattle. T1-T3 were control samples of muscle, tongue, and heart, while T4-T6 were *S. fusiformis*-infected muscle, tongue, and heart samples, respectively.




Fig. 8: Transcript levels of IFN- γ in different organs infected with *S. fusiformis* in cattle. T1-T3 were control samples of muscle, tongue, and heart, while T4-T6 were *S. fusiformis*-infected muscle, tongue, and heart samples, respectively.




Fig. 9: Transcript levels of TNF- α in different organs infected with *S. fusiformis* in cattle. T1-T3 were control samples of muscle, tongue, and heart, while T4-T6 were *S. fusiformis*-infected muscle, tongue, and heart samples, respectively.




Fig. 10: Transcript levels of IL-8 in different organs infected with *S. fusiformis* in cattle. T1-T3 were control samples of muscle, tongue, and heart, while T4-T6 were *S. fusiformis*-infected muscle, tongue, and heart samples, respectively.

Based on the assessed incidence of sarcocystosis in this study, Egypt has several appropriate conclusive and intermediate hosts for various parasite species (El-Morsey *et al.*, 2021). The variation in the prevalence percentage between the other studies may be attributed to the differences in sampling time, localities, hygienic measures applied on the various farms, and preventive medication and control strategies applied on each farm (Belina *et al.*, 2021; Alsulami *et al.*, 2023; Alsulami *et al.*, 2024). It is noteworthy that the animals afflicted with the macroscopic sarcocyst forms also contracted the microscopic ones suggests that they had previously come into contact with both canines and felines that were infected (Ashmawy *et al.*, 2014; Idland *et al.*, 2022).

Previous studies can distinguish between different types of *Sarcocystis* by examining the microscopic structure of the parasite's cyst wall. This detailed analysis involves using powerful imaging techniques like TEM and histological staining (El-Kady *et al.*, 2018; Swar and Shnawa, 2020). Microscopic examination of infected muscle tissue showed the presence of *Sarcocystis* cysts with thick walls containing numerous merozoites (Morsy *et al.*, 2018; Salem *et al.*, 2022b). The surrounding muscle tissue exhibited signs of damage and inflammation, including interstitial edema as fluid buildup between muscle fibers and hyaline degeneration as the change in muscle fibers where they appear glassy or translucent. Inflammatory cell infiltration is the presence of immune cells, predominantly eosinophils, in the spaces between

muscle fibers. Some cases also displayed a significant presence of mononuclear inflammatory cells, indicating that the parasite triggers a significant inflammatory response in the affected muscle tissue (Faghiri *et al.*, 2019). Our results agreed with those recorded by Farhangpazhouh *et al.* (2020) and Shams *et al.* (2022) reported similar histopathological changes in infected cattle tissues with *Sarcocystis* infection in Iran.

The immune response to sarcocystosis is species-specific, and the host protection against one species of the parasite cannot provide immunity against heterologous challenges (Hettiarachchi and Rajapakse, 2008; Florin-Christensen *et al.*, 2021). Cell-mediated and humoral immunity can be developed after immunizing the intermediate hosts against crude antigens isolated from cystozoites and merozoites of *Sarcocystis* species (Salem *et al.*, 2022c; Mousa *et al.*, 2024). In cell-mediated immunity, immune cells such as macrophages and lymphocytes migrate and disseminate through visceral and muscular tissues (El Shanawany *et al.*, 2019; Delgado and Lennon-Duménil, 2022).

The humoral immune response usually appears after 30 days of infection. Cell-mediated immunity in Sarcocystosis is more significant than humoral immunity (Hussein *et al.*, 2025). Skin or delayed-type hypersensitivity tests enable crude antigen stimulation of cell-mediated immunity response to the specific antigen injected intradermally (Hussein *et al.*, 2025).

This study explores how *Sarcocystis* infection affects different tissues in livestock, by analyzing microscopic

changes in infected tissues and examining the roles of various immune system molecules (cytokines and chemokines) to develop effective disease prevention strategies. Consistent with our study Teufel *et al.* (2022) focused on understanding the activity of IL-1 cytokines, which are important in regulating the body's inflammatory response and natural defenses against infection. This knowledge could lead to new treatments using recombinant cytokines to combat *Sarcocystis* infection. While IL-1Ra binds to IL-1RI to have a potent anti-inflammatory impact, IL-1 α and IL-1 β are potent inducers of inflammation (Mingala *et al.*, 2007; Makaremi *et al.*, 2022). The IL-1 family members that have been investigated the most are IL-1 α and IL-1 β (Bhadaniya *et al.*, 2019). IL-1 α is typically connected to the plasma membrane of several cell types (Ezzat Alnakip *et al.*, 2014; Pyriellou *et al.*, 2020). Only in its fully developed state does IL-1 β function, and this protein is released, circulates throughout the body, and functions as a paracrine factor (Cavalli *et al.*, 2021).

The current study agreed with El Shanawany *et al.* (2019), who stated that older female cattle were much more likely to be infected (90.7%) with *Sarcocystis* than young males (20%). Analysis of the immune response showed a shift towards a Th-2 type response in infected animals, with suppression of pro-inflammatory signals: Levels of the Th-1 cytokine IFN- γ , which normally helps activate a strong immune response, were reduced. Increase in anti-inflammatory signals: Levels of the Th-2 cytokine IL-5, which is associated with suppressing inflammation, were elevated. Antibody production: Higher levels of IgG and IgE antibodies were also found in infected animals. At the infection site, the immune response involved a mix of inflammatory cells, granuloma formation, eosinophil infiltration, fluid buildup (edema), and tissue death (necrosis). This indicates a complex immune reaction to the parasite.

In the current study, the analysis of IL-1 β levels in *S. fusiformis* -infected cattle revealed (8.5 ± 1) levels in muscles, while 9.7 ± 1 in the esophagus and 10.5 ± 1 in hearts, which were greater than those of the healthy cattle (3.5 ± 0.50) however, IFN- γ levels in *S. fusiformis* -infected cattle were 7 ± 1 in muscles, while in esophagus 7.5 ± 1.5 and 9.6 ± 1.49 in hearts, The reported values were greater than those of healthy cattle (3 ± 0.55), also, the analysis of TNF- α levels in *S. fusiformis* -infected cattle revealed (9 ± 1) levels in muscles, while 10.5 ± 1 in the esophagus, and 10.0 ± 1 in hearts compared to control (3 ± 0.50), and the analysis of IL-8 levels in *S. fusiformis* -infected cattle revealed 10 ± 1 in muscles and 12 ± 1 in the esophagus, and 95 ± 1 in hearts, compared to control animals (3.5 ± 0.59). There were different studies in concur with our observation, such as the parallel research by (Aboelenin *et al.*, 2017), who noticed that TNF α is a cytokine signaling protein crucial for regulating the immune system and influencing female reproductive function and its pathogen response. Also, (Abou Mossallam *et al.*, 2015) found that the inflammatory cytokines released during infection are endometriosis, which is prompted to release cytokines and chemokines, such as IL-8. This attracts and activates neutrophils and monocytes to clear germs and prevent illness. Dini *et al.*

(2023) reported the first case of bovine eosinophilic myositis (BEM) in a beef cow where *Toxoplasma gondii* DNA was detected alongside *S. hominis*. This finding suggests that co-infection with these two parasites may contribute to the development of BEM, a muscle disease that causes significant economic losses and potential public health concerns. IL-8 was recognized as the first member of the chemokine superfamily. Comparative analysis of IL-8 expression was conducted in Egyptian buffalo with and without endometritis (Abou Mossallam *et al.*, 2015; Farghali *et al.*, 2022).

The expression of IL8 was examined in the previous study in Egyptian buffalo's ovaries, uteruses, and mammary glands, both with and without endometritis (Raheel *et al.*, 2020). Only the uteruses of buffalo with endometritis exhibited a considerable increase (26.6-fold) in IL8 expression compared to buffalo without endometritis (Elsayed *et al.*, 2020; Taha *et al.*, 2021).

Prior research on the key cytokine genes in the buffalo immune system, the conservation of this high-level sequence (Abdallah *et al.*, 2021; Yehia *et al.*, 2023). For the host to defend itself against pathogenic invasion, IL-12 must be expressed early. The transcription of the p40 chain regulates the secretion of the physiologically active heterodimer, or p70. Non-mammalian p40, such as avian (Zhang and Su, 2023) and Fugu p40 (Yoshiura *et al.*, 2003), has been shown to exhibit constitutive expression in tissues of non-lymphoid origin, such as the kidney, brain, heart, and intestine. Constitutive expression of buffalo p35 was seen in both lymphoid tissues, such as the spleen, and non-lymphoid tissues, such as the brain. Exposure to polyclonal mitogens in lymphoid cells leads to immunological activation, which in turn promotes the expression of IL-12 p40 (Lima-Junior *et al.*, 2022; McDaniel *et al.*, 2022).

One powerful inducer of the acute phase response is IL-6. Quick IL6 synthesis aids in tissue damage and infection response on the part of the host, but overproduction of IL6 leads to disease pathology (Sreekumar *et al.*, 2002). Upon recognition of pathogens by toll-like receptors (TLRs) at the infection site or tissue damage, myeloid cells, including macrophages and dendritic cells, initiate a portion of the innate immune response (Ni and Chen, 2023). The transcript levels of different genes were upgraded in different tissues according to inflammatory reactions.

Conclusions: This study found a 10% prevalence of *S. fusiformis* in cattle. The parasite's ultrastructure was characterized, and its impact on host tissues, including upregulation of inflammatory cytokines and histopathological changes, was documented. Advanced molecular studies are necessary to determine the relationship between *Sarcocystis* species infecting buffalo and cattle in this region. Considering the economic importance of cattle, more research is crucial to fully grasp the effects of *Sarcocystis* infection. This includes deeper investigation into the parasite's biology, life cycle, and variations. Furthermore, controlling the movement of canine and feline hosts on farms should be explored as a potential method for mitigating the impact of *Sarcocystis*.

Competing interests: The authors declare that they have no competing interests.

Acknowledgements: The authors gratefully acknowledge Princess Nourah bint Abdulrahman University Researchers Supporting Project number (PNURSP2024R459), Princess Nourah bint Abdulrahman University, Riyadh, Saudi Arabia. The authors extend their appreciation to the Deanship of Research and Graduate Studies at King Khalid University for funding this work through Large Research Project under grant number (R.G.P2/478/45).

Authors contributions: Conceptualization, SMAB, MAA, HSAR, AMA, and SAK, formal analysis, HAM, MMAQ, NAH, MMA, and MMA, investigation, SMAB, MAA, HSAR, AMA, and SAK, data curation, HAM, MMAQ, NAH, MMA, and MMA, writing original draft preparation, SMAB, MAA, HSAR, AMA, and SAK, writing final manuscript and editing, HAM, MMAQ, NAH, MMA, and MMA, visualization and methodology, SMAB, MAA, HSAR, AMA, SAK, HAM, MMAQ, NAH, MMA, and MMA. All authors have read and agreed to the published version of the manuscript.

REFERENCES

- Abdallah F, Coindre S, Gardet M, et al., 2021. Leukocyte immunoglobulin-like receptors in regulating the immune response in infectious diseases: A window of opportunity to pathogen persistence and a sound target in therapeutics. *Front Immunol* 12:717998.
- Aboelenin M, Mahrous K, Rashed M, et al., 2017. Molecular characterization of tumor necrosis factor- α (tnfa) gene in egyptian river buffaloes. *Arab Univ J Agric Sci* 25:367-375.
- Abou Mossallam AA, El Nahas SM, Mahfouz ER, et al., 2015. Characterization of buffalo interleukin 8 (IL-8) and its expression in endometritis. *J Genet Eng Biotechnol* 13:71-77.
- Ahmed AM, Elshraway NT, Youssef AI, 2016. Survey on *Sarcocystis* in bovine carcasses slaughtered at the municipal abattoir of El-Kharga, Egypt. *Vet World* 9:1461.
- Al-Nabati EA, Almahallawi R, Alzahrani AM, et al., 2024. Estimation of in vitro acaricidal activities of ethanolic and ethyl acetate extracts of *Nicotiana tabacum* against *Hyalomma* species of livestock. *Pak Vet J* 44(3): 939-943.
- Attia MM, Alzahrani AM, Hanna MI, et al., 2022a. The biological activity of *Illicium verum* (Star Anise) on *Lernaea cyprinacea*-Infested *Carassius auratus* (Goldfish): In Vivo study. *Life* 12(12): 2054.
- Attia MM, Soliman SM, Salaeh NM, et al., 2022c. Evaluation of immune responses and oxidative stress in donkeys: Immunological studies provoked by *Parascaris equorum* infection. *Saudi J Biol Sci* 29(4): 2173-2179.
- Attia MM, Yehia N, Soliman MM, et al., 2022b. Evaluation of the antiparasitic activity of the chitosan-silver nanocomposites in the treatment of experimentally infested pigeons with *Pseudolynchia canariensis*. *Saudi J Biol Sci* 29(3): 1644-1652.
- Ashmawy KI, Abu-Akkada SS, Ghashir MB, 2014. Prevalence and molecular characterization of *Sarcocystis* species in water buffaloes (*Bubalus bubalis*) in Egypt. *Trop Anim Health Prod* 46:1351-1356.
- Badr A, EL-Malek A, Ashraf M, Arafa M. 2024. Prevalence of *Sarcocystis fusiformis* in slaughtered buffaloes in assiut abattoir, and study the effect of chilling on their viability and infectivity. *Assiut Vet Med J* 70:86-97.
- Bancroft JD, Gamble M, 2008. Theory and practice of histological techniques: Elsevier health sciences.
- Barakat AZ, Abdel-Aty AM, Ibrahim MK, et al., 2023. Purification and characterization of cysteine protease of *Sarcocystis fusiformis* from infected Egyptian water buffaloes. *Sci Rep* 13:16123.
- Belina D, Hailu Y, Gobena T, et al., 2021. Prevalence and epidemiological distribution of selected foodborne pathogens in human and different environmental samples in Ethiopia: a systematic review and meta-analysis. *One health outlook* 3:19.
- Bhadaniya A, Prasad M, Savsani H, et al., 2019. Pro-inflammatory cytokine expression studies of subclinical and clinical endometritis in endometrial tissues of buffaloes. *Trop Anim Health Prod* 51:1161-1166.
- Castro-Forero S, Bulla-Castañeda D, Buitrago HL, et al., 2022. *Sarcocystis* spp., a parasite with zoonotic potential. *Bulg J Vet Med* 25:175-186.
- Cavalli G, Colafrancesco S, Emmi G, et al., 2021. Interleukin 1 α : a comprehensive review on the role of IL-1 α in the pathogenesis and treatment of autoimmune and inflammatory diseases. *Autoimmun Rev* 20:102763.
- Constable PD, Hinchcliff KW, Done SH, et al., 2016. Veterinary medicine: a textbook of the diseases of cattle, horses, sheep, pigs and goats: Elsevier Health Sciences.
- Cordeiro M, Mengistu G, Pogue S, et al., 2022. Assessing feed security for beef production within livestock-intensive regions. *Agric Syst* 196:103348.
- Delgado MG, Lennon-Duménil A-M, 2022. How cell migration helps immune sentinels. *Front Cell Dev Biol* 10:932472.
- Dini FM, Caffara M, Jacinto JG, et al., 2023. A case of bovine eosinophilic myositis (BEM) associated with co-infection by *Sarcocystis hominis* and *Toxoplasma gondii*. *Animals* 13(2): 311.
- Dubey J, Fayer R, Rosenthal B, et al., 2014. Identity of *Sarcocystis* species of the water buffalo (*Bubalus bubalis*) and cattle (*Bos taurus*) and the suppression of *Sarcocystis sinensis* as a nomen nudum. *Vet Parasitol* 205:1-6.
- Dubey J, Gupta A, de Araujo LS, et al., 2023. *Sarcocystis cruzi* (Hasselmann, 1923) Wenyon, 1926: redescription, molecular characterization and deposition of life cycle stages specimens in the Smithsonian museum. *Parasitology* 150:1192-1206.
- Dubey J, Rosenthal B, 2023. Bovine sarcocystosis: *Sarcocystis* species, diagnosis, prevalence, economic and public health considerations, and association of *Sarcocystis* species with eosinophilic myositis in cattle. *Int J Parasitol* 53:463-475.
- Dubey JP, Hilali M, Van Wilpe E, et al., 2015. Redescription of *Sarcocystis fusiformis* sarcocysts from the water buffalo (*Bubalus bubalis*). *Parasitology* 142:385-394.
- Dubey JP, Sykes JE, 2021. Sarcocystosis. In: Greene's Infectious Diseases of the Dog and Cat: Elsevier. p 1172-1178.
- El-Kady AM, Hussein NM, Hassan AA, 2018. First molecular characterization of *Sarcocystis* spp. in cattle in Qena Governorate, Upper Egypt. *J Parasitic Dis* 42:114-121.
- El-Mahdi MB, Rabie SA, Hassanine RMES, et al., 2023. Molecular identification, pathogenesis, and life cycle of *Sarcocystis cruzi* from Cattle (*Bos taurus*) in New Valley Governorate, Egypt. *J Parasitol Res* 2023(1), 7829290.
- El-Morsey A, Abdo W, 2024. Recent insights into the morphology, molecular characterization and tissue localization of the caprine *Sarcocystis* species infecting domestic goats (*Capra hircus*): *Sarcocystis moulei*, *Sarcocystis capracanis*, and *Sarcocystis hircanicis*. *Parasitol Res* 123:55.
- El-Morsey A, Abdo W, Zaid AAA, et al., 2021. Morphologic and molecular identification of three macroscopic *Sarcocystis* species infecting domestic sheep (*Ovis aries*) and cattle (*Bos taurus*) in Egypt. *Parasitol Res* 120:637-654.
- Alsulami MN, & El-Saadony MT, 2024. The enhancing effect of bacterial zinc nanoparticles on performance, immune response, and microbial load of Nile Tilapia (*Oreochromis niloticus*) by reducing the infection by *Trichodina heterodontata*. *Pak Vet J* 2024, 44(3): 599-610.
- Alsulami MN, & El-Saadony MT, 2023. Supplementing broiler diets with bacterial selenium nanoparticles enhancing performance, carcass traits, blood indices, antioxidant status, and caecal microbiota of *Eimeria tenella*-infected broiler chickens. *Poult Sci* 102(12):103111.
- El Shanawany EE, Nassar SA, Ata EB, 2019. Detection of humoral and cellular immune responses in buffaloes naturally infected with sarcocystosis with risk factor assessment. *Acta Vet* 69:275-289.
- Elsayed DH, El-Azzazi FE, Mahmoud YK, et al., 2020. Subclinical endometritis and postpartum ovarian resumption in respect to TNF- α , IL-8 and CRP in Egyptian buffaloes. *Anim Reprod* 17:e20190027.
- Ezzat Alnakiq M, Quintela-Baluja M, Böhme K, et al., 2014. The immunology of mammary gland of dairy ruminants between healthy and inflammatory conditions. *J Vet Med* 2014:659801.
- Faghiri E, Davari A, Nabavi R, 2019. Histopathological survey on *Sarcocystis* species infection in slaughtered cattle of zabol-iran/zabol-iran-da kesilen sigirlarda *Sarcocystis* turlerinin yol actigi enfeksiyonlar uzerine histopatolojik inceleme. *Turkish J Parasitol* 43(4): 182-187.

- Farghali HA, AbdElKader NA, Fathi M, et al., 2022. The efficiency of intrauterine infusion of platelet-rich plasma in the treatment of acute endometritis as assessed by endoscopic, Doppler, oxidative, immunohistochemical, and gene expression alterations in jennies. *Theriogenology* 181:147-160.
- Farhangpazhouh F, Yakhchali M, Farshid AA, et al., 2020. Prevalence and pathologic changes due to *Sarcocystis* species in naturally infected sheep in Urmia city, Iran. *J Zoonotic Dis* 4:54-60.
- Florin-Christensen M, Schnittger L, Bastos RG, et al., 2021. Pursuing effective vaccines against cattle diseases caused by apicomplexan protozoa. *CABI Reviews* 16: 1-24.
- Gjerde B, De la Fuente C, Alunda JM, et al., 2020. Molecular characterization of five *Sarcocystis* species in domestic sheep (*Ovis aries*) from Spain. *Parasitol Res* 119:215-231.
- Hamidinejat H, Moetamedi H, Alborzi A, et al., 2014. Molecular detection of *Sarcocystis* species in slaughtered sheep by PCR-RFLP from south-western of Iran. *J Parasit Dis* 38(2): 233-237.
- Hettiarachchi D, Rajapakse R, 2008. Antigenic analysis of bovine *Sarcocystis* spp. in Sri Lanka. *J Natn Sci Foundation Sri Lanka* 36 (3):239-244
- Hussein D, Abu-Akkada S, Bessat M, et al., 2017. Molecular identification of *Sarcocystis* species in imported frozen beef in Egypt. *Alex J Vet Sci* 53(2):72.
- Hussein SN, Ibrahim AA, Shukur MS, 2025. Immunization of rabbits against whole crude antigen of sarcocystosis (*S. gigantea*) isolated from macrocystis of naturally infected sheep in duhok province, Iraq. *Egyptian J Vet Sci* 56: 1289-1299.
- Idland L, Granquist EG, Aspholm M, et al., 2022. The prevalence of *Campylobacter* spp., *Listeria monocytogenes* and Shiga toxin-producing *Escherichia coli* in Norwegian dairy cattle farms: A comparison between free stall and tie stall housing systems. *J Appl Microbiol* 132:3959-3972.
- Kinley RD, Martinez-Fernandez G, Matthews MK, et al., 2020. Mitigating the carbon footprint and improving productivity of ruminant livestock agriculture using a red seaweed. *J Clean Prod* 259:120836.
- Köchl S, Niederstätter H, Parson W, 2023. DNA extraction and quantitation of forensic samples using the phenol-chloroform method and Real-Time PCR, 2005, Springer, Berlin, Germany, 13-29.
- Lima-Junior DS, Collins N, Han S-J, 2022. Spleen, lymph nodes, and lymphoid tissues—Inflammatory system. In: *Macrophages in the human body*: Elsevier. p 67-105.
- Lovarelli D, Bacenetti J, Guarino M, 2020. A review on dairy cattle farming: Is precision livestock farming the compromise for an environmental, economic and social sustainable production? *J Clean Prod* 262:121409.
- Makaremi S, Asgarzadeh A, Kianfar H, et al., 2022. The role of IL-1 family of cytokines and receptors in pathogenesis of COVID-19. *Inflamm Res* 71:923-947.
- McDaniel MM, Chawla AS, Jain A, et al., 2022. Effector memory CD4+ T cells induce damaging innate inflammation and autoimmune pathology by engaging CD40 and TNFR on myeloid cells. *Sci Immunol* 7:eabk0182.
- Mendoza Roldan JA, Otranto D, 2023. Zoonotic parasites associated with predation by dogs and cats. *Parasites & Vectors* 16:55.
- Menshawry SM, Essa BH, Shaaban SI, et al., 2023. Prevalence and genetic characterization of *Sarcocystis fusiformis* in water buffaloes (*Bubalus bubalis*) in two northern provinces of Egypt. *Slov Vet Res* 60 (Suppl 25): 195-203
- Mingala CN, Odbileg R, Konnai S, et al., 2007. Molecular cloning, sequencing and phylogenetic analysis of inflammatory cytokines of swamp type buffalo contrasting with other bubaline breeds. *Comp Immunol Microbiol Infect Dis* 30:119-131.
- Mohamed T, Hussein S, Shukur M, et al., 2020. Survey on *Sarcocystis* infection in imported male cattle carcasses slaughtered at Duhok abattoir, Kurdistan region of Iraq. *Microb Biotechnol* 5(1): 128-134.
- Morsy K, Abdel-Ghaffar F, Bin Dajem S, et al., 2018. First molecular characterization and morphological aspects of *Sarcocystis fusiformis* infecting water buffalo *Bubalus bubalis* in Egypt. *Acta Parasitol* 63:333-345.
- Motamedi GR, Dalimi A, Nouri A, et al., 2011. Ultrastructural and molecular characterization of *Sarcocystis* isolated from camel (*Camelus dromedarius*) in Iran. *Parasitol Res* 108 4: 949-954.
- Mottet A, de Haan C, Falcucci A, et al., 2017. Livestock: On our plates or eating at our table? A new analysis of the feed/food debate. *Glob Food Secur* 14:1-8.
- Mousa MM, El Sokkary MY, Hamouda Abd El Naby WS, et al., 2021. The prevalence of *Sarcocystis* affecting slaughtered cattle and buffalo at sirselian abattoir in Egypt. *Alex J Vet Sci* 69(2): 49-58.
- Mousa MM, Hiekal FA, El-Hoshay SM, et al., 2016. Characterization of *Sarcocystis* species based on traditional and molecular methods in imported frozen buffalo meat in Egypt. *Alex J Vet Sci* 51(1): 155-161.
- Mousa MR, Attia MM, Salem, HM, et al., 2024. Coinfection of the gut with protozoal and metazoal parasites in broiler and laying chickens. *Poult Sci* 103(1):103227.
- Ni H, Chen Y, 2023. Differentiation, regulation and function of regulatory T cells in non-lymphoid tissues and tumors. *Int Immunopharmacol* 121:110429.
- Pal M, Tolawak D, Garedaghi Y, 2023. A comprehensive review on major zoonotic parasites from dogs and cats. *Int J Med Parasitol Epidemiol Sci* 4(1):3-11.
- Palange A, Dhatrak P, 2021. Lean manufacturing a vital tool to enhance productivity in manufacturing. *Materials Today: Proceedings* 46:729-736.
- Pyrillou K, Burzynski LC, Clarke MC, 2020. Alternative pathways of IL-1 activation, and its role in health and disease. *Front Immunol* 11:613170.
- Raheel IAER, Hassan WH, Salem SSR, et al., 2020. Biofilm forming potentiality of *Escherichia coli* isolated from bovine endometritis and their antibiotic resistance profiles. *J Adv Vet Anim Res* 7:442.
- Ras RA, 2021. Prevalence and molecular identification of *Sarcocystis* spp. infecting water buffaloes (*Bubalus bubalis*) in Sharkia Province, Egypt. *Egyptian Vet Med Soc Parasitol J* 17:1-19.
- Saad AM, Samy A, Soliman MA, et al., 2017. Genotypic and pathogenic characterization of genotype VII Newcastle disease viruses isolated from commercial farms in Egypt and evaluation of heterologous antibody responses. *Arc. Virol.*, 162(7), 1985-1994.
- Salem HM., Khattab MS, Yehia N, et al., 2022a. Morphological and molecular characterization of *Ascaridia columbae* in the domestic pigeon (*Columba livia domestica*) and the assessment of its immunological responses. *Poult Sci* 101(2): 101596.
- Salem HM, Salem MA, Soliman MM, et al., 2022b. Parasitological and histopathological examination of cocktail lovebirds infected with *Eimeria aratinga* (Apicomplexa: Eimeriidae). *Poult Sci* 101(5):101781.
- Salem HM, Salaeh NM, Ragni M, et al., 2022c. Incidence of gastrointestinal parasites in pigeons with an assessment of the nematocidal activity of chitosan nanoparticles against *Ascaridia columbae*. *Poult Sci* 101(6): 101820.
- Sarafraz N, Spotin A, Haniloo A, et al., 2020. Prevalence and molecular analysis of *Sarcocystis* infections in cattle in Northwest Iran and the first global report of *S. gigantea* in cattle. *Comp Immunol Microbiol Infect Dis* 73: 101566.
- Shams M, Shamsi L, Asghari A, et al., 2022. Molecular epidemiology, species distribution, and zoonotic importance of the neglected meat-borne pathogen *Sarcocystis* spp. in cattle (*Bos taurus*): a global systematic review and meta-analysis. *Acta Parasitol* 67:1055-1072.
- Šneideris D, Moskaliova D, Butkauskas D, et al., 2024. The distribution of *Sarcocystis* species described by ungulates-canids life cycle in intestines of small predators of the family Mustelidae. *Acta Parasitol* 69(1): 747-758.
- Soliman SM, Attia MM, Al-Harbi MS, et al., 2022a. Low host specificity of *Hippobosca equina* infestation in different domestic animals and pigeon. *Saudi J Biol Sci* 29:2112-2120.
- Soliman SM, Salem HM, Saad AM, et al., 2024. *Haemonchus contortus* infection of goats and the use of anthelmintic natural alternative: An updated review. *J Hellenic Vet Med Soc* 75(2): 7201-7210.
- Soliman SM, Aljhdali NH, Majrashi KA, et al., 2022b. Immune cell reaction associated with *Coenurus cerebralis* infection in sheep with particular reference to ELISA as a diagnostic tool. *Life* 12(10):1515.
- Sreekumar E, Premraj A, Saravanakumar M, et al., 2002. Buffalo (*Bubalus bubalis*) interleukin-2: sequence analysis reveals high nucleotide and amino acid identity with interleukin-2 of cattle and other ruminants. *Eur J Immunogenet* 29:341-345.
- Strydom T, Lavan RP, Torres S, et al., 2023. The economic impact of parasitism from nematodes, trematodes and ticks on beef cattle production. *Animals* 13:1599.
- Swar SO, Shnawa BH, 2020. Ultrastructural and molecular characterization of *Sarcocystis* species derived from macroscopic sarcocysts of domestic sheep and goats in Soran City, Erbil, Iraq. *World's Vet J* 10(4): 540-550.
- Taha DA, Mahfouz ER, Bibars MA, et al., 2021. Cytokine genes expression in uteri of *Bubalus bubalis* associated with endometritis infection. *Jordan J Biol Sci* 14(2): 245 - 251.

- Tenu F, Şerban A, Donosă R-E, *et al.*, 2023. Study regarding the beef meat production evolution worldwide and national level. *Scientific Papers. Series D. Animal Science*, 1(2285-5750): 352-362.
- Terry SA, Basarab JA, Guan LL, *et al.*, 2020. Strategies to improve the efficiency of beef cattle production. *Can J Anim Sci* 101:1-19.
- Teufel LU, Arts RJ, Netea MG, *et al.*, 2022. IL-1 family cytokines as drivers and inhibitors of trained immunity. *Cytokine* 150:155773.
- Thomas D, Beletse Y, Dominik S, *et al.*, 2021. Net protein contribution and enteric methane production of pasture and grain-finished beef cattle supply chains. *Animal* 15:100392.
- Thornton P, Nelson G, Mayberry D, *et al.*, 2022. Impacts of heat stress on global cattle production during the 21st century: a modelling study. *The Lancet Planetary Health* 6:e192-e201.
- Valizadeh H, Ebrahimi A, 2022. A study on the abundance of infection with *Sarcocystis* in goats slaughtered in the Tabriz slaughterhouse. *J Food Safe Hyg* 7(3):164-170.
- Wielgus E, Caron A, Bennitt E, *et al.*, 2021. Inter-group social behavior, contact patterns and risk for pathogen transmission in cape buffalo populations. *J Wildl Manag* 85:1574-1590.
- Yehia N, Salem HM., Mahmmud Y, Said D, *et al.* 2023. Common viral and bacterial avian respiratory infections: an updated review. *Poult. Sci.*, 102(5), 102553.
- Yoshiura Y, Kiryu I, Fujiwara A, *et al.*, 2003. Identification and characterization of Fugu orthologues of mammalian interleukin-12 subunits. *Immunogenetics* 55:296-306.
- Yu HB, Wang WJ, Tang S, *et al.*, 2021. Immune responses and pathogenesis in persistently PCR-positive patients with SARS-CoV-2 infection. *J Med Virol* 93(2): 760-765.
- Zhang Y, Su J, 2023. Interleukin-2 family cytokines: An overview of genes, expression, signaling and functional roles in teleost. *Dev Comp Immunol* 141:104645.

IB-Press