



Reference Values of Oxidative Stress Parameters in Adult Iranian Fat-Tailed Sheep

S. Nazifi*, N. Ghafari¹, F. Farshneshani, M. Rahsepar and S. M. Razavi¹

Department of Clinical Studies; ¹Department of Pathobiology, School of Veterinary Medicine, Shiraz University, Shiraz, Iran

*Corresponding author: nazifi@shirazu.ac.ir

ARTICLE HISTORY

Received: August 08, 2009

Revised: September 10, 2009

Accepted: October 24, 2009

Key words:

Malondialdehyde

Superoxide dismutase

Catalase

Glutathione peroxidase

Reference values

Iranian fat-tailed sheep

ABSTRACT

The present study was performed on 111 adult clinically healthy Iranian fat-tailed sheep from both sexes (28 male and 83 female). Blood concentrations of oxidative stress parameters viz. malondialdehyde (MDA), superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPX) were determined. The reference values for oxidative stress parameters of adult clinically healthy Iranian fat-tailed sheep were as followed: MDA 0.53-0.60 $\mu\text{mol/l}$, SOD 948.65-1011.50 U/gHb, CAT 1834.29-1915.63 U/gHb and GPX 191.67-196.52 U/gHb. There were no significant differences in oxidative stress parameters among animal of the two sexes.

©2010 PVJ. All rights reserved

To cite this article: Nazifi S, N Ghafari, F Farshneshani, M Rahsepar and SM Razavi, 2010. Reference values of oxidative stress parameters in adult Iranian fat-tailed sheep. *Pakistan Vet J*, 30(1): 13-16.

INTRODUCTION

A stressful condition leads to the excessive production of the radicals, which results in oxidative stress, an imbalance in the oxidant/antioxidant system (Khadija *et al.*, 2009). Generation of free radicals is an integral feature of normal cellular function. In contrast, excessive generation and/or inadequate removal of free radicals results in destructive and irreversible damage to the cell (Lopaczyski and Zeisel, 2001). Reactive oxygen species (ROS) including superoxide radical, hydrogen peroxide and hydroxyl radical have a great impact on the normal function of biomolecules like nucleic acids, proteins and cell membrane phospholipids. Free radicals are generated during stepwise reduction of molecular oxygen (Singh *et al.*, 1999). Hallwell and Gutteridge (1999) described several lines of defense against reactive oxygen species in animals.

Enzymes with important antioxidant functions include: i) superoxide dismutase (SOD), which catalyses the dismutation of superoxide radical to hydrogen peroxide and water, ii) catalase (CAT), which catalyses the breakdown of hydrogen peroxide to oxygen and water, and iii) glutathione peroxidase (GPX), which facilitates the destruction of both hydrogen peroxide and organic peroxides. Reduced glutathione (GSH), a tri-peptide thiol, is an important antioxidant, as well as a co-factor for various antioxidant enzymes (Kidd, 1997). SOD is the first line of defense against ROS and is active in catalyzing detoxification of superoxide radical (Gonzales *et al.*, 1984). The hydrogen peroxide generated in this

reaction is restored to water in the presence of CAT and GPX. Polyunsaturated fatty acids present in membrane phospholipids are the main target substrates for oxygen radical activity which results in disorganization of cell framework and function (Patterson and Leacke, 1998). Lipid peroxidation is an indicator of oxidative stress in cells and tissues. Lipid peroxides derived from polyunsaturated fatty acids are unstable and are decomposed to form a series of compounds, including malondialdehyde (MDA). The quantization of MDA is widely used as an indicator of lipid peroxidation (Simsek *et al.*, 2006).

Increased levels of lipid peroxidation products such as MDA have been reported in a variety of diseases like *Dicrocoelium dendriticum* infection in sheep (Simsek *et al.*, 2006) and kidney diseases in dogs (Kargin and Fidanci, 2001). The brain injury is reported to be caused by superoxide radical and hydrogen peroxide (Kotos and Wel, 1986). Distomatosis (*Fasciola hepatica*, *Fasciola gigantica* and *Dicrocoelium dendriticum* infections) in sheep causes production of reactive oxygen species and lipid peroxidation by significant increase in liver MDA (Deger *et al.*, 2008).

There is little information concerning the oxidative stress enzymes in sheep. Comparative aspects of plasma antioxidant status in sheep and goats, and the influence of experimental abomasal nematode infection were investigated by Lightbody *et al.* (2001). Also, Kizil *et al.* (2007) reported oxidative stress and antioxidant status in goats naturally infected with *Mycoplasma agalactiae*. The aim of this study was to present the reference values

of oxidative stress parameters in adult clinically healthy Iranian fat-tailed sheep. Such information would allow us to make comparisons between normal and abnormal states and provide a better understanding in diseases accompanied by oxidative stress.

MATERIALS AND METHODS

Experimental animals

The study was performed on 111 adult, clinically healthy Iranian fat-tailed sheep from both sexes (28 male and 83 female; 3-4 years old) reared mainly in South Iran (Fars province). Animals were fed with hay (mainly alfalfa and grass). All animals were treated against internal and external parasites. They were treated with Fenbendazole (Damloran Company, Borujerd, Iran) 10 mg/kg, 30 days prior to the study. A healthy condition was established by clinical and laboratory examination (haematology, clinical biochemistry and other relevant tests).

Sample analysis

Blood samples for determination of oxidative stress parameters were obtained between 8 and 9 am, to avoid diurnal influences, by jugular venepuncture into vacutainers containing EDTA (in a 10:1 ratio) as an anticoagulant. After centrifugation at 750 g for 15 min, plasma was separated and erythrocyte lysate was prepared (Ivanov, 1999). After triple washing of erythrocyte mass with physiological solution, 0.5 mL of cell suspension was dissolved in 2 mL cold water for lysis of erythrocytes. Haemoglobin was then precipitated by adding 1.8 mL water and 0.2 mL ethanol/chloroform (3:5/v:v) to 0.2 mL lysate. The tubes were shaken for 5 min and centrifuged at 750 g for 20 min. The supernatant was used for the determination of enzyme activities.

SOD activity was measured by a modified method of iodophenyl nitrophenol phenyltetrazolium chloride (RANSOD Kit, Randox, UK). This method employs xanthine and xanthine oxidase (XOD) to generate superoxide radicals which react with 2-(4-iodophenyl)-3-(4-nitrophenol)-5-phenyl-tetrazolium chloride (INT) to form a red formazan dye. SOD activity was then measured by the degree of inhibition of this reaction. One unit of SOD was that which caused a 50% inhibition of the reduction rate of INT under the assay condition.

GPX activity was measured by the method of Paglia and Valentine (1967), using RANSEL Kit, (Randox, UK). GPX catalyses the oxidation of glutathione (GSH) by cumene hydroperoxide. In the presence of GR and NADPH, the oxidized glutathione is immediately converted to the reduced form with a concomitant oxidation of NADPH to NADP⁺. The decrease in absorbance was measured at 340 nm. The thiobarbituric acid method was used to quantitate MDA-reactive products (Plaser and Cushman, 1966). Thiobarbituric acid (TBA) and MDA react to form a Schiff base adduct under high temperature/acidic conditions to produce a chromogenic/fluorescent product that can be easily measured employing various analytical techniques such as spectrophotometric or fluorometric methods. CAT activity was estimated in erythrocyte lysate by the method of

Beers and Sizer (1952), using the ferrous oxidation in xylenol orange (FOX) assay. Samples containing CAT were incubated with H₂O₂ for varying time intervals prior to rapid mixing of aliquots of the incubation mixtures with FOX reagent, which measures residual H₂O₂. Absorbance was then read at 560 nm after 30-min incubation at room temperature. Decay of H₂O₂ is proportional to CAT activity in the original sample. The haemoglobin concentration of the lysate was determined by the cyanmethaemoglobin method (Mahoney *et al.*, 1993).

Data analysis

Results are presented as mean, standard error of mean (SEM), median, mode, standard deviation (SD), range and minimal and maximal values. Raw data were tested for normal distribution using SPSS software and the Kolmogorov-Smirnov method. All reference ranges had a normal distribution by confidence interval 90%. The reference range was determined according to mean \pm 2.397 \times SD. Statistical analysis was performed using t-test for comparison of the differences between animals of two sexes.

RESULTS

Reference values of superoxide dismutase (SOD), glutathione peroxidase (GPX), catalase (CAT) and malondialdehyde (MDA) in 111 adult male and female clinically healthy Iranian fat-tailed sheep are shown in Table 1. The values were as followed: SOD 948.65-1011.50 U/gHb, GPX 191.67-196.52 U/gHb, CAT 1834.29-1915.63 U/gHb and MDA 0.53-0.60 μ mol/l. There were no significant differences in oxidative stress parameters (SOD, GPX, CAT and MDA) between both sexes ($P > 0.05$).

DISCUSSION

There is little information about the oxidative stress parameters (SOD, GPX, CAT and MDA) of sheep. The activity of antioxidant enzymes in adult clinically healthy Iranian fat-tailed sheep were comparable to the values reported by Lightbody *et al.* (2001) and Kizil *et al.* (2007). A comparison of our results to those reported by Todorova *et al.* (2005) showed that reference values of oxidative stress indices such as MDA, CAT and SOD in carnivores (dogs and cats) were higher than the values obtained in Iranian fat-tailed sheep. This outcome was consistent in both sexes. This status may be due to their different diets, in other word, generation of free radicals in carnivores is more than that in herbivores. The oxidative status is variable and can be changed by different factors. In the present study, all Iranian fat-tailed sheep were adult. In normal conditions, age influences greatly free radical generation and consequently, the level of antioxidant defense enzymes. In similar investigations on rats at different ages, decreased plasma levels of antioxidant vitamins C and E, decreased SOD activity, and increased CAT activity was observed (De and Durad, 1991).

Table 1: Reference values of superoxide dismutase (SOD), glutathione peroxidase (GPX), catalase (CAT) and malondialdehyde (MDA) in adult male and female clinically healthy Iranian fat-tailed sheep

| Sex | Parameter | No | Mean | SEM | SD | Reference range | Median | Mode | Minimum | Maximum |
|--------|-------------------|-----|--------|-------|-------|-----------------|---------|--------|---------|---------|
| Male | SOD(U/gHb) | 28 | 987.3 | 15.9 | 84.26 | 954.68-1020.03 | 968 | 920.00 | 860 | 1160 |
| | GPX(U/gHb) | 28 | 193.5 | 1.34 | 7.13 | 190.77-196.30 | 194.00 | 186.00 | 180 | 205 |
| | CAT(U/gHb) | 28 | 1880.0 | 24.5 | 129.8 | 1829.68-1930.38 | 1910.50 | 1826.0 | 1587 | 2067 |
| | MDA(μ mol/l) | 28 | 0.56 | 0.01 | 0.08 | 0.53-0.60 | 0.54 | 0.53 | 0.46 | 0.77 |
| Female | SOD(U/gHb) | 83 | 973.24 | 8.33 | 75.95 | 956.65-989.82 | 961.00 | 954.00 | 861 | 1136 |
| | GPX(U/gHb) | 83 | 193.68 | 0.81 | 7.41 | 192.06-195.30 | 194.00 | 198.00 | 180 | 205 |
| | CAT(U/gHb) | 83 | 1867.5 | 16.33 | 148.8 | 1835.05-1900.05 | 1913.00 | 1836.0 | 1518 | 2062 |
| | MDA(μ mol/l) | 83 | 0.57 | 0.01 | 0.09 | 0.55-0.59 | 0.57 | 0.48 | 0.41 | 0.76 |
| Total | SOD(U/gHb) | 111 | 984.58 | 7.48 | 80.91 | 948.65-1011.50 | 964.50 | 947.00 | 861 | 1145 |
| | GPX(U/gHb) | 111 | 193.54 | 1.00 | 7.54 | 191.67-196.52 | 194.00 | 194.00 | 180 | 205 |
| | CAT(U/gHb) | 111 | 1874.9 | 12.33 | 141.7 | 1834.29-1915.63 | 1912.00 | 1833.0 | 1561 | 2064 |
| | MDA(μ mol/l) | 111 | 0.57 | 0.008 | 0.009 | 0.53-0.60 | 0.5600 | 0.4900 | 0.44 | 0.76 |

In the present study, there were no significant differences in oxidative stress parameters between both sexes ($P>0.05$). There is a good evidence to show that sex differences in oxidative status exist in different species. In many species, females live longer than males and it is probably associated with free radicals which are in lower amount in the mitochondria of females than males (Sastre *et al.*, 2002). The longer lifespan in females may be due to the higher gene expression of antioxidants and the lower oxidative damage of mitochondria (Borras *et al.*, 2003). Moreover, there is evidence for the strong antioxidant properties of estrogen (Tudus, 2000) but not for progesterone and testosterone (Barp *et al.*, 2002). In animal species, further investigations are needed to interpret these changes.

In conclusion, the results of the present investigation provide a baseline data for comparisons between normal and abnormal states; therefore, can provide a better understanding in diseases accompanied by oxidative stress.

REFERENCES

- Barp J, AS Araujo, TR Fernands, KV Rigatto, S Llesuy, A Belloklein and P Singal, 2002. Myocardial antioxidant and oxidative stress changes due to sex hormones. *Brazilian J Med Biol Res*, 35: 1075-1081.
- Beers R and T Sizer, 1952. Spectrophotometric method for measuring the breakdown of hydrogen peroxide by catalase. *J Biol Chem*, 195: 133-138.
- Borras C, J Sastre, D Garcia-Sala, A Lloret, FV Pallardo and J Vina, 2003. Mitochondria from females exhibit higher antioxidant gene expression and lower oxidative damage than males. *Free Radical Biol Med*, 34: 546-552.
- De AK and R Durad, 1991. Age-associated changes in antioxidants and antioxidative enzymes in rat. *Mech Ageing Dev*, 59: 123-128.
- Deger Y, A Ertekin, S Deger and H Mert, 2008. Lipid peroxidation and antioxidant potential of sheep liver infected naturally with distomatosis. *Turkiye, Parazitoloji Dergisi*, 32: 23-26.
- Gonzales R, C Auclair, E Voisin, H Gautero, D Dherym and P Boivin, 1984. Superoxide dismutase, catalase and glutathione peroxydase in red blood cells from patients with malignant diseases. *Cancer Res*, 44: 4137-4139.
- Hallwell B and JMC Gutteridge, 1999. *Free radicals in Biology and Medicine*. 3rd Ed, Oxford University Press, Oxford, UK.
- Ivanov IT, 1999. Low pH-induced hemolysis of erythrocytes is related to the entry of the acid into cytosole and oxidative stress on cellular membranes. *Biochem. Biophys Acta*, 1415: 349-360.
- Kargin F and UR Fidanci, 2001. Kidney diseases and antioxidative metabolism in dogs. *Turkish J Vet Anim Sci*, 25: 607-613.
- Khadija, AA Ati, S Mohammed, AM Saad and HE Mohamed, 2009. Response of broiler chicks to dietary monosodium glutamate. *Pakistan Vet J*, 29(4): 165-168.
- Kidd PM, 1997. Glutathione: Systemic protectant against oxidative and free radical damage. *Alter Med Rev*, 2: 155-176.
- Kizil O, H Ozdemir, M Karahan and M Kizil, 2007. Oxidative stress and antioxidant status in goats naturally infected with *Mycoplasma agalactiae*. *Rev Med Vet*, 158: 326-330.
- Kotos HA and EP Wel, 1986. Superoxide production in experimental brain injury. *J Neurosurgery*, 64: 803-807.
- Lightbody JH, LM Stevenson, F Jackson, K Donaldson and DG Jones, 2001. Comparative aspects of plasma antioxidant status in sheep and goats, and the influence of experimental abomasal nematode infection. *J Comp Pathol*, 124:192-199.
- Lopaczyski W and SH Zeisel, 2001. Antioxidants, programmed cell death and cancer. *Nutr Res*, 21: 295-307.
- Mahoney JJ, HJ Vreman, DK Stevenson and AL Van Kessel, 1993. Measurement of carboxyhemoglobin and total hemoglobin by five specialized spectrophotometers (Co-oximeters) in comparison with reference methods. *Clin Chem*, 39: 1693-1700.
- Paglia DE and WN Valentine, 1967. Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. *J Lab Clin Med*, 70: 158-169.
- Patterson RA and DS Leacke, 1998. Human serum, cysteine and histidine inhibit the oxidation of low

- density lipoprotein less at acidic pH. Feder European Biochem Soc Lett, 434: 317-321.
- Plaser ZA and LL Cushman, 1966. Estimation of product of lipid peroxidation (malonyl dialdehyde) in biochemical systems. Analy Biochem, 16: 359-364.
- Sastre J, C Borrás, D Garcia-Sala, A Lloret, FV Pallardo and J Vina, 2002. Mitochondrial damage in aging and apoptosis. Annl New York Acad Sci, 959: 448-451.
- Simsek S, A Yuce and AE Utuk, 2006. Determination of serum malondialdehyde levels in sheep naturally infected with *Dicrocoelium dendriticum*. Firat Univ Saglik Bil Dergisi, 20: 217-220.
- Singh SK, TD Dua, A Tondon, S Kumari, G Ray and S Batra, 1999. Status of lipid peroxidation and antioxidant enzymes in hypoxic ischemic encephalopathy. Indian Ped, 26: 659-668.
- Todorova I, G Simeonova, D Kyuchukova Dinev and V Gadjeva, 2005. Reference values of oxidative stress parameters (MDA, SOD, CAT) in dogs and cats. Comp Clin Pathol, 13:190-194.
- Tudus PM, 2000. Estrogen and gender effects on muscle damage, inflammation, and oxidative stress. Canadian J Appl Physiol, 25: 274-287.