



## Chemical Composition of Different Varieties of Linseed

M. Laiq Khan\*, M. Sharif, M. Sarwar, Sameea<sup>1</sup> and M. Ameen

Institute of Animal Nutrition and Feed Technology, University of Agriculture, Faisalabad, <sup>1</sup>Government Islamia College, Faisalabad, Pakistan

\*Corresponding author: [drlaiqkhan@gmail.com](mailto:drlaiqkhan@gmail.com)

### ARTICLE HISTORY

Received: June 30, 2009  
Revised: November 04, 2009  
Accepted: November 21, 2009

#### Key words:

Linamarin  
Linseed  
Minerals  
Proximate composition  
Varieties

### ABSTRACT

The present study was conducted to investigate chemical composition of six varieties of linseed (Chandni, LS-29, LS-49, LS-70, LS-75 and LS-76). Proximate composition, mineral profile and cyanogenic glycosides (linamarin) were determined. Average proximate composition values for linseed i.e. crude protein, ether extract, crude fiber, ash and nitrogen free extract were 24.18, 37.77, 4.78, 3.50 and 25.86%, respectively. Higher values of crude protein, ether extract, crude fiber and nitrogen free extract were observed in varieties LS-49, LS-70, LS-29 and Chandni, respectively. Average mineral contents in linseed i.e. Ca, Mg, K, Na, Cl, P, Cu, Fe, Mn and Zn were 0.39, 0.09, 1.41, 0.05, 0.08, 0.89, 4.67, 50.56, 8.29 and 13.55 ppm, respectively. Among micro minerals, varieties LS-29 and LS-70 were higher in Cu contents; LS-75 was higher in Fe content, while LS-49 was higher in Mn and Zn contents. Among macro minerals, level of Ca was higher in LS-70, levels of Mg, K and Na were higher in Chandni, while P was higher in LS-49. Average amount of linamarin in linseed was 31.05mg/100 gm DM. The variety LS-75 had the highest (35.22 mg/100 gm) linamarin content, while variety LS-70 had least (26.22 mg/100 gm) amount of linamarin. In conclusion, there is significant difference in chemical composition among linseed varieties. The varieties LS-49 showed higher crude protein content, LS-70 showed greater oil content, while LS-75 had higher content of linamarin.

©2010 PVJ. All rights reserved

**To cite this article:** Khan ML, M Sharif, M Sarwar, Sameea and M Ameen, 2010. Chemical composition of different varieties of linseed. *Pak Vet J*, 30(2): 79-82.

### INTRODUCTION

Linseed (*Linum usitatissimum L*), also called flaxseed, has got significance due to its nutrients and pharmaceutical uses. It is used for edible and lightening purposes in some Asian countries and can also be used as a substitute part of animal fat in poultry diets.

The linseed contains about 36-48% oil content which is high in unsaturated fatty acids, especially linolenic acid (Enser *et al.*, 2000; Kouba *et al.*, 2003; Kouba, 2006). Inclusion of linseed in the diet results in increased polyunsaturated fatty acids (Matthews *et al.*, 2000; Riley *et al.*, 2000), which protect the body against cancer (Rose and Connolly, 1999), reduce the chances of cardiovascular diseases and certain other health related problems (Alexander, 1998). So, linseed may play a vital role in reducing these health risk problems due to its nutritive value which depends upon different factors like cultivar, locality,

sowing date and year of production, with cultivar being the most important factor (Oomah *et al.*, 1992). However, information regarding chemical composition of various varieties of linseed is limited.

In view of the availability of linseed with high protein contents, good amino acids profile and essential polyunsaturated fatty acids, there was a need to explore the nutritive value of different available varieties of linseed. The present study was, therefore, conducted to determine the comparative proximate composition, macro and micro-minerals and linamarin contents of six varieties of linseed grown in Pakistan.

### MATERIALS AND METHODS

Six varieties of linseed were procured from Directorate of Ayub Agriculture Research Institute, Faisalabad, Pakistan. These varieties included Chandni, LS-29, LS-49,

LS-70, LS-75 and LS-76. Proximate analysis, minerals profile and linamarin content/cyanogenic glycosides of these varieties were determined. For this purpose, six aliquot samples of each variety were taken for analysis.

In proximate analysis, dry matter (DM) of linseed was determined by drying it in a forced air oven at 105°C till constant weight was attained (AOAC, 1990). Linseeds were milled in a hammer mill to produce finely ground powder. Percent nitrogen was estimated by Kjeldhal method (AOAC, 1990) and crude protein (CP) was calculated by multiplying nitrogen (%) with factor 6.25 (AOAC, 1990). Crude fat determination was carried out by petroleum ether (BP 40-60°C) extraction, using the soxhlet procedure. Crude fiber and ash were determined according to the procedure described by AOAC (1990), while Na, K, Ca, Cl, S and Mg were analyzed using atomic absorption spectrophotometry (model 4, Perkin-Elmer, Norwalk) and P was determined photometrically via Spectronic 1001 (Milton Roy Co., Cincinnati, OH). Among micromineral elements, Fe, Mn, Zn and Cu were analyzed using atomic absorption spectrophotometry.

Linamarin content/cyanogenic glycosides were determined using distillation and spectrophotometric method (Krishna and Ranjhan, 1981). For this purpose, 5g of finely ground sample was taken into 250 ml round bottom flask and 5 ml chloroform was added in it. It was then steam distilled and distillate was collected upto 60 ml in 100 ml flask containing 5 ml of 2% KOH solution. The cyanide content with the reaction of KOH was converted into KCN and was collected in distillate. It was then estimated by spectrophotometry.

The data were subjected to statistical analysis by using analysis of variance technique with completely randomized design. The significance of difference among means was compared using Duncan's multiple range test (Steel and Torrie, 1984).

## RESULTS AND DISCUSSION

### Proximate composition

The mean values for crude protein content varied from 22.37 to 27.24% ( $P < 0.05$ ). The data indicated that the variety LS-49 had the highest protein content (27.24%), followed by the variety LS-75 (25.08%) and LS-76 (24.02%), whereas the variety LS-70 had the lowest (22.37%) protein content. The variety LS-29 was not significantly different from the variety LS-70 but it differed significantly from rest of the varieties for this trait (Table 1). Contrary to our findings, Gambus *et al.* (2003) reported that linseed varieties did not differ in protein contents.

There was wide variation in ether extract value, which ranged from 35.03 to 41.23% among different varieties under study ( $P < 0.05$ ) as shown in Table 1. The highest ether extract was observed in the variety LS-70 (41.23%), followed by LS-75 (38.43%) and LS-76 (38.40%), while the variety LS-49 had the lowest ether extract content (35.03%). Results of Gambus *et al.* (2003) supported our findings. The results of present study are also in agreement with the findings of other researchers (Kouba *et al.*, 2003; Kouba,

2006). In our findings, protein and oil contents varied negatively with each other (Table 1). These are affected by different factors like date of sowing, seed rate, level of nitrogen used (Taylor and Morrice, 1991) and genetic variability (Karmuka *et al.*, 1988). Green and Marshal (1981) reported a positive correlation between oil content and seed weight.

Average crude fiber contents varied from 4.16 to 5.14% ( $P < 0.05$ ), as shown in Table 1. The variety LS-29 had the greatest crude fiber content (5.14%), followed by LS-76 (5.01%) and Chandni (5.00%), whereas the variety LS-49 had the lowest crude fiber content (4.16%). Gambus *et al.* (2003) also observed variation in crude fiber in linseed varieties. This may be due to genetic variability and other environmental factors.

Nitrogen free extract contents varied from 23.57 to 29.05% ( $P < 0.05$ ). It was observed that variety Chandni had the highest nitrogen free extract content (29.05%), whereas the variety LS-75 had the lowest nitrogen free extract content (23.57%). The mean value for the ash content varied from 3.18 to 4.35% (Table 1). The data indicated that variety LS-29 had higher ash content (4.35%) than all other varieties of linseed. It has been observed that mineral contents varied with different levels of inorganic matter applied, interaction among minerals and soil types (Olomu and Racz, 1974).

The data indicates that the varieties LS-49 and LS-70 are more nutritious than all other varieties of linseed.

### Mineral profile

Average Cu content varied from 2.30 to 5.77 ppm ( $P < 0.05$ ) as shown in Table 2. The data indicated that the varieties LS-29, LS-70 and Chandni had the highest Cu contents (5.75 to 5.77 ppm), whereas the variety LS-75 had the lowest Cu content (2.31 ppm). The mean values for Fe content varied ( $P < 0.05$ ) from 42.25 to 71.16 ppm (Table 2). The variety LS-75 had the highest Fe content (71.16 ppm), followed by the variety LS-49 (52.21 ppm) and LS-70 (48.72 ppm). The variety LS-29 had the lowest Fe content (42.25 ppm). The concentrations of Fe in seed differ due to the interaction of different minerals like cadmium, zinc (Moraghan, 1993) and chlorosis (Olomu and Racz, 1974). The mean value of Mn content varied from 6.85 to 9.28 ppm (Table 2). The variety LS-49 had the greatest Mn content (9.28 ppm), followed by the variety LS-70 (8.75 ppm) and LS-76 (8.71 ppm), whereas the variety LS-29 had the lowest Mn content (6.85 ppm). This difference depends upon the type of soil (Olomu and Racz, 1974). The mean values for Zn content varied from 11.72 to 16.52 ppm (Table 2). The variety LS-49 had the greatest Zn content (16.52 ppm), whereas LS-75 had the lowest Zn content (11.72 ppm). The Zn content varied due to soil temperature and level of Fe source (Moraghan, 1978). The Zn fertilizer is the major factor influencing seed Zn concentration (Moraghan, 1993).

The mean value of Ca content varied from 0.31 to 0.46% ( $P < 0.05$ ), as shown in Table 3. Our findings are consistent with NRC (1988) which indicated that linseed contains about 0.37% Ca content. The variety LS-70 had the highest ( $P < 0.05$ ) Ca content (0.46%), followed by the

variety Chandni (0.41%) and LS-75 (0.40%), whereas the variety LS-49 had the lowest Ca content (0.31%). This indicates variability in Ca contents among different linseed varieties. The mean value for Mg content varied from 0.07 to 0.10% (Table 3). The data indicated that the variety Chandni had the highest Mg content (0.10%), followed by the variety LS-29 (0.09%), LS-75 (0.09%) and LS-76 (0.09%), whereas the variety LS-49 had the lowest Mg content (0.07%). Singh *et al.* (1990) reported that Mg contents decreased with increased K level and applied Mg contents.

The mean values for K content varied from 1.14 to 1.92% (Table 3). The results showed that the variety Chandni had the highest ( $P < 0.05$ ) K content (1.92%), followed by the variety LS-49 (1.42%), whereas the variety LS-29 had the lowest K content (1.14%). The K contents increased with increased level of K fertilizer (Singh *et al.*, 1990). The mean values for Na content varied from 0.04 to 0.06% (Table 3). Variety Chandni had the greatest Na content (0.06%), whereas the variety LS-75 had the lowest Na content (0.04%). The average values for Cl content varied from 0.06 to 0.08% (Table 3). The variety Chandni, LS-29 and LS-70 had the highest ( $P < 0.05$ ) Cl content, whereas the variety LS-75 had the lowest Cl content (0.06%). The mean value for P content varied from 0.80 to 0.98%. Variety LS-49 had the greatest P content (0.98%), followed by the variety LS-70 (0.92%), whereas the variety Chandni had the lowest P content (0.80%). There is

variability in P content among various linseed varieties. Its concentration depends upon its uptake and level of fertilizer applied (Chaubey and Dwivedi, 1995). The variation in mineral contents is due to inorganic matter applied and different soil types (Olomu and Racz, 1974).

#### Cyanogenic glycoside

The mean values for linamarin content varied from 26.22 to 35.22 mg/100 gm DM (Table 3). Variety L.S-70 had the least linamarin content (26.22 mg/100 gm), followed by the variety LS-29 (28.22 mg/100 gm), whereas the variety LS-75 had the highest linamarin content (35.22 mg/100 gm). This shows that there is variability in linamarin contents among various linseed varieties. Oomah *et al.* (1992) supported our findings and stated that linamarin contents varied among various varieties of linseed. Majak *et al.* (1981) also reported that cyanogenic glycoside is found in trace amounts in various plants. Presence of cyanogenic glycoside in higher quantity limits its use in animal ration (Oomah *et al.*, 1992).

This study revealed that there are significant differences in chemical composition among various varieties of linseed. The variety LS-49 was higher in crude protein content; LS-70 rich in oil content, whereas LS-75 showed higher linamarin content. The varieties L.S-49 and Chandni are good sources of micro and macro minerals. The chemical composition of linseed will be helpful in selecting various varieties for human consumption.

**Table 1: Proximate composition of different linseed varieties (%)**

Parameter	Varieties						Average value	SEM
	Chandni	L.S-29	L.S-49	L.S-70	L.S-75	L.S-76		
Crude protein	23.57 <sup>c</sup>	22.82 <sup>d</sup>	27.24 <sup>a</sup>	22.37 <sup>d</sup>	25.08 <sup>b</sup>	24.02 <sup>b</sup>	24.18	0.40
Ether extract	36.58 <sup>bc</sup>	36.93 <sup>bc</sup>	35.03 <sup>c</sup>	41.23 <sup>a</sup>	38.43 <sup>b</sup>	38.40 <sup>b</sup>	37.77	0.53
Crude fiber	5.00 <sup>ab</sup>	5.14 <sup>a</sup>	4.16 <sup>d</sup>	4.92 <sup>b</sup>	4.46 <sup>c</sup>	5.01 <sup>ab</sup>	4.78	0.09
Ash	3.33 <sup>b</sup>	4.35 <sup>a</sup>	3.25 <sup>b</sup>	3.43 <sup>b</sup>	3.18 <sup>b</sup>	3.43 <sup>b</sup>	3.50	0.10
Nitrogen free extract	29.05 <sup>a</sup>	25.28 <sup>bc</sup>	26.82 <sup>b</sup>	25.63 <sup>bc</sup>	23.57 <sup>c</sup>	24.78 <sup>bc</sup>	25.86	0.49

Means in a row with different superscripts differ significantly ( $P < 0.05$ ).

**Table 2: Average micro mineral profile of different varieties of linseed**

Minerals (ppm)	Varieties						Average value	SEM
	Chandni	L.S-29	L.S-49	L.S-70	L.S-75	L.S-76		
Cu	5.75 <sup>a</sup>	5.77 <sup>a</sup>	5.38 <sup>b</sup>	5.76 <sup>a</sup>	2.31 <sup>d</sup>	3.07 <sup>c</sup>	4.67	0.03
Fe	44.52 <sup>d</sup>	42.25 <sup>e</sup>	52.21 <sup>b</sup>	48.72 <sup>c</sup>	71.16 <sup>a</sup>	44.52 <sup>d</sup>	50.56	0.24
Mn	8.27 <sup>c</sup>	6.85 <sup>e</sup>	9.28 <sup>a</sup>	8.74 <sup>b</sup>	7.86 <sup>d</sup>	8.71 <sup>b</sup>	8.29	0.02
Zn	13.24 <sup>d</sup>	12.35 <sup>e</sup>	16.52 <sup>a</sup>	14.12 <sup>b</sup>	11.72 <sup>f</sup>	13.32 <sup>c</sup>	13.55	0.04

Means in a row with different superscripts differ significantly ( $P < 0.05$ ).

**Table 3: Average macro mineral profile and linamarin contents of different varieties of linseed (%)**

Minerals	Varieties						Average value	SEM
	Chandni	L.S-29	L.S-49	L.S-70	L.S-75	L.S-76		
Ca	0.41 <sup>b</sup>	0.38 <sup>d</sup>	0.31 <sup>f</sup>	0.46 <sup>a</sup>	0.40 <sup>c</sup>	0.37 <sup>e</sup>	0.39	0.01
Mg	0.10 <sup>a</sup>	0.09 <sup>b</sup>	0.07 <sup>d</sup>	0.08 <sup>c</sup>	0.09 <sup>b</sup>	0.09 <sup>b</sup>	0.09	0.01
K	1.92 <sup>a</sup>	1.14 <sup>f</sup>	1.42 <sup>b</sup>	1.39 <sup>c</sup>	1.37 <sup>d</sup>	1.24 <sup>e</sup>	1.41	0.06
Na	0.06 <sup>a</sup>	0.05 <sup>b</sup>	0.05 <sup>b</sup>	0.05 <sup>b</sup>	0.04 <sup>c</sup>	0.05 <sup>b</sup>	0.05	0.01
Cl	0.08 <sup>a</sup>	0.08 <sup>a</sup>	0.07 <sup>b</sup>	0.08 <sup>a</sup>	0.06 <sup>c</sup>	0.07 <sup>b</sup>	0.08	0.01
P	0.58 <sup>f</sup>	0.89 <sup>c</sup>	0.98 <sup>a</sup>	0.92 <sup>b</sup>	0.78 <sup>e</sup>	0.85 <sup>d</sup>	0.89	0.01
Linamarin (mg/100gm)	34.23 <sup>b</sup>	28.22 <sup>e</sup>	29.22 <sup>d</sup>	26.22 <sup>f</sup>	35.22 <sup>a</sup>	33.22 <sup>c</sup>	31.05	0.81

Means in a row with different superscripts differ significantly ( $P < 0.05$ ).

## REFERENCES

- Alexander JW, 1998. Immunonutrition: the role of 0-3 fatty acids. *Nutrition*, 14: 627-633.
- AOAC, 1990. *Official Methods of Analysis*. 15<sup>th</sup> Ed, Association of Analytical Chemists. Arlington, Virginia, USA.
- Chaubey AK and KN Dwivedi, 1995. Effect of N, P and S and their interaction on yield of, and nutrient uptake by, linseed (*Linum usitatissimum L.*). *J Indian Soc Soil Sci*, 43: 72-75.
- Enser M, RI Richardson, JD Wood, BP Gill and PR Sheard, 2000. Feeding linseed to increase the n-3 PUFA of pork: Fatty acid composition of muscle, adipose tissue, liver and sausages. *Meat Sci*, 55: 201-212.
- Gambus H, F Borowiec and T Zajac, 2003. Chemical composition of linseed with different colour of bran layer. *Polish J Food Nutr Sci*, 12: 67-70.
- Green AG and DR Marshal, 1981. Variation for oil quantity and quality in linseed (*Linum usitatissimum L.*). *Austr J Agri. Res*, 32: 599-607.
- Karmuka KS, T Dawa and GS Sethi, 1988. Combining ability analysis for oil content and iodine value in linseed (*Linum usitatissimum L.*). *Indian J Agri Sci*, 58: 252-254.
- Krishna G and SK Ranjhan, 1981. *Laboratory Manual for Nutrient Research*. 1<sup>st</sup> Ed, Vikas Publishing House Pvt Ltd New Delhi, India.
- Kouba M, M Enser, FM Whittington, GR Nute and JD Wood, 2003. Effect of a high-linolenic acid diet on lipogenic enzyme activities, fatty acid composition, and meat quality in the growing pig. *J Anim Sci*, 81: 1967-1979.
- Kouba M, 2006. Effect of dietary omega-3 fatty acids on meat quality of pigs and poultry. In: MC Teale (ed). *Omega 3 Fatty Acid Research*. Nova Publishers. New York, USA, pp: 225-239.
- Majak W, RE McDiarmid and JW Hall, 1981. The cyanide potential of Saskatoon serviceberry and chokecherry. *Canadian J Anim Sci*, 61: 681-686.
- Matthews KR, DB Homer, F Thies and PC Calder, 2000. Effect of whole linseed (*Linum usitatissimum*) in the diet of finishing pigs on growth performance and on the quality and fatty acid composition of various tissues. *British J Nutr*, 83: 637-643.
- Moraghan JT, 1978. Chlorotic dieback in flax. *Agron J*, 70: 501-505.
- Moraghan JT, 1993. Accumulation of cadmium and selected elements in flax seed grown on a calcareous soil. *Plant Soil*, 150: 61-68.
- NRC, 1988. *Nutrient Requirements of Dairy Cattle*. 6<sup>th</sup> Ed, National Research Council, Natl Acad Sci, Washington, DC, USA.
- Olomu MO and GJ Racz, 1974. Effect of soil water and aeration on Fe and Mn utilization by flax. *Agron J*, 66: 523-526.
- Oomah BD, G Mazza and EO Kenaschuk, 1992. Cyanogenic compounds in flaxseed. *J Agri Food Chem*, 40: 1346-1348.
- Riley PA, M Enser, GR Nute and JD Wood, 2000. Effects of dietary linseed on nutritional value and other quality aspects of pig muscle and adipose tissue. *Anim Sci*, 71: 483-500.
- Rose DP and JM Connolly, 1999. Omega-3 fatty acids as cancer chemopreventive agents. *Pharmacol Therap*, 83: 217-244.
- Singh RP, V Singh and V Singh, 1990. Effects of S, Mg and K on yield and uptake by linseed. *J Indian Soc Soil Sci*, 38: 169-170.
- Steel RGD and JH Torrie, 1984. *Principles and Procedures of Statistics*. 2<sup>nd</sup> Ed, McGraw-Hill Book Co, Inc, New York, NY, USA.
- Taylor BR and LAF Morrice, 1991. Effects of husbandry practices on the seed yield and oil content of linseed in Northern Scotland. *J Sci Food Agri*, 57: 189-198.