



SHORT COMMUNICATION

Serological Signs of West Nile Virus Infection in Horse Serum Samples Collected for Equine Infectious Anemia Virus screening in the Northeastern Turkey: Traces of Past

Zafer Yazici^{1*}, Cuneyt Tamer¹, Semra Gumusova¹, Emre Ozan², Sabri Hacıoğlu³, Aykut Ozkul³ and Harun Albayrak¹

¹Department of Virology, Faculty of Veterinary Medicine, Ondokuz Mayıs University, 55139, Samsun, Turkey

²Virology Laboratory, Veterinary Control Institute, Samsun, Turkey; ³Department of Virology, Faculty of Veterinary Medicine, Ankara University, 06110 Ankara, Turkey

*Corresponding author: zyazici@omu.edu.tr

ARTICLE HISTORY (18-215)

Received: June 24, 2018
Revised: July 06, 2018
Accepted: July 07, 2018
Published online: September 04, 2018

Key words:

Horse
Serology
West Nile virus

ABSTRACT

West Nile virus (WNV) is an important arthropod-borne pathogen and a leading cause of morbidity and mortality in human and animal populations in many parts of the world. The presence of WNV in Turkey has been reported in various studies during the last 15 years. In this study, 1500 horse serum samples collected from Ardahan Province in Northeastern Turkey were screened by using a virus neutralization test against WNV and then positives were confirmed by a plaque reduction test-₉₀ (PRNT₉₀). The overall seropositivity was 0.13% (2/1500), with neutralizing antibody titers determined to be 1/40 and 1/80. These findings suggest that some of horses in Ardahan had been exposed to WNV and also indicates the evidence of WNV circulation in the Northeastern Turkey.

©2018 PVJ. All rights reserved

To Cite This Article: Yazici Z, Tamer C, Gumusova S, Ozan E, Hacıoğlu S, Ozkul A and Albayrak H, 2019. Serological signs of West Nile virus infection in horse serum samples collected for Equine Anemia Infectious Virus screening in the Northeastern Turkey: traces of past. *Pak Vet J*, 39(1): 121-123. <http://dx.doi.org/10.29261/pakvetj/2018.083>

INTRODUCTION

West Nile virus (WNV) is an important arthropod-borne pathogen transmitted by mosquitoes and can result in serious illness in both human and animal populations. The virus is a member of the Japanese encephalitis virus (JEV) antigenic serocomplex and is a member of the genus *flavivirus* within the family *Flaviviridae* (McVey *et al.*, 2015). WNV has become endemic in many countries of Europe, America, Africa, the Middle East and Asia following the first isolation of virus in Uganda in 1937 (David and Abraham, 2016).

Mosquitoes belonging to the genus *Culex* within the family *Culicoidae* are the primary vectors of WNV. The natural cycle of WNV is mosquito-bird-mosquito (Lafri *et al.*, 2017). Birds are considered to be the primary reservoir host of WNV, with high levels of viremia all owing transmission to *Culex* mosquitoes. Both humans and horses represent dead-endhosts, which develop low-level viremia that precluding onward infection of feeding mosquitoes (Lafri *et al.*, 2017).

Horses are more susceptible to WNV as compared to the other animal species. Fatal meningo encephalitis can occur in horses infected with WNV (Lafri *et al.*, 2017). Historically, many WNV outbreaks in horses have previously been reported in numerous parts of the world

e.g., Morocco, Israel, Northern Italy, France, USA, Canada and Tunisia (Di Sabatino *et al.*, 2014), with several countries, such as Algeria, South France and Corsica, reporting outbreaks more recently (Di Sabatino *et al.*, 2014). In Turkey, serological and virological evidence of WNV infection in humans and domestic animals has previously been reported (Albayrak and Ozan, 2013; Toplu *et al.*, 2015). In light of the historical presence of WNV, the virus continues to represent a public health importance due to its epizootic and zoonotic potential (David and Abraham, 2016).

The aim of this study was to determine the presence of neutralizing antibodies to WNV in horse sera collected from Northeastern Turkey due to the proximity to countries where human cases have been reported e.g., Armenia, Georgia and Russia.

MATERIALS AND METHODS

This study was conducted in Ardahan province, located in the northeastern part of Turkey, using archived horse serum samples collected in 2009 as part of equine infectious anemia virus (EIAV) surveillance program with the approval of Turkish Ministry of Food, Agriculture and Livestock (Approved date/ no:27.12.2017/3336566). A total of 1500 horse serum samples were randomly selected

among archived sera. All horses were non-vaccinated against WNV and were EIAV seronegative. Serum samples were kept at -20°C until used and were inactivated at 56°C for 30 min prior to performing serological tests.

The Vero-E6 cell line was used in this study. Cells were grown in Dulbecco's modified Eagle's medium (DMEM) (Gibco, UK) supplemented with 10% fetal bovine serum (FBS) (Sigma, UK) and 1% antibiotic (Sigma, UK). The NY99 strain of WNV was kindly provided by Aykut OZKUL from Faculty of Veterinary Medicine Ankara University. Virus was propagated on Vero-E6 cells with DMEM supplemented with 3% FBS.

The WNV titre was established in Vero-E6 using the conventional TCID₅₀ method and were calculated as log₁₀ TCID₅₀/ml. The presence of neutralizing antibodies against the WNV-NY99 strain was investigated by using a conventional virus neutralization test (VNT). Positive serum samples were confirmed with plaque reduction neutralization test (PRNT) using the NY99 strain of WNV as described before (Ozkul *et al.*, 2006). A four-fold greater reciprocal 90% neutralization titer is required for a sample to be considered positive for WNV infection due to the fact that high levels of cross-reaction are observed among *flavi viruses*.

RESULTS AND DISCUSSION

Out of the 1500 serum samples tested, 2 (0.13%) were found to be seropositive for WNV as assessed using VNT. These serum samples were also confirmed using PRNT₉₀ as being WNV seropositive. The positive serum samples had 1/40 and 1/80 mean neutralizing antibody titers.

WNV is an important arthropod-borne pathogen that is considered to be a public health concern. Like humans, horses can become infected with WNV and neurological disorders including fetal meningo encephalitis in severe infections can develop in 8% of exposed horses (Lafri *et al.*, 2017). However, WNV infection is subclinical in the vast majority of horses following exposure to WNV. Transmission is directly related to environmental conditions; so, there are many factors that strongly affect WNV infection, including humidity, wetlands, an abundance of avifauna and mosquitoes etc (Lafri *et al.*, 2017).

In Turkey, Ozkul *et al.* (2006) first serologically reported WNV in horse serum samples collected from the Marmara and Mediterranean Regions with a rate of 13.5%. In subsequent years, seropositivity rates of WNV in horses ranged from 4.1% to 31.6% (Ozkul *et al.*, 2013; Kale *et al.*, 2017). Despite the previous studies indicating WNV circulation in Turkey, there was no previously reported WNV seropositivity in human and horses as well as any notification concerning the WNV identification from mosquitoes, human and horses in Northeast Turkey (Yazici *et al.*, 2012; Albayrak and Ozan, 2013). The only report of WNV in this region of Turkey was reported in goats, with a seroprevalence of 2.85% (Albayrak and Ozan, 2013). Furthermore, there have been no reports of WNV in human, horse or mosquito populations from Ardahan, Georgia, Armenia and Russia, which are considered to be endemic for WNV and border the north-east Turkey and the current study is the first report

concerning the seroprevalence of WNV in horses in the Northeast Turkey. After screening 1500 horse sera we observed a seropositivity rate of 0.13%. This rate could be considered remarkably low, both compared to previous WNV studies in Turkey, and given the abundant breeding habitats for mosquitoes in the Ardahan region. However, this result may have been affected by some factors including the use of archived sera and difference in climate. For example, these serum samples had been collected and stored for an extended period of time. In this context, it cannot be ruled out that the antibody titer in some samples may have dropped depending as a result of any freezing and thawing processes. Furthermore, the climatic condition of a region, such as amount of rainfall, wind status and insecticide applications are particularly important factors in vector biology. In Ardahan Province, yearly averages of temperature, humidity and rain are reported as 3.8°C, 72.6%, 543mm, respectively. When compared, these values are lower than the Western Anatolia region of Turkey where the disease is seen frequently, reportedly 10.3°C, 63%, 2, 950mm respectively. These data are important factors affecting vector abundance. In addition, the existence of *Culicoides* spp was far less in the Northeast Turkey than in the South and West parts of Turkey.

Despite the low seroprevalence, the results reported here indicate that WNV was in circulation in the Northeastern regions of Turkey. Moreover, it could get a warning for WNV infection for human beings living Ardahan and its neighbor areas due to the fact that horses might be one of the sentinel animals for WNV.

Conclusions: The determination of neutralizing antibodies in the horse, human, chicken, and dogs may be thought to have pointed toward a sustained arboviral infection. In this context, the results of this study should be taken into consideration even though the seroprevalence rate is lower than previous reports considering the fact that WNV may have impacts on public health together with its zoonotic potential. It is also recommended that the new serological screening on humans and animals with more recent and /or actual samples should be planned in order to realize dynamic of the virus in the region.

Authors contribution: This study was conducted by ZY, CT and SG, EO helped in sample collection analysis. All authors have approved the final version of the manuscript.

REFERENCES

- Albayrak H and Ozan E, 2013. Seroepidemiological study of West Nile virus and Rift Valley fever virus in some of mammalian species (herbivores) in Northern Turkey. *J Arthropod Borne Dis* 7:90-3.
- David S and Abraham AM, 2016. Epidemiological and clinical aspects on West Nile virus, a globally emerging pathogen. *Infect Dis* 8:571-86.
- Di Sabatino D, Bruno R, Sauro F, *et al.*, 2014. Epidemiology of West Nile disease in Europe and in the Mediterranean Basin from 2009 to 2013. *Biomed Res Int*: 907852.
- Kale M, Gur S, Yapık O, *et al.*, 2017. Serological investigation of West Nile virus infection in domestic horses and donkeys in Turkey. *Pak Vet J* 37:51-4.
- Lafri I, Prat CM, Bitam I, *et al.*, 2017. Seroprevalence of West Nile virus antibodies in equids in the North-East of Algeria and

- detection of virus circulation in 2014. *Comp Immunol Microbiol Infect Dis* 50:8-12.
- Maquart M, Dahmani M, Marie JL, *et al*, 2017. First serological evidence of West Nile virus in horses and dogs from Corsica Island, France. *Vector-borne Zoonotic Dis* 17:275-7.
- McVey DS, Wilson WC, Gay CG, 2015. West Nile Virus. *Rev Sci Tech Off Int Epiz* 34:431-9.
- Ozkul A, Yildirim Y, Pinar D, *et al*, 2006. Serological evidence of West Nile Virus (WNV) in mammalian species in Turkey. *Epidemiol Infect* 134:826-9.
- Ozkul A, Ergunay K, Koysuren A, *et al*, 2013. Concurrent occurrence of human and equine West Nile virus infections in Central Anatolia, Turkey: the first evidence for circulation of lineage I viruses. *Int J Infect Dis* 17:e546-51.
- Toplu N, Oğuzoğlu TÇ, Ural K, *et al*, 2015. West Nile virus infection in horses. *Vet Pathol* 52:1076-6.
- Yazici Z, Albayrak H, Ozan E, *et al*, 2012. The first investigation of West Nile virus in horses using real time rt-PCR in middle black sea region in Turkey. *J Arthropod Borne Dis* 6:151-5.