



RESEARCH ARTICLE

Epidemiological Study of Mastitis in Three Different Strains of Beetal Goat in Selected Districts of Punjab, Pakistan

Muhammad Ijaz Saleem^{*1}, Muhammad Saqib¹, Muhammad Sajjad Khan², Ghulam Muhammad¹ and Sajjad ur Rehman³

¹Department of Clinical Medicine and Surgery; ²Institute of Dairy and Animal Sciences; ³Institute of Microbiology, University of Agriculture, Faisalabad, Pakistan

*Corresponding author: drijazsaleem@gmail.com

ARTICLE HISTORY (18-401)

Received: October 21, 2018
Revised: December 21, 2018
Accepted: December 24, 2018
Published online: December 27, 2018

Key words:

Beetal
Breed
Goat
Mastitis prevalence
Strains

ABSTRACT

Mastitis in goats, analogous to dairy cattle, is among the momentous diseases of economic value in all parts of the world. Local information of epidemiology of goat mastitis is scanty. In the present study, considering geographical distribution of Beetal goat, 172 flocks were randomly pulled out by "Survey tool box" to reduce biased sampling from perie-urban and urban areas of six districts of Punjab. By following the proportionate sampling strategy and assuming the highest population (70%) of Beetal all over the Punjab, Beetal strains were screened for the prevalence of mastitis from Faisalabad, Jhang, Chiniot, Bahawalpur, Muzzafargarh and Rajanpur districts. Screening of the flocks with surf field mastitis test indicated that overall prevalence of caprine mastitis in Beetal breed was 17.39% while in Beetal Faisalabadi strain 16.53% from 116 flocks, in Beetal Makhicheeni 18.06% from 32 flocks and in Beetal Nuqri 19.33% from 24 flocks. Microbiological examination of milk samples showed various types of mastitic microorganisms. The predominant isolates included *Staphylococcus (S). hyicus* (45%), *S. xylosus* (19%), *S. simulans* (16%), *S. aureus* (9%) and unidentifiable *Staphylococcal* species (11%). Antibiotic sensitivity test was conducted on the most prevalent isolates. *S. aureus* and *S. hyicus* showed sensitivity to eight antibiotics: amoxicillin, ampicillin, lincomycin, sulfamethoxazole+trimethoprim, novobiocin, enrofloxacin, amoxicillin+ clavulanic acid and oxytetracycline while *S. xylosus* was sensitive to all antibiotics except novobiocin and *S. simulans* was sensitive to lincomycin, sulfamethoxazole+trimethoprim, enrofloxacin and resistant to all other antibiotics. The study found over all 16.74% mastitis in different strains of Beetal. The most important parameters as; type of housing, farm hygiene, age, stage of lactation, parity, color, body condition score, teat ends, SFMT score and decreased milk yield were found pertinent risk factors so that prompt attention required for mastitis control in goats.

©2018 PVJ. All rights reserved

To Cite This Article: Saleem MI, Saqib M, Khan MS, Muhammad G and Rehman SU, 2019. Epidemiological study of mastitis in three different strains of beetal goat in selected districts of Punjab, Pakistan. *Pak Vet J*, 39(3): 389-394. <http://dx.doi.org/10.29261/pakvetj/2018.118>

INTRODUCTION

The goat is an important animal in many parts of the world, where it is kept as a source of meat, milk and fiber (DaMassa *et al.*, 1992). It is often dubbed as the poor man's cow in subcontinent. Pakistan is the 3rd largest goat rearing country in the world (Khan *et al.*, 2008). In Pakistan, the economic contribution from goat source is 0.719 million (M) tons of milk, 0.6 M tons of mutton and 22 million hides (Economic Survey of Pakistan 2016-17). Goats, particularly of milk breeds, are worthy source of income for small households and are well recognized for providing

nutritional surrogate, particularly for the growing infants in the rural areas of Pakistan.

Goats can bear hard as well as harsh environments in various agro-climatic conditions, predominantly in the tropics and sub-tropics. About 90% goat population is found in the developing countries whereas continent wise, Asia always stand first by producing 80% of total milk mainly from Pakistan, Turkey, China, Bangladesh, Iran and India (Khan *et al.*, 2008).

More than 25 breeds of goat have been reported in Pakistan including the Northern Areas and Azad Jammu and Kashmir. Goats of Beetal, Diara Din Panah (DDP),

Nachi, Damani, Kacchan and Kamori breeds are well recognized for their milk and meat production. The province of Punjab has mainly three autochthonous milk breeds (Beetal, Diara Din Panah and Nachi). There is no current report on milk production potential of various breeds of goat in Pakistan (Iqbal *et al.*, 2008). Among all the breeds of goat in Pakistan, Beetal is the best in terms of milk and meat production. Recently, five different strains of Beetal have been documented including Faisalabadi or Lyalpuri strain found in Faisalabad, Lahore, Sheikhpura, Gujranwala, Jhang, Chiniot, Sargodha, Toba Tek Singh and Khanewal; Makhicheeni strain found in Bahawalpur, Bahawalnagar and Muzaffargarh; Nuqri strain, pure white color, found in Dera Ghazi Khan, Rajanpur and Bahawalpur district; Gujrati strain found in Gujrat, Sargodha, Mandi Bhaudhin, Jhelum and Sialkot; Nagri strain found in Faisalabad, Okara, Sahiwal and Pakpattan (Khan and Okeo, 2016).

Mastitis is the inflammation of mammary glands along with physical, chemical, pathological, microbiological changes and increased somatic cells, especially leukocytes in the milk (Fox, 2009). Mastitis is known as the most important and expensive disease of dairy animals. It is also the most prevalent production disease causing heavy financial penalty. So, any factor that harmfully deteriorates the milk quality and quantity from the goat is of great economic interest. Milk value is mostly pretentious by intra-mammary infections that become causes of mastitis and mostly mastitis in goats is mainly of bacterial species origin (Bergonier *et al.*, 2003).

Generally, mastitis occurs in two forms viz. clinical (overt) and sub-clinical (hidden). In the clinical mastitis, all the five fundamental signs of inflammation (pain, heat, swelling, redness and loss of milk production) are present while sub-clinical form is bereft of any noticeable sign of inflammation. About 75-80% mastitis is sub-clinical, characterized by a significantly increased leukocyte count in milk (Bradley, 2002). It exists for long time, adversely affects quantity and quality of milk and constitutes a reservoir of microorganisms due to their contagious nature for other animals within the herd (Schultz *et al.* 1978). Diagnosis of subclinical mastitis can be performed by various tests like California Mastitis Test (CMT) and Surf Field Mastitis Test (SFMT) under field conditions (Bachaya *et al.*, 2005; Muhammad *et al.*, 2010)

Mastitis in countries like Pakistan increases due to poor disease prevention and reporting system (Javed *et al.*, 2009). Goat mastitis has not been accorded due consideration by the scientific community of the country (Arshad *et al.*, 2006; Ali *et al.*, 2010) because considerable efforts and resources have been focused on the control of mastitis in cattle and buffaloes previously. Similarly, in Pakistan, the scientific research data on caprine mastitis in general and especially in various breeds of goat and recently documented strains of Beetal goat is meager or absent (Rizwana *et al.*, 2016). Notwithstanding the fact that Pakistan is the 3rd largest goat raising country of the world but at present a 'do nothing' policy *vis-à-vis* control of caprine mastitis is followed throughout the country. Keeping in view this scenario, the current study was planned to estimate mastitis distribution in different strains of Beetal goats, risk factors estimation and isolation of various strains of staphylococcus genus.

MATERIALS AND METHODS

Study locales and settings: For making the present research more demographic and diverse, three areas of Central Punjab (Faisalabad, Chiniot and Jhang) and three areas from Southern Punjab (Rajanpur, Bahawalpur and Muzaffargarh) were selected to conduct this study. Among these districts, three sites were selected based on home tracts of important strains of Beetal goat: Site-I Bahawalpur and Muzaffargarh districts for Beetal Makhicheeni, Site-II Rajanpur district for Beetal Nuqri and Site-III Faisalabad Division for Beetal Faisalabadi or Lyalpuri. The duration of research was about 18 months (January, 2016 to June, 2017).

Sample size and sampling: Sample size was calculated from the expected prevalence of about 20% (known disease status) with confidence interval of 95% and absolute desire precision 5%. Expected prevalence of 20% was deduced from the average prevalence of goat mastitis in three previous studies (Rashid *et al.*, 2017, Rizwana *et al.*, 2016 and Najeeb *et al.*, 2013). The sample size was estimated using the following equation (Thrusfield, 2007).

$$n = 1.96^2 P_{\text{exp}} (1 - P_{\text{exp}}) / d^2$$

Where:

n = required sample size

P_{exp} = expected prevalence

d = desired absolute precision

The number of samples thus calculated was subjected to the following equation for adjusted number to reach the maximum number of samples (Thrusfield, 2007):

$$n_{\text{adj}} = (N \times n) / (N + n)$$

Where:

N = total population

n = calculated sample size through formula

Following the proportionate sampling strategy and assuming the highest population (70%) of Beetal all over Punjab, 172 flock of Beetal from the selected districts and screened out respectively for the prevalence of mastitis in three strains of Beetal goat.

Epidemiological parameters: Epidemiological parameters including age, stage of lactation, parity, stage of pregnancy, length of lactation period, amount of milk, farming system, housing and management conditions, previous disease, vaccination, deworming, and treatment history were recorded on a predesigned questionnaire. Previous milk production and any mastitis control strategy (post milking teat dipping and dry period antibiotic therapy) were also recorded.

Diagnostic procedures: Field diagnosis of both clinical and sub-clinical mastitis was performed by complete physical examination of udder and Surf Field Mastitis Test (SFMT) in all milking animals of selected goat farms. SFMT score 1 was considered as negative whereas score 2 traces, 3 positive (weak degree of gel formation), 4 distinct positive and 5 for strong positive for sub-clinical mastitis (Schalm *et al.*, 1971). The SFMT score of 2 (traces) was considered healthy milk because goat milk has intra-cytoplasmic particles that resemble with somatic cells (Smith and Roguinsky, 1977).

Surf Field Mastitis Test (SFMT) scores: This test was performed as described previously (Schalm *et al.*, 1971; Muhammad *et al.*, 1995 and Muhammad *et al.*, 2010). The SFMT is a CMT-like test which has been demonstrated to be comparable to CMT in terms of sensitivity and specificity of mastitis detection (Badiuzzaman *et al.*, 2015). The reactions of these tests were interpreted as shown in Table 1 (Schalm *et al.*, 1971).

Microbiological examination of milk: Surf Field Mastitis Test (SFMT) positive milk samples were collected aseptically in sterile vials by recording the udder halves and SFMT score (Muhammad *et al.*, 2010). These milk samples were transferred to the Mastitis Research Laboratory, University of Agriculture, Faisalabad, for further microbiological examination. The microbiological assays were performed as described by National Mastitis Council Inc., USA (Anonymous, 1990). Briefly, for the isolation of mastitic pathogens, milk samples were aseptically cultured at 37°C for 24 hours on blood agar and MacConkey's agar. Gram-positive, catalase-positive, α and β hemolytic coccal isolates were presumptively identified as Staphylococci and subjected to coagulase test using rabbit plasma (at 4 hours) for grit of coagulation property of organism. For species identification of Staphylococci, biotyping of the isolates was conducted by using a commercial kit (api-STAPH; BioMerieux, France). Latex Slide Agglutination test using Staphytest plus kit (Oxoid, Ltd, Basingstoke, Hampshire, UK) was conducted to check the presence of protein A, clumping factor and certain polysaccharides that are found entirely in *S. aureus*.

Antibiotic sensitivity test: Disc diffusion method on Mueller-Hinton medium was used for the sake of antibiotic susceptibility profiles of the prevalent mastitic organisms following the guidelines (Anonymous, 2012).

Statistical analysis: The data obtained were analyzed by Chi Square test for the prevalence, incidence and other risk factors associated with mastitis (Steel *et al.*, 1997) in Table 3.

RESULTS AND DISCUSSION

Goat population of selected area: The total reported goat population in selected districts was 4,648,234 heads (Anonymous, 2006). According to Livestock Census, it was observed that Muzaffargarh is the biggest city of keeping goats among selected districts followed by Bahawalpur (1044722 head), Faisalabad (904918 head), Rajanpur (633524 head) and 503496 head for Jhang and Chiniot because after Livestock Census 2006, Chiniot was declared as independent district so that their goat population is divided equally.

Locales and Beetal goat strains distribution: A total of 3195 goats from 172 flocks were collected for screening of mastitis from 6 goat populated districts of Punjab (3 from central and 3 from southern Punjab) to make this study more diverse in form of climate and geography. Depending upon the population of goats in each district, random flocks were pulled out by using 'survey tool box' to remove biased sampling. By considering the reported goat population of selected districts (Anonymous, 2006), Beetal Faisalabadi 1673 goats were screened out for mastitis from 85 flocks from district Faisalabad. From Jhang, Beetal Faisalabadi 201 goats from 14 flocks were added in this study, from Chiniot, Beetal Faisalabadi 359 goats from 17 flocks, from Bahawalpur, 631 Beetal Makhicheeni goats from 32 flocks and from Rajanpur, 331 Beetal Nuqri goats from 24 flocks were selected.

Goat strains related distribution of samples: Proportionate sampling was conducted according to strains population in six selected districts. Out of 3195 goats sampled, 2233 (69.89%) were selected for Beetal Faisalabadi goats from the districts of Faisalabad, Chiniot, Jhang, Bahawalpur and Rajanpur. For Beetal Makhicheeni 631 goats (19.74%) were selected from districts Bahawalpur and Muzaffargarh while for Beetal Nuqri 331 goats (10.35%) from Rajanpur were screened out for mastitis (Table 2).

Table 1: Guidelines used for interpretation (scoring) of Surf Field Mastitis Test (SFMT)

Symbol (score)	Suggested meaning	Description of visible reaction
1	Negative	Mixture remains liquid.
2	Traces	A slight slime forms and was seen by tipping the paddle back and forth.
3	Weak Positive	A distinct slime but with no tendency towards gel formation.
4	Distinct Positive	The mixture thickens immediately with the gel formation.
5	Strong Positive	A gel forms with convex surface of mixture, which is so viscous that it adheres to the bottom of the receptacle of the test paddle.

Table 2: Prevalence of mastitis according to geographical area

Area	No. of goats	Mastitic goats	Prevalence (%)	p-value	Confidence interval (95%)
Faisalabad	1673	258	15.42	0.000	13.77-17.23
Jhang	201	28	13.93		09.82-19.40
Chiniot	359	71	19.78		15.99-24.21
Bahawalpur	425	80	18.82		15.39-22.81
Rajanpur	331	64	19.33		15.45-23.94
Muzaffar Garh	206	34	16.50		12.05-22.17
Total	3195	535	15.20		15.49-18.07
Strain wise prevalence					
Strain (Beetal)	No. of goats	Mastitic goats	Prevalence (%)	p-value	Confidence interval (95%)
Faisalabadi	2233	357	15.99	0.192	15.30-18.40
Makhi Cheeni	631	114	18.07		15.26-21.26
Nuqri	331	64	19.34		15.45-23.94
Total	3195	535	16.74		15.49-18.07

Table 3: Association of variables on mastitis

Variables	Class of Variable	Mastitis (%)	Healthy (%)	df	Chi Square value	P-Value
Color	Black& white	24(31.9)	3(0.4)	7	17.512	0.014*
	Brown & white	65(8.5)	2(0.3)			
	Brown black	6(0.8)	1(0.1)			
	Black	158(20.6)	2(0.3)			
	Brown Splashed	86(11.2)	0(0.0)			
	Brown	11(1.4)	0(0.0)			
	Blackish brown	125(16.3)	0(0.0)			
	White	64(8.3)	0(0.0)			
Education	Illiterate	284(37.0)	7(0.9)	2	8.17	0.014*
	Up to Middle	364(47.4)	1(0.1)			
	Matric & Above	112(14.6)	0(0.0)			
Body Condition Score (BCS)	2 (poor)	326(42.4)	0(0.0)	3	41.585	0.000**
	3 (Normal)	316(41.1)	0(0.0)			
	4 (Good)	117(15.2)	8(1.0)			
	5 (Excellent)	1(0.1)	0(0.0)			
Housing	Backyard Shed	226(29.4)	4(0.5)	2	19.705	0.000**
	Street	4(0.5)	1(0.1)			
	Open Area	530(69.0)	3(0.4)			
Farm hygiene	Very poor	258(33.6)	0(0.0)	3	8.283	0.041*
	Poor	124(16.1)	4(0.5)			
	Normal	362(47.1)	4(0.5)			
	Good	16(2.1)	0(0.0)			
Parity No.	1-2 Kidding	230(29.9)	6(0.8)	2	7.663	0.022*
	3-4Kidding	407(53.0)	2(0.3)			
	5 & > Kidding	123(16.0)	0(0.0)			
Age	1-3 Year	258(33.6)	6(0.8)	2	6.105	0.047*
	4-5 Year	402(52.3)	2(0.3)			
	>5 Year	100(13.0)	0(0.0)			
Ease of milking	Easy	95(51.4)	8(1.0)	1	7.322	0.007**
	Hard	365(47.5)	0(0.0)			
Milk leakage	Yes	33(4.3)	3(0.4)	1	19.482	0.000**
	No	727(94.7)	5(0.7)			
Blood in milk	Yes	88(11.5)	3(0.4)	1	5.093	0.024*
	No	672(87.5)	5(0.7)			
SFMT Score	*2 (Traces)	70(9.2)	0(0.0)	4	4.69.506	0.000**
	Weak positive	355(46.2)	0(0.0)			
	Distinct positive	258(33.6)	0(0.0)			
	Strong positive	77(10.0)	0(0.0)			
Milk Taste	Sweet	161(21.0)	8(1.0)	2	28.654	0.000**
	Bitter	75(9.8)	0(0.0)			
	Salty	524(68.2)	0(0.0)			
Milk yield	Decreased	756(98.4)	0(0.0)	1	509.305	0.000**
	Not decreased	4(0.5)	8(1.0)			

** Highly Significant * Significant.

Table 4: Antibiotic sensitivity test

Designation of vaccinal isolates	Antibiotics sensitivity profiles							
	AMC(10ug)	AMP(10ug)	L.S(109ug)	SXT(25ug)	NV(10ug)	EN(5ug)	AML(10ug)	OT(10ug)
<i>S. aureus</i> *(1)	28(S)	25(S)	22(S)	29(S)	25(S)	23(S)	32(S)	26(S)
<i>S. xylosum</i> *(2)	33(S)	31(S)	29(S)	33(S)	9(R)	32(S)	39(S)	28(S)
<i>S. hyicus</i> *(3)	29(S)	25(S)	28(S)	21(S)	25(S)	27(S)	32(S)	24(S)

AMC=Amoxicillin; AMP=Ampicillin; L.S=Lincomycin; SXT=Sulfamethoxazole+Trimethoprim; NV=Novobiocin; EN=Enrofloxacin; AML=Amoxicillin+clavulanic acid; OT=Oxytetracycline; S=Sensitive; R= Resistant; *Prevalent isolates.

Epidemiology of mastitis in different strains of Beetal:

The study inferred overall prevalence of 16.74% caprine mastitis in Beetal goat while strains of Beetal Faisalabadi, Beetal Makhicheeni, and Beetal Nuqri suffered 15.99%, 18.07%, and 19.34%, respectively. The overall strain-wise prevalence in Beetal goat was observed as highest 19.34% in Beetal Nuqri followed by 18.06% in Beetal Makhicheeni and 15.99% in Beetal Faisaladadi (Table 2). The major species of Staphylococcus genus found in caprine mastitis were *S. hyicus*, *S. xylosum*, *S. simulans*, *S. aureus*. Some unidentifiable Staphylococcal species were also observed.

Bacteriological study of caprine mastitic pathogens:

A total of 535 mastitis positive milk samples out of 3195 goats were screened out by using SFMT, with recording goat identity no., udder half marking and score of SFMT (Muhammad *et al.*, 2010). In this study, 357 mastitic

samples from Beetal Faisalabadi, 114 from Beetal Makhicheeni and 64 from Beetal Nuqri were studied for microbiological assays. Biotyping of isolates showed that coagulase negative Staphylococci are the main class of bacteria frequently observed. Furthermore, different types of bacteria were found in mastitic samples. *E. coli* from *Enterobacteriaceae*, Klebsiella, Proteus, Enterobacter from *Micrococcaceae*, *S. aureus*, Micrococcus spp and various spp. of Streptococcus, Bacillus, Corynebacterium and Pseudomonas spp. were also found.

Mastitis prevalence in small ruminants has been reported by Fox and Gay, 1993; Bergonier *et al.*, 2003; Contreras *et al.* 2003; Contreras *et al.* 2007; Ali *et al.* 2010; Megersa *et al.*, 2010; Islam *et al.*, 2011; Islam *et al.*, 2102; Bourabah *et al.*, 2013 and Rashid *et al.*, 2017. The current study's findings agree with outcomes of study reported by Megersa *et al.* (2010). Contradiction in prevalence outcome was found in studies conducted by

Contreras *et al.*, 2003 who revealed 9.4-47%, Rashid *et al.*, 2017 showed 21.68% in winter and 25.70% in summer respectively and Rizwan *et al.*, 2016 who reported average 18.29% prevalence of mastitis in different breeds of goat in Pakistan. The findings of this study is also in line with the results of Contreras *et al.* 2007 and Fox and Gay, 1993 who showed the range of prevalence from 5-30% and 7-40% respectively. The current finding is strongly differing with the results of Hall and Rycroft, 2007 showed 33%, Ali *et al.* 2010 revealed 30.6%, Islam *et al.*, 2011 reported 6%, Islam *et al.*, 2011 reported 4%, Aqib *et al.*, 2018 showed more than 50%, Bergronier *et al.*, 2003 showed 20-50% and Bourabah *et al.*, 2013 reported 33.9% prevalence. This difference could be due to environmental differences and differences in animal breeds and strains, fluctuation of immune response, housing and management system, all other farmers' and host's related determinants, with different diagnostic methods and different level of expertness for diagnosis as well as result interpretations.

Farm management and housing related determinants for mastitis: Current study revealed that type of housing, farm hygiene, age, stage of lactation, parity, color, body condition score, teat ends, SFMT score, milk taste, milk consistency, milk color and milk yield have significantly ($p < 0.005$) associated with mastitis. But it was observed that floor type, condition of floor, source of drinking water, feeding practices, use of mineral, general vaccination, deworming, milking practices, teat shape, teat symmetry, teat injury and supernumerary teat has non-significant ($P < 0.05$) effect on mastitis. This difference of non-significance could be due to environmental differences, differences in strains, fluctuation of immune response, housing and management system, all other farmers' as well as host's related determinants. Table 3 depicts that there is significant relation of color with mastitis prevalence in all three strains of Beetal. It was observed that color Black & White 31.9% followed by Black 20.6%, Blackish brown 16.3%, Brown Splashed 11.2%, Brown & White 8.5% while 0.8% in Brown Black. This information is reported for the first time in Pakistan and the reference availability for comparison might be difficult but its significant effect has been observed.

Varying level of education has a significant effect ($P < 0.05$) on mastitis as it was observed that as the level of education increases, marked decline was seen in mastitis prevalence that is due to better awareness about farm hygiene, sanitation, disinfectant, social as well as farmer meetings, watching TV for news in educated farmers community. The flock size has significant effect ($P < 0.05$), as the flock size of 21-40 goats has 46.5% followed by 1-20 goat head 28.8% and more than 40 head 23.7%. This finding is very close to the results about flock size of Megersa *et al.* 2010 and Rashid *et al.*, 2017. Body condition score has shown that there was highly significant ($P < 0.001$) difference presenting poor and good having 42.4% and 15.2%, respectively while excellent to be 0.1%. The findings were again in agreement with Megersa *et al.* 2010 and Rashid *et al.*, 2017. The calculated housing system has shown highly significant ($P < 0.05$) effect on mastitis as open area showed 69.0% followed by backyard shed 29.4% and street presenting

0.5% mastitis prevalence. Such variation speaks of cleaning as difficult or impossible for open area as there is no restriction for microbial contamination. The results were concordant with findings of Megersa *et al.* 2010 and Rashid *et al.*, 2017.

Farm hygiene has shown significant ($P < 0.05$) effect on mastitis as normal 47.1% followed by very poor 33.6%, poor 16.1% and good 2.1% was observed. Table 3 depicts that as the level of farm hygiene is attaining better score, marked decline was seen in mastitis. This finding is also strongly supported by Ameh *et al.* 2000 and Rashid *et al.*, 2017. It was observed that parity of 3-4 kidding has 53.0% followed by 1-2 kidding 29.9% while 5 and above kidding 16.4%. The chance of mastitis is directly proportional to parity as animal to face more numbers of infections depending upon the environmental conditions, milking practices and sanitations. The results are in line with findings of Ndegwa *et al.*, 2000, Moroni *et al.*, 2005 and McDougall *et al.*, 2002 showing strong correlation between occurrence of mastitis and the passage of time. As the age increases, there is greater risk of diseases likewise in case of mastitis. Age group 4-5 year showed 52.3% followed by 1-3 year 33.6% and more than 5 year 13.0%. Prevalence of mastitis has increased with increasing age in goats as reported by Ameh and Tari, 2000, McDougall *et al.*, 2002 and Moroni *et al.*, 2005 as observed in our study that occurrence of disease was more in 3-5 years of age in all strains of Beetal.

Ease of milking, milk leakage, blood in milk, SFMT score, milk taste and milk yield have very strong connection with the occurrence of mastitis and showing significant ($P < 0.05$) effect. Occurrence of mastitis was accompanied with marked hardness of teat during milking, milk leakage, blood in milk in very severe attack of mastitis, severity of SFMT score and decreased milk yield up to 20%. These findings are very close to the results presented by Ameh and Tari, 2000; McDougall *et al.*, 2002; Ndegwa *et al.*, 2000; Moroni *et al.* 2005; Athar *et al.*, 2007; Ahmad *et al.* 2008, Ali *et al.*, 2010; Megersa *et al.* 2010; Islam *et al.*, 2011 and Rashid *et al.*, 2017.

Antibiotic sensitivity test of prevalent isolates: In the present study, among the selected isolates, *S. aureus* and *S. hyicus* were sensitive to amoxicillin, ampicillin, lincomycin, sulfamethoxazole+Trimethoprim, novobiocin, enrofloxacin, amoxicillin+clavulanic acid and oxytetracycline while *S. xylosus* was sensitive to all except novobiocin. *S. simulans* was sensitive to lincomycin, sulfamethoxazole+Trimethoprim and enrofloxacin (Table 4). These study findings are comparable with that of Athar *et al.*, 2007; Ali *et al.*, 2010; Islam *et al.*, 2011 and Najeeb *et al.*, 2013 who also reported similar kind of trend in antibiotic susceptibility of *S. aureus* and Coagulase Negative Staphylococcal Spp. Contrarily, Chaudhry and Azam (1995) reported gentamicin to be the most effective *in vitro* antibiotic followed by chloramphenicol, kanamycin, oxytetracycline, cotrimoxazole, penicillin, doxycycline, ampicillin, and nystatin. Haphazard use of certain antibiotics might have conferred resistance in the present microorganisms and might lead to therapeutic failure. Therefore, use of antibiotics based on antibiotic susceptibility profile is highly warranted to achieving a better cure rate.

Conclusions: The study found over all 16.74% mastitis in different strains of Beetal goat with significantly variable mastitis distribution in studied areas. Type of housing, farm hygiene, age, stage of lactation, parity, color, body condition score, teat ends, SFMT score, milk taste, milk consistency, milk color and milk yield were found pertinent risk factors. The major staphylococcal species with decreasing order were *S. hyicus*, *S. xylosus*, *S. simulans*, *S. aureus* and unidentifiable staphylococcal species. The epidemiology of goat mastitis is fulminating with pathogenic involvement of bacteria requiring immediate attention.

Acknowledgments: The study was supported by Agriculture Innovation Program entitled “Cattle and goat value chain” and by GEF-UNEEP-ILRI FAnGR Asia Project “Development and Application of Decision Support Tools to Conserve and Sustainably Use Genetic Diversity in Indigenous Livestock & Wild Relatives”. Technology transfer project of Endowment Fund Secretariate, University of Agriculture, Faisalabad, “Dissemination of farmer’s friendly diagnostic techniques for the control of caprine mastitis in selected districts of Punjab-Pakistan” also provided traveling support as well as helped in farmer’s interaction, feedback and samples collection. The goat farmer’s cooperation was indispensable input for the accomplishment of this research. All contributions and supports are gratefully acknowledged.

Authors contribution: MIS and MS designed and conducted the study. MS helped in microbiological assay. MSK performed statistical analysis. GM and SUR observed and guided in research work. All authors contributed in drafting the manuscript.

REFERENCES

Ahmad T and Muhammad G, 2008. Evaluation of *staphylococcus aureus* and *streptococcus agalactiae* aluminium hydroxide adjuvanted mastitis vaccine in rabbits. Pak J Agri Sci 45:353-61.

Ali Z, Muhammad G, Ahmad T, et al., 2010. Prevalence of caprine sub-clinical mastitis, its etiological agents and their sensitivity to antibiotics in indigenous breeds of Kohat, Pakistan. Pak J Life Soc Sci 8:63-7.

Ameh JA and Tari IS, 1999. Observations on the prevalence of caprine mastitis in relation to predisposing factors in Maiduguri. Small Rum Res 35:1-5.

Anonymous, 1990. Microbiological Procedures for the Diagnosis of Bovine Udder Infection. National Mastitis Council, Inc., Arlington, Virginia, USA.

Anonymous, 2006. Livestock Census, Agricultural Census Organization, Statistics Division, Government of Pakistan, 3: 16-60.

Anonymous, 2012. Performance standards for antimicrobial susceptibility testing, 25th Informational Supplement, M-100-S-25.

Aqib AI, Nighat S, Ahmed R, et al., 2018. Drug susceptibility profile of *Staphylococcus aureus* isolated from mastitic milk of goats and risk factors associated with goat mastitis in Pakistan. Pak J Zool 51:307-15.

Arshad M, Muhammad G, Siddique M, et al., 2006. Staphylococcal mastitis in bovines and some properties of staphylococcal isolates. Pak Vet J 26:20.

Athar M, Muhammad G, Shakoor A, Saqib M, Ahmad R & Naureen A, 2007. A preliminary study on the effect of inactivated polyvalent mastitis vaccines on milk quantity and quality in dairy buffaloes. Pak Vet J 27:85-91.

Bachaya HA, Iqbal Z, Muhammad G, et al., 2005. Subclinical mastitis in buffaloes in Attock district of Punjab, Pakistan. Pak Vet J 25:134-6.

Badiuzzaman M, Samad MA, Siddiki SHMF, et al., 2015. Subclinical mastitis in lactating cows: comparison of four screening tests and effect of animal factors on its occurrence. Bangl J Vet Med 13:41-50.

Bergonier D, De CR, Rupp R, et al., 2003. Mastitis of dairy small ruminants. Vet Res 34:689-716.

Bourabah A, Ayad A, Boukraa L, et al., 2013. Prevalence and etiology of subclinical mastitis in goats of the Tiaret region, Algeria. Global Vet 11:604-8.

Bradley AJ, 2002. Bovine mastitis: an evolving disease. The Vet J 164:116-28.

Chaudhary TM and Azam MW, 1995. Sensitivity of various organisms isolated from clinical cases of mastitis in cattle, buffalo and goats. Pak J Livestock Poultr 1:74-6.

Contreras A, Luengo C, Sanchez A, et al., 2003. The role of intramammary pathogens in dairy goats. Livestock Prod Sci 79:273-83.

Contreras A, Sierra D, Sánchez A, et al., 2007. Mastitis in small ruminants. Small Rum Res 68:145-53.

DaMassa AJ, Wakenell PS and Brooks DL, 1992. Mycoplasmas of goats and sheep. J Vet Diag Inv 4:101-13.

Economic survey of Pakistan (2016-17). Economic Affairs Division, Islamabad, Government of Pakistan, Islamabad. pp:30.

Fox LK and Gay JM, 1993. Contagious mastitis. Vet Clin North Am, Food Anim Pract 9:475-87.

Fox LK, 2009. Prevalence, incidence and risk factors of heifer mastitis. Vet Micro 134:82-8.

Hall SM and Rycroft AN, 2007. Causative organisms and somatic cell counts in subclinical intramammary infections in milking goats in the UK. The Vet Rec 160:19-22.

Iqbal A, Khan BB, Tariq M et al., 2008. Goat A potential dairy animal Present and future prospects. Pak J Agri Sci 45:227-30.

Islam A, Samad A and Rahman A, 2012. Prevalence of subclinical caprine mastitis in Bangladesh based on parallel interpretation of three screening tests. Int J Anim Vet Adv 4:225-8.

Islam MR, Ahamed MS, Alam MS, et al., 2012. Identification and antibiotic sensitivity of the causative organisms of subclinical mastitis in sheep and goats. Pak Vet J 32:179-82.

Javaid SB, Gadahi JA, Khaskeli M, et al., 2009. Physical and chemical quality of market milk sold at Tandojam, Pakistan. Pak Vet J 29:27-31.

Khan MS and Okeyo AM, 2016. Handbook of Judging and Selection in Beetal Goats GEF-UNEP-ILRI FAnGR Asia Project, Academic Press, University of Agriculture Faisalabad, Pakistan pp:2-3.

Khan MS, Khan MA and Mahmood S, 2008. Genetic resources and diversity in Pakistani goats. Int J Agric Biol 10:227-31.

McDougall S, Pankey W, Delaney C, et al., 2002. Prevalence and incidence of subclinical mastitis in goats and dairy ewes in Vermont, USA. Small Rum Res 46:115-21.

Megersa B, Tadesse C, Abunna F, et al., 2010. Occurrence of mastitis and associated risk factors in lactating goats under pastoral management in Borana, Southern Ethiopia. Trop Anim Health Prod 42:1249-55.

Moroni P, Pisoni G, Vimercati C, et al., 2005. Characterization of *Staphylococcus aureus* isolated from chronically infected dairy goats. J Dairy Sci 88:3500-9.

Muhammad G, Athar M, Shakoor A, et al., 1995. Surf Field Mastitis Test: An inexpensive new tool for evaluation of wholesomeness of fresh milk. Pak J Food Sci 5:91-3.

Muhammad G, Naureen A, Asi MN, et al., 2010. Evaluation of a 3% surf solution (surf field mastitis test) for the diagnosis of subclinical bovine and bubaline mastitis. Trop Anim Health Prod 42:457-64.

Najeeb MF, Anjum AA, Ahmad MUD, et al., 2013. Bacterial etiology of subclinical mastitis in dairy goats and multiple drug resistance of the isolates. J Anim Plant Sci 23:1541-4.

Ndegwa EN, Mulei CM and Munyua JM, 2000. The prevalence of subclinical mastitis in dairy goats in Kenya. J South Afr Vet Assoc 71:25-7.

Rashid M, Saleem MI, Deeba F, et al., 2017. Effect of season on occurrence of caprine mastitis in beetal in faisalabad premises. Mat Sci Med 1:19-21.

Rizwana M, Durranib AZ, Ijaz M, et al., 2016. Clinio-bacteriological investigation of sub-clinical and clinical mastitis in dairy goats. Veterinaria 4:4-6.

Schalm OW, Carroll EJ and Jain NC, 1971. Bovine mastitis. 1971 Ed. Lea and Febiger, Philadelphia, USA.

Schultz LH, Broom RW, Jasper DE, et al., 1978. Current concepts of bovine mastitis. In The National Mastitis Council pp:1-16.

Smith MC and Roguinsky M, 1977. Mastitis and other diseases of the goat's udder. J American Vet Med Assoc 171:1241-8.

Steel RGD, Torrie JH and Dickey DA, 1997. Principles and procedures of statistics: a biometrical approach, 3rd Ed., McGraw-Hill: New York.

Thrusfield M, 2007. Veterinary Epidemiology. Blackwell Science, Oxford, Chap 13:228-42.