



RESEARCH ARTICLE

Protective Effect of Yeast Sludge and Whey Powder against Ochratoxicosis in Broiler Chicks

Huma Mujahid^{1*}, Abu Saeed Hashmi², Muhammad Zargham Khan³, Muhammad Tayyab¹ and Wasim Shehzad¹

¹Institute of Biochemistry and Biotechnology, University of Veterinary and Animal Sciences, Lahore; ²Ripha College of Veterinary Sciences, Lahore; ³Department of Pathology, University of Agriculture, Faisalabad

*Corresponding author: huma.mujahid@uvas.edu.pk

ARTICLE HISTORY (19-119)

Received: March 17, 2019
Revised: April 19, 2019
Accepted: May 09, 2019
Published online: May 25, 2019

Key words:

Broilers
Ochratoxin A
Toxicity
Whey
Yeast sludge

ABSTRACT

The aim of the present study was to determine the detoxification potential of the food industry by-products such as yeast sludge (YS) and whey powder (WP) against harmful effects of ochratoxin A (OTA) in broilers. One day old broilers chicks (n=1250) were randomly divided into five groups replicated five times with each replicate having 50 birds. The experimental feed in different groups was as; group A (basal feed), group B (200ppb OTA), group C (200ppb OTA and 0.2% Dried YS), group D (200ppb OTA and 0.2% WP) and group E (200ppb OTA and 0.2% protomyc). OTA adversely affected body weight gain, feed consumption and feed conversion ratio (FCR) of broiler chicks. Haematobiochemical parameters such as alanine amino transferase (ALT), aspartate amino transferase (AST), and creatinine levels raised by OTA feeding were significantly ($P>0.05$) reduced in YS and WP supplemented group. Residues of OTA were detected in all the tissues studied, with highest levels observed in kidneys, YS and WP significantly reduced the tissue residues of OTA. In conclusion, present study suggested that addition of YS and WP in broilers feed reduce the harmful effects of OTA in broiler chicks as efficiently as protomyc a commercial mycotoxin binder.

©2019 PVJ. All rights reserved

To Cite This Article: Mujahid H, Hashmi AS, Khan MZ, Tayyab M and Shehzad W, 2019. Protective effect of yeast sludge and whey powder against ochratoxicosis in broiler chicks. Pak Vet J, 39(4): 588-592. <http://dx.doi.org/10.29261/pakvetj/2019.077>

INTRODUCTION

Ochratoxins are mycotoxins produced by toxigenic strains of *Aspergillus ochraceus*, *Aspergillus westerdijkiae*, *Aspergillus niger* and some spp. of *Penicillium*, it has three main types, i.e., Ochratoxin A (OTA), ochratoxin B (OTB) and Ochratoxin C (OTC). Of these, OTA has potent immunotoxic, nephrotoxic, mutagenic and teratogenic effects (Paola and Marco, 2015). The presence of OTA in poultry feedstuffs appears to put a severe hazard for the native poultry industry. It also induces alterations in the biochemical parameters (reduction in, hematocrit, hemoglobin, blood cells, serum total proteins; whereas the uric acid, creatinine, liver enzymes were increased). It disturbs protein synthesis, carbohydrate and lipid metabolism blood coagulation and hormones. Ochratoxin A is also characterized as severe immunosuppressive in avian species (Saleemi *et al.*, 2017; Elaroussi *et al.*, 2008).

Due to prevalence of high level toxicity of OTA, it is imperative to find procedures for elimination of OTA from poultry food and feedstuffs (Amézqueta *et al.*,

2009). An effective approach for detoxification of mycotoxin in animal feedstuff is the usage of nutritionally inert substances, which can minimize the level of toxicity. Toxin binders, such as aluminosilicate, yeast, lactic acid producing bacteria and detoxifying agents such as vitamin E, are used in the feed to reduce the harmful effects of these mycotoxins on poultry birds (Khan *et al.*, 2010; 2014a; Chen *et al.*, 2016).

Distillery yeast sludge (DYS) a by-product of alcohol production industry contains high amount of glucomannans (Sharif *et al.*, 2012), which can adsorb different mycotoxins. An improvement in the blood biochemistry parameters and productive performance of broilers has been reported by the supplementation of DYS in mycotoxin contaminated feed (Mujahid *et al.*, 2012; Hashmi *et al.*, 2006; Khatoon *et al.*, 2017).

Dried concentrated whey is a by-product of cheese making industry, it enhances the immunity, improve survival rates, and stimulate the growth of beneficial intestinal bacteria in broilers because it is a natural prebiotic. Greater abundance of *L. salivarius* with improved feed conversion ratio (FCR) has been reported

by the supplementation of whey in Broiler diet (Pineda-Quirogaa *et al.*, 2017). Anti-ochratoxin effect of whey can be attributed to its capacity to increase the *Lactobacillus* count in the gastrointestinal tract of the poultry (Hameed *et al.*, 2017b; Zarei *et al.*, 2018) Moreover Mansour *et al.* (2011) and (2015) proved that whey successfully neutralizes the drastic toxic effects of OTA in *Oreochromis niloticus*. Studies conducted to investigate the effect of whey on ochratoxicosis in broilers are not available. The objective of this paper is to discuss the changes on broiler performance and biochemical parameters associated with ochratoxicosis induced in broiler chicks by feeding them with a diet containing known concentration of OTA at level of 200ppb. In addition, to determine the protective effect of YS and WP against harmful effects of OTA on broiler performance, biochemical parameters and tissue residues.

MATERIALS AND METHODS

Collection of industrial waste: Yeast sludge (YS) was collected from Shukkar Gunj Sugar Mills, Jhang and whey powder (WP) was purchased from local market. Protomyc (yeast cells and bentonite) Biorigin, Brazil was purchased from local distributors.

Production of OTA: OTA was produced from *Aspergillus ochraceus* (CECT 2948, Culture Collection - Center, University De Valencia, Spain) by solid state fermentation on wheat grain Trenk *et al.* (1971). Briefly, 50g of wheat grains were soaked in 50 mL of distilled water for 2 hours in a 500 mL Erlenmeyer flask. The flask was autoclaved, inoculated with *A. ochraceus* spores and incubated for two weeks at 28°C in dark and shaken once daily. OTA was extracted from the fermented wheat by solvent extraction in acetonitrile-water and quantified by HPLC (Bayman *et al.*, 2002).

Biological trial: A Biological trial of 35 days duration was conducted at Hi-Tech Research and Development Centre, Lahore. An approval for the study was granted by ethical review committee, University of Veterinary and Animal sciences, Lahore.

Birds and experimentation: A total of 1250 one day old broiler chicks of Arbreaker breed were divided randomly into five groups (A-E) replicated five times in such a way that each replicate contained 50 birds. Five diets formulated were A (basal poultry feed, 22% protein contents), B (OTA 200ppb), C (OTA 200ppb and 0.2% dried YS), D (OTA 200ppb and 0.2% WP) and E (200ppb OTA and 0.2% protomyc). Experiment was conducted under completely randomized design (CRD). The Broilers were assigned these rations *ad libitum* for 35 days. During the trial weekly weight gain (g) and feed consumption (g) was recorded. At the end of experimental trial feed conversion ratio (FCR) and mortality rate was calculated Khan *et al.* (2017).

Sample collection: At end of trail two birds were selected randomly per replicate. Blood (2 mL) was collected per bird by syringe from wing vein. Serum was separated and stored at -20°C. The thigh muscles, liver, kidney and heart samples of slaughtered birds were taken in sterilized plastic containers, labelled and stored at -20°C.

Estimation of serum Biochemical parameters: Estimation of serum total protein, albumin, creatinine, Activity of ALT and AST was carried out by using commercially available kits (Human, Wiesbaden, Germany).

Hemagglutination inhibition (HI) test against Newcastle disease virus (NDV): The Humoral Immune response was evaluated by HI test against NDV (Anon, 1971; Mahmood *et al.*, 2018). In brief the hemagglutination Inhibition pattern was detected of the highest dilution of the virus giving complete HA pattern. The values of the last dilutions which caused total inhibition of hemagglutination were calculated as the logarithm base 2.

Extraction of OTA from tissues: Two millilitres of serum sample was mixed with 2.5mL of phosphoric acid and twenty gram of tissues samples (muscles, kidneys, heart and livers) were homogenized with 7.5 mL of 1 M phosphoric acid in a homogenizer. Two millilitres of phosphoric acid treated samples were extracted thrice with Ethyl acetate (5 mL), and centrifuged at 350 g for 5 min. The organic phase concentrated and extracted with 2 mL of 0.5 M NaHCO₃. The aqueous extract was purified further by immune affinity column (OchraTest WB column, Vicam, USA) (Bozzo *et al.*, 2008).

Estimation of OTA in tissue samples: The purified extract was analyzed by high performance liquid chromatography (HPLC) Agilent 1260 Infinity II system. The mobile phase was acetonitrile: water: acetic acid (99:99:2) (Biro *et al.*, 2002).

Statistical analysis: The data was analyzed statistically by One way ANOVA and significant differences among means were compared using the Duncan Multiple Range Test using SPSS 16 software.

RESULTS

Body weights of OTA treated group was slightly but not significantly decreased (Table 1) from the control (A) at any of the 5 weeks, weight gain in group D fed on WP at week 2 and 4, was significantly higher than Group B. Group E treated with protomyc showed significantly higher weight gain ($P < 0.05$) than other groups at week 4 and 5 ($P < 0.05$). On day35, the ameliorating effect of protomyc was clear: though group D was non significantly different at 1, 3 and 5 weeks but the body weights of group D were significantly greater ($P < 0.05$) at weeks 2 and 4 when compared with group B.

As evident from Table 2, no significant difference was observed in feed consumption in all groups. The quantity of feed consumed in groups D and E fed on WP and protomyc showed significantly higher feed consumption in week 2, the feed consumption of YS, WP and protomyc was slightly (but not significantly) improved.

No significant differences between groups were found when FCR was calculated (Table 3). Only in week 3 was the FCR of group E increased ($P < 0.05$). At week 5, the FCR of YS, WP and protomyc treated group was slightly, but not significantly improved than group B.

Table 1: Effects of Feeding OTA, YS, WP and protemyc supplementation on weight gain of broiler chicks (Means±SD)

Treatment	Weeks					
	Bodyweight (g)					
	1 st Day	1	2	3	4	5
A (basal feed)	42±1.5	171.6±5.7 ^{ab}	372.6±12.6 ^{ab}	754.8±34.3 ^a	1003±98.6 ^a	1382.4±74.2 ^{ab}
B (200ppb OTA)	42±1.5	166.2±17.4 ^{ab}	338.8±9.4 ^a	747.8±39.3 ^a	993±56.05 ^a	1361.8±37.0 ^a
C (200ppb OTA+YS)	42±1.5	172.9±5.1 ^b	395.6±34.3 ^{ab}	765.5±69.0 ^a	1111.4±86.9 ^{ab}	1428.6±4.1 ^{ab}
D (200ppb OTA+ WP)	42±1.5	161.8±6.7 ^{ab}	410.8±39.5 ^b	816.8±70.1 ^a	1166.2±71.9 ^b	1461±54.6 ^{ab}
E (200ppb OTA+ protemyc)	42±1.5	159.3±9.3 ^a	394.6±30.9 ^{ab}	770.6±63.9 ^a	1191.8±104.6 ^b	1476.2±84.9 ^b

Different superscripts on means in a column show significant difference among groups (P<0.05); OTA (Ochratoxin A); YS (yeast sludge); WP (whey powder).

Table 2: Effects of Feeding OTA, YS, WP and protemyc supplementation on Feed Consumption of broiler chicks (Means±SD)

Treatment	Weeks				
	Feed Consumption (g)				
	1	2	3	4	5
A (basal feed)	161.6±11.8 ^a	445.8±15.1 ^{ab}	694.5±15.6 ^a	1669.2±42.0 ^a	2312±73.9 ^a
B (200ppb OTA)	156.1±14.3 ^a	396.9±38.8 ^a	690.4±6.1 ^a	1668.4±167.5 ^a	2303.7±178.8 ^a
C (200ppb OTA+YS)	167.6±10.46 ^a	450.7±35.3 ^{ab}	701.0±71.7 ^a	1747.8±157.1 ^a	2391.8±175.0 ^a
D (200ppb OTA+ WP)	162.8±12.2 ^a	462±37.1 ^b	698.9±43.3 ^a	1767.6±116.8 ^a	2416.3±183.2 ^a
E (200ppb OTA+ protemyc)	166.2±11.9 ^a	469.96±26.6 ^b	735.6±93.5 ^a	1769.8±146.9 ^a	2460.4±147.16 ^a

Different superscripts on means in a column show significant difference among groups (P<0.05); OTA (Ochratoxin A); YS (yeast sludge); WP (whey powder).

Table 3: Effects of Feeding OTA, YS, WP and protemyc supplementation on FCR of broiler chicks (Means±SD)

Treatment	Weeks				
	FCR				
	1	2	3	4	5
A (basal feed)	0.941±0 ^a	1.19±0 ^{ab}	0.92±0 ^a	1.67±0.1 ^a	1.67±0 ^a
B (200ppb OTA)	0.933±0 ^a	1.13±0.1 ^a	0.93±0 ^a	1.56±0.1 ^a	1.68±0 ^a
C (200ppb OTA+YS)	0.966±0 ^a	1.14±0 ^a	0.91±0 ^a	1.57±0 ^a	1.67±0 ^a
D (200ppb OTA+ WP)	1.007±0 ^a	1.19±0 ^{ab}	0.86±0 ^a	1.51±0.1 ^a	1.65±0 ^a
E (200ppb OTA+ protemyc)	1.024±0 ^a	1.33±0.1 ^b	0.95±0.1 ^a	1.48±0 ^a	1.67±0 ^a

Different superscripts on means in a column show significant difference among groups (P<0.05); OTA (Ochratoxin A); YS (yeast sludge); WP (whey powder).

Table 4: Effects of Feeding OTA, YS, WP and protemyc supplementation on serum Biochemical parameters of broiler chicks (Means±SD)

Treatments	Serum Total Protein (g/dL)	Serum Albumin (g/dL)	Activity of ALT (IU/dL)	Activity of AST (IU/dL)	Seum Creatinine Levels (g/dL)	HI Titer against NDV
A (basal feed)	3.02±0.1 ^a	1.74±0.1 ^d	7.1±0.7 ^a	159±1.5 ^a	0.92±0.1 ^{ab}	5.4±0.8 ^{ab}
B (200ppb OTA)	2.88±0.3 ^a	1±0.16 ^a	42±2.2 ^e	401.6±2.0 ^e	1.24±0.1 ^c	4.46±0.5 ^a
C (200ppb OTA+ YS)	3.64±0.3 ^b	1.5±0.21 ^c	32.6±1.8 ^d	373±10.5 ^d	0.78±0 ^a	5.6±1.1 ^{ab}
D (200ppb OTA+ WP)	3.72±0.1 ^b	1.16±0.11 ^{ab}	18±1.5 ^c	251.6±6.3 ^c	1.10±0.1 ^{bc}	6.2±0.4 ^{ab}
E (200ppb OTA+ protemyc)	4.78±0.7 ^c	1.34±0.15 ^{bc}	12.8±1.4 ^b	211.8±1.0 ^b	1.06±0.2 ^{bc}	6.4±2.1 ^b

Different superscripts on means in a column show significant difference among groups (P<0.05); OTA (Ochratoxin A); YS (yeast sludge); WP (whey powder).

Table 5: Effects of Feeding OTA, YS, WP and protemyc supplementation on Tissue residues of Ochratoxin A in broiler chicks (Means±SD)

Treatments	Tissue Levels of ochratoxin A µg/Kg				
	Heart	Thigh muscle	Liver	kidney	Serum
A (basal feed)	0 ^a	0 ^a	0 ^a	0 ^a	0 ^a
B (200ppb OTA)	0.073±0 ^b	0.016±0 ^c	0.93±0.3 ^b	1.14±0.2 ^b	3.4±0.2 ^d
C (200ppb OTA+YS)	0.068±0 ^b	0.004±0 ^{ab}	0.03±0 ^a	0.08±0 ^a	1.4±0 ^c
D (200ppb OTA+WP)	0.024±0 ^a	0.003±0 ^{ab}	0.0022±0 ^a	0.06±0 ^a	0.34±0 ^b
E (200ppb OTA+ protemyc)	0.024±0 ^a	0.006±0 ^b	0.07±0 ^a	0.08±0 ^a	0.53±0.1 ^b

Different superscripts on means in a column show significant difference among groups (P<0.05); OTA (Ochratoxin A); YS (yeast sludge); WP (whey powder).

Highest mortality (9.2%) was recorded in birds received diet contaminated with 200ppb OTA followed by Group E (7.6%) fed with protemyc. The Mortality rate in Group C supplemented with YS was 5.6%. The lowest mortality (4%) was recorded in group D treated with WP in the presence of OTA. Supplementing broiler diets with YS, WP and protemyc in presence of OTA alleviate the toxic effect of OTA on broiler mortality.

The results in Table 4 indicate that serum total protein was slightly reduced in OTA treated group but serum albumin levels were significantly reduced (P<0.05) in OTA treated group as compared to control. Significant increase in serum total protein and albumin was observed in YS, WP and protemyc treated groups. Activity of ALT

and AST was significantly increased (P<0.05) in OTA treated group as compared to control group. YS, WP and protemyc significantly improved the level of these enzymes. Serum creatinine level was also significantly increased (P<0.05) in OTA treated group as compared to control group and significant improvement was observed in YS, WP and protemyc supplemented groups. HI titre was slightly but not significantly (P<0.05) reduced in OTA treated group as compared to control group and it was significantly increased by protemyc treatment.

Toxin residues were detected in all the tissue samples of group B (200ppb OTA fed group), and the concentration of OTA in serum, liver and kidney was significantly higher in group B (P<0.05) as compared to

other groups as shown in Table 5. Heart and muscle contain very low level of OTA, WP and protemyc significantly reduced the tissue residues in these organs. Tissue level of ochratoxin A was found to be Serum>Kidney>Liver>Heart>Muscle. Whereas in the groups fed with diet supplemented with YS, WP and protemyc a considerable decrease in serum and tissue levels was detected.

DISCUSSION

Results obtained in the present study showed that body weight, feed consumed and FCR was slightly but non significantly impaired by OTA at a level of 200ppb as compared to the control group. At the first week of experiment, mortality was observed in all groups due to stress of travelling, weak immune system and adjustment to the new environment, thereafter mortality was increased in OTA group as compared with the other groups. YS, WP and protemyc improved the performance parameters in OTA treated broiler chicks. The slight alteration in body weight, FC and FCR due to OTA in the present study was in accordance with several previous investigations using dietary OTA inclusion rates of 0.1, 0.5, 1.0, 3.0, 5.0 and 10.0 mg/Kg feed (Zahoor *et al.*, 2010). Dönmez *et al.*, 2012 indicated that addition of yeast sludge to broiler diets providing the partial protection against the harmful effects of OTA. Similar results of L-carnitine are also reported by Bhatti *et al.* (2018). The addition of whey as a prebiotic to diets may influence broiler weight gain. The improvement in growth rate of birds fed with diets containing the tested prebiotic shows that the use of these products is a feasible alternative to antibiotics and antimycotoxins used as growth promoters (Nagarachi *et al.*, 2007).

Anti-ochratoxin effect of whey can be attributed to its capacity to increase the *Lactobacillus* count in the gastrointestinal tract of the poultry and *Lactobacillus* strains were conducive to detoxification of OTA when added to a feed mixture for chickens (Śliżewska *et al.*, 2014).

Present results are in agreement with those reported by Mansour *et al.* (2011) and (2015) where in *Oreochromis niloticus* addition of whey to fish diets with OTA improved the productive performance, histopathological alterations and Biochemical parameters.

OTA administrations to broilers also altered different serum biochemical parameters adversely. Activity of ALT, AST and creatinine levels were significantly increased in OTA fed group as compared to all other groups. Feeding of OTA increased the serum levels of ALT (Khan *et al.*, 2017), urea and creatinine. Zahoor *et al.*, (2010) also reported that Serum levels of total proteins and albumin were significantly decreased in the groups fed OTA.

The findings of present study revealed that supplementation of feed with YS and WP reduce the OTA levels in serum, kidneys and liver significantly. We previously reported that level of OTA is highest in kidney as compared to liver in chicks fed with 500 ppb OTA and 2% YS effectively reduced the level of ochratoxin A in tissues (Mujahid *et al.*, 2012). Supplementation of WP also reduces the tissue residues of OTA indicating the gastrointestinal removal of OTA probably due to increase

in *Lactobacillus* count in gastrointestinal tract of poultry. Lactic acid bacteria are actively involved in the adsorption of OTA (Śliżewska *et al.*, 2014).

Conclusions: The present study described performance parameters and biochemical alterations in OTA fed broilers. Improvement in FCR, serum biochemical changes, tissue residues and decreased mortality in YS and WP supplemented groups suggested a decrease in severity of the ochratoxicosis with supplementation of these industrial by products in feed.

Acknowledgments: I am thankful to Dr Muhammad Athar from Hi-Tech feeds PVT Limited for providing us the facility to run the biological trial.

Authors contribution: HM: Designed and performed the experiments, analysed the data and wrote the manuscript. ASH proposed the idea of the research work, designed the study and also help in planning and arrangement of trials. MZK provided the fungal strain and methodology for toxin production and also helps to prepare the final manuscript. MT worked out almost all of the technical details, and analyzed the data obtained from the experiments. WS provided the chemicals and raw materials required in the study, contributed to the design and implementation of the research, to the analysis of the results and to the writing of the manuscript. All the authors read and approved the manuscript.

REFERENCES

- Amézqueta S, Schorr-Galindo S, Murillo-Arbizu M, *et al.*, 2009. OTA-producing fungi in foodstuffs: a review. *Food Control* 4:326-33.
- Anon N, 1971. Methods for examining poultry biologics and for identifying avian pathogens. National Academy of Sciences, Washington DC pp:270-94.
- Bayman P, Baker JL, Doster MA, *et al.*, 2002. Ochratoxin production by the *Aspergillus ochraceus* group and *Aspergillus alliaceus*. *Appl Environ Microbiol* 68:2326-9.
- Bhatti SA, Khan MZ, Hassan ZU, *et al.*, 2018. Dietary L-carnitine and vitamin-E; a strategy to combat ochratoxin-A induced immunosuppression. *Toxicon* 153:62-71.
- Biro K, Solti L, Barna-Vetro I, *et al.*, 2002. Tissue distribution of ochratoxin A as determined by HPLC and ELISA and histopathological effects in chickens. *Avian Pathol* 31:141-8.
- Bozzo G, Ceci E, Bonerba E, *et al.*, 2008. Ochratoxin A in laying hens: high-performance liquid chromatography detection and cytological and histological analysis of target tissues. *J Appl Polym Res* 17:151-6.
- Chen L, Jiang T and Li X, 2016. Immunomodulatory activity of β -glucan and mannan-oligosaccharides from *Saccharomyces cerevisiae* on broiler chickens challenged with feed-borne *Aspergillus fumigatus*. *Pak Vet J* 36:297-301.
- Dönmez N, Dönmez HH, Keskin E, *et al.*, 2012. Effects of aflatoxin on some haematological parameters and protective effectiveness of esterified glucomannan in Merino rams. *Sci World J* 87:125-31.
- Elaroussi MA, Mohamed FR, Elgendy MS, *et al.*, 2008. Ochratoxicosis in broiler chickens: Functional and histological changes in target organs. *Int J Poult Sci* 7:414-22.
- Hameed MR, Khan MZ, Saleemi MK, *et al.*, 2017a. Study of ochratoxin A (OTA) induced oxidative stress markers in broiler chicks. *Toxin Rev* 36:270-4.
- Hashmi I, Pasha TN, Jabbar MA, *et al.*, 2006. Study of adsorption potential of yeast sludge against aflatoxins in broiler chicks. *J Anim Plant Sci* 16:12-4.
- Khan WA, Khan MZ and Khan A, 2010. Pathological effects of aflatoxins and their amelioration by vitamin E in White Leghorn layers. *Pak Vet J* 30:155-62.
- Khan WA, Khan MZ and Khan A, 2014a. Potential for amelioration of aflatoxin B1-induced immunotoxic effects in progeny of white

- leghorn breeder hens co-exposed to vitamin E. J Immunotoxicol 11:116-25.
- Khan A, Aalim MM, Khan MZ, et al., 2017. Does distillery yeast sludge ameliorate moldy feed toxic effects in White Leghorn hens? Toxin Rev 36:275-81.
- Khatoon A, Khan MZ, Abidin ZU, et al., 2017. Mitigation potential of distillery sludge (DS) against ochratoxin A induced immunological alterations in broiler chicks. World Mycotoxin J 10:255-62.
- Mahmood MS, Amir HVV, Abbas RZ, et al., 2018. Evaluation of antiviral activity of *Azadirachta indica* (Neem) bark extract against Newcastle Disease Virus. Pak Vet J 38:25-8.
- Mansour TA, Safinaz G, Mohamed S, et al., 2011. Ameliorate the drastic effect of ochratoxin a by using yeast and whey in cultured *Oreochromis niloticus* in Egypt. Life Sci J 8:68-81.
- Mansour TA, Omar AE, Soliman KM, et al., 2015. The antagonistic effect of whey on ochratoxin a toxicity on the growth performance, feed utilization, liver and kidney functions of Nile tilapia (*Oreochromis niloticus*). Middle East J App Sci 5:176-83.
- Nagharchi MM, Jouybari GM, Rezaei-pour V, et al., 2010. The effects of fermented and dried whey powder on performance and nutrient digestibility in broilers. Analele IBNA 26:76-82.
- Paola B and Marco CL, 2015. OTA-Grapes: A mechanistic model to predict ochratoxin a risk in grapes, a step beyond the systems approach. Toxins 7:3012-29.
- Pineda-Quiroga C, Atxaerandioa R, Zubiria I, et al., 2017. Productive performance and cecal microbial counts of floor housed laying hens supplemented with dry whey powder alone or combined with *Pediococcus acidilactici* in the late phase of production. Livest Sci 195:9-12.
- Saleemi MK, Khan MZ, Khan A, et al., 2017. Study of fungi and their toxigenic potential isolated from wheat and wheat-bran. Toxin Rev 36:80-8.
- Sharif M, Shahzad MA, Rehman S, et al., 2012. Nutritional evaluation of distillery sludge and its effect as a substitute of canola meal on performance of broiler chickens. Asian-Australas J Anim Sci 25:401-9.
- Śliżewska K and Piotrowska M, 2014. Reduction of ochratoxin a in chicken feed using probiotic. Annals Agric Env Med 21:676-80.
- Trenk HL, Butz ME and Chu FS, 1971. Production of ochratoxins in different cereal products by *Aspergillus ochraceus*. Appl Microbiol 21:1032-5.
- Zahoor-ul-Hassan, Khan MZ, Khan A, et al., 2010. Pathological responses of White Leghorn breeder hens kept on ochratoxin a contaminated feed. Pak Vet J 30:118-23.
- Zarei A, Lavvaf A and Motlagh MM, 2018. Effects of probiotic and whey powder supplementation on growth performance, microflora population and ileum morphology in broilers. J App Anim Res 46:840-4.